



KAISER PERMANENTE®

DOCUMENT NUMBER: SBMC-PPP-0032
DOCUMENT TITLE: Drawing Blood from Patients with an IV
DOCUMENT NOTES:

LOCATION: SBMC-rel	VERSION: 02
DOC TYPE: SBMC PPP	STATUS: Release

EFFECTIVE DATE: 10 May 2018	NEXT REVIEW DATE: 03 Apr 2022
RELEASE DATE: 10 May 2018	EXPIRATION DATE:

AUTHOR:	PREVIOUS NUMBER: SCPMG-PPP-0110
OWNER: SBMC Preatalytic Mgr	CHANGE NUMBER:

Drawing Blood from Patients with an IV

Introduction Drawing blood from an IV arm has the potential risk for erroneous and misleading test results. Whenever possible, blood should be drawn from the opposite arm when an IV fluid is being administered into a patient's arm. Blood should not be drawn from that limb unless there is no alternative and the tests are critical to the care of the patient, as determined by the provider.

Scope This procedure is intended for use by any person performing phlebotomy.

- Policy**
- If a patient has an IV in one arm, the Phlebotomist must obtain the blood specimen from the arm without the IV.
 - If a patient has an IV in both arms or if the arm with an IV is the only site, the Phlebotomist must follow the procedure below.
 - Collection above the IV is not recommended and should only be attempted when other alternatives are exhausted.
 - Acceptable sites for draw are the antecubital vein, hand and wrist. Feet and ankles sites may only be used after consulting with the patient's provider and obtaining a written order, if applicable per local policies.
 - Do not draw specimens when a patient is being transfused. The transfusion may be paused for 5 minutes to have blood drawn.
-

Procedure Follow the steps below to draw a blood specimen from a patient that has an IV and there is no other alternative for specimen collection.

Step	Action
1	Identify the patient using two patient identifiers.

Continued on next page

Drawing Blood from Patients with an IV, Continued**Procedure** (continued)

Step	Action				
2	<p>Ask the patient's nurse to turn off the IV for 5 minutes.</p> <p>NOTE: If the drip is an anticoagulant and the draw is above the IV, ask the patient's nurse to turn off the IV for 10 minutes.</p> <table border="1"> <tbody> <tr> <td>If the draw is being performed... below the IV</td> <td>Then place the tourniquet... between the IV and the draw site and perform the venipuncture.</td> </tr> <tr> <td>above the IV</td> <td>3-4 inches above the antecubital vein and perform the venipuncture.</td> </tr> </tbody> </table>	If the draw is being performed... below the IV	Then place the tourniquet... between the IV and the draw site and perform the venipuncture.	above the IV	3-4 inches above the antecubital vein and perform the venipuncture.
If the draw is being performed... below the IV	Then place the tourniquet... between the IV and the draw site and perform the venipuncture.				
above the IV	3-4 inches above the antecubital vein and perform the venipuncture.				
3	Indicate that the draw was performed on the arm where an IV was present [KPPI, sticker on tubes], as required by your Medical Center policy.				
4	Have the patient's nurse turn the IV back on.				

Non-controlled documents

- CLSI – GP41-A6, Procedures for the Collection of Diagnostic Blood Specimens by Venipuncture; Approved Standard – Sixth Edition, October 2007, page 23.
- Phlebotomy Tutorial from library.med.utah.edu/WebPath/Tutorial/Phleb/Phleb.html
- Annotations from Phlebotomy Procedures from Los Angeles and Orange County Medical Centers

Controlled documents

Regional Parent Document Reference Number: SCPMG-PPP-0110 Rev. 02

Signature Manifest

Document Number: SBMC-PPP-0032

Revision: 02

Title: Drawing Blood from Patients with an IV

Effective Date: 10 May 2018

All dates and times are in Pacific Standard Time.

Review: SBMC-PPP-0032 02 Drawing Blood from Patients with an IV

Review

Name/Signature	Title	Date	Meaning/Reason
Qiyamaa Portillo (K237031)	Assistant Director Operations	20 Mar 2020, 10:57:36 AM	Reviewed
Janice Wolf (K119893)	Director Operations Area Lab	03 Apr 2020, 12:16:29 PM	Reviewed