

Document Name: Tube Coagulase Test

Approved By:

Status: **DRAFT**

PURPOSE: The tube coagulase test is used to differentiate *Staphylococcus aureus* from other *Staphylococcus* spp. by determining the organism's ability to clot plasma using the enzyme coagulase.

SAMPLE INFORMATION:

Type	<p>Several, well-isolated colonies that are:</p> <ul style="list-style-type: none"> • Gram-positive cocci • Catalase positive • From a 18 to 24 hour culture
-------------	---

REAGENTS and/or MEDIA:

Type	BD BBL Coagulase Plasma, Rabbit with EDTA
Reagent Preparation	Reconstitute lyophilized reagent with 15 mL of sterile water
Stability and Storage Requirements	<ul style="list-style-type: none"> • Store unopened lyophilized coagulase plasma at 2°C to 8°C. • Store reconstituted plasma at 2°C to 8°C for up to 14 days. • Do not use if the product is caked, discolored or shows other signs of deterioration. • Examine reconstituted reagents for evidence of contamination, evaporation or other signs of deterioration, such as cloudiness or partial clotting.

SUPPLIES:

- | | |
|--|---|
| <ul style="list-style-type: none"> • Sterile water • 10 mL plastic syringe • Glass test tubes | <ul style="list-style-type: none"> • Sterile pipettes • Disposable loops • 35° ambient air incubator |
|--|---|

NOTE: This is a controlled document for internal use only. Any documents appearing in paper form are not controlled and should be checked against electronic version prior to use.

FILENAME:

Print Date:

SPECIAL SAFETY PRECAUTIONS:

Containment Level 2 facilities, equipment, and operational practices for work involving infectious or potentially infectious materials or cultures.

- Lab gown must be worn when performing activities with potential pathogens.
- Gloves must be worn when direct skin contact with infected materials is unavoidable.
- Eye protection must be used when there is a known or potential risk of exposure to splashes.
- All procedures that may produce aerosols, or involve high concentrations or large volumes should be conducted in a biological safety cabinet (BSC).
- The use of needles, syringes and other sharp objects should be strictly limited.

All patient specimens are assumed to be potentially infectious. Universal precautions must be followed. Since viable micro-organisms are used, all cultures must be handled with appropriate precautions. All equipment in contact with cultures should be decontaminated by appropriate methods.

QUALITY CONTROL:

- Quality control is performed daily:
 - Positive: *Staphylococcus aureus* ATCC 25923
 - Negative: *Staphylococcus epidermidis* ATCC 12228
- A TQC order is automatically generated daily to record the QC results.

NOTE: This is a controlled document for internal use only. Any documents appearing in paper form are not controlled and should be checked against electronic version prior to use.

FILENAME:

Print Date:

PROCEDURE INSTRUCTIONS:

Step	Action
Performing the tube coagulase test	
1	Using a sterile pipette, add approximately 0.5 mL of rehydrated coagulase reagent to a labelled glass test tube.
2	Using a disposable loop, thoroughly emulsify several colonies from a non-inhibitory agar plate in the tube of plasma and mix gently.
3	Incubate in the O ₂ incubator for up to 6 hours.
4	Examine the tubes periodically by gently tipping the tube. Avoid shaking or agitating the tube, which could cause breakdown of the clot and false negative results.
5	If no clot is visible after 6 hours, leave at room temperature overnight and observe for clot formation.

INTERPRETATION OF RESULTS:

IF	THEN
Any degree of clot formation	Tube coagulase = Positive
No clot formation	Tube coagulase = Negative



NOTE: This is a controlled document for internal use only. Any documents appearing in paper form are not controlled and should be checked against electronic version prior to use.

FILENAME:

Print Date:

LIMITATIONS/PRECAUTIONS:

1. Some species of organisms utilize citrate in their metabolism and will yield false-positive reactions for coagulase activity. Normally, this does not cause problems since the coagulase test is performed almost exclusively on *Staphylococci*. However, it is possible that bacteria that utilize citrate may contaminate *Staphylococcus* cultures on which the coagulase test is being performed. These contaminated cultures may, upon prolonged incubation, give false-positive results due to citrate utilization.
2. Some strains of *S.aureus* produce staphylokinase, which may lyse clots. If the tubes are not read until 24 hours of incubation, false-negative results may occur.
3. Do not use plasma if a heavy precipitate or clot has formed before inoculation.
4. Do not shake or agitate the tube during the test. This may cause a breakdown of the clot, which will not reform upon additional incubation.
5. False negative coagulase reactions may occur if the test culture is older than 18 to 24 hours or if there is scant inoculum.
6. False-positive results may occur if the test isolate is removed from agar containing high concentrations of salt. Use only strains grown on non-selective media.

REFERENCES:

- BD BBL, Coagulase Plasmas package insert, 2017-02
- Clinical Microbiology Procedures Handbook, 4th edition, ASM Press, 2016

REVISION HISTORY:

REVISION	DATE	Description of Change	REQUESTED BY
1.0	4 APR 19	Initial Release	L. Steven

NOTE: This is a controlled document for internal use only. Any documents appearing in paper form are not controlled and should be checked against electronic version prior to use.

FILENAME:

Print Date: