

PROGRAM Standard Operating Procedure – Laboratory Services	
Title: MIC51400 – ALA Test	Policy Number:
Program Name: Laboratory Services	
Applicable Domain: Lab, DI and Pharmacy Services	
Additional Domain(s):	
Effective Date:	Next Review Date:
Issuing Authority: Director of Health Services	Date Approved:
Accreditation Canada Applicable Standard: N/A	

GUIDING PRINCIPLE:

The ALA (Aminolevulinic Acid) test is used to rapidly detect porphyrin as a means of speciating *Haemophilus* species.

PURPOSE/RATIONALE:

This standard operating procedure describes how to perform the ALA test.

SCOPE/APPLICABILITY:

This procedure applies to Medical Laboratory Technologists (MLTs) performing the ALA test.

SAMPLE INFORMATION:

Type	Few, well isolated colonies that are: <ul style="list-style-type: none"> • Tiny, gram-negative bacilli or coccobacilli • Growing only on Chocolate agar • Possess typical <i>Haemophilus</i> colonial morphology • Do not grow on Blood agar • 18 to 24 hours old
-------------	--

REAGENTS and/or MEDIA:

Type	remel A.L.A. Disk
Stability and Storage Requirements	<ul style="list-style-type: none"> • Store at 2°C to 8°C • Allow to come to room temperature before use • Protect disks from light, as the substrate is highly light sensitive

Disclaimer Message: This is a **CONTROLLED** document for internal use only. Any documents appearing in paper form are not controlled and should be checked against the electronic file version prior to use.

SUPPLIES:

- Forceps
- Wooden sticks
- Disposable loops
- Blood agar
- Filter paper
- Sterile water

EQUIPMENT

- 35° ambient air incubator
- Long wave ultraviolet lamp

NOTE:

- **The A.L.A test is only to be used on small, Gram-negative bacilli resembling *Haemophilus* spp. and must be confirmed with Vitek 2 NH card or API NH if from sterile site**
- **Any positive result from a sterile site for *Haemophilus influenzae* must be sent immediately to Provincial Lab Edmonton for typing as soon as identification is confirmed. Assure there is a purity plate made that can be used for this purpose and can be sent out the day the identification is confirmed. Refer to MIC10510- Referral of Category B Specimens to DynaLIFE and Alberta Precision Labs**

SPECIAL SAFETY PRECAUTIONS:

Containment Level 2 facilities, equipment, and operational practices for work involving infectious or potential infectious materials or cultures.

- Ensure that appropriate hand hygiene practices be used.
- Lab gown must be worn when performing activities with potential pathogens.
- Gloves must be worn when direct skin contact with infected materials is unavoidable.
- Eye protection must be used when there is a known or potential risk of exposure of splashes.
- All procedures that may produce aerosols, or involve high concentrations or large volumes should be conducted in a biological safety cabinet (BSC).
- The use of needles, syringes and other sharp objects should be strictly limited.

All patient specimens are assumed to be potentially infectious. Routine Practices must be followed. Since viable micro-organisms are used, all cultures must be handled with appropriate precautions. All equipment in contact with cultures should be decontaminated by appropriate methods.


QUALITY CONTROL:

- Quality control is performed weekly:
 - Positive: *Aggregatibacter aphrophilus* ATCC 7901
 - Negative: *Haemophilus influenzae* ATCC 10211
- A TQC order is automatically generated on Wednesdays to record the QC results

PROCEDURE INSTRUCTIONS:

Step	Action
Performing the ALA test	
1	Verify the test isolate resembles <i>Haemophilus</i> spp. by gram-stain and colonial morphology.
2	Place the ALA disk, with the "A" side down, on the agar surface of a Blood agar plate.
3	Inoculate the disk with a heavy, visible inoculum from a pure 18 to 24 hour culture of the test isolate.
4	Place a piece of filter paper saturated with water on the lid of the agar plate.
5	Incubate for up to 6 hours in the O ₂ incubator.
6	Examine the disk at 1 hour under the ultraviolet light in the dark room for orange fluorescence. If negative, re-incubate the test and examine hourly for up to 6 hours before reporting as negative.

INTERPRETATION OF RESULTS:

IF	THEN
Bright orange fluorescence	ALA = Positive
No fluorescence	ALA = Negative
	

LIMITATIONS:

1. The ALA test is used for differentiating *Haemophilus* spp. only.
2. Best results are obtained using a heavy inoculum. False negative reactions may occur if the inoculum is insufficient or if the culture is greater than 24 hours old.
3. Many organisms will give a positive reaction. If test is performed only on Gram-negative bacillus colonies that grow well on Chocolate agar within 24 hours and not on Blood agar, results are for *Haemophilus* spp.
4. Organisms that are strongly oxidase positive or catalase positive may give a false positive result. Such organisms make heme and its precursors from ALA in the process of synthesizing oxidase or catalase. Verify that the test organism resembles *Haemophilus* both by Gram stain and colonial morphology before testing.
5. Product should not be used if the colour has changed from white or the desiccant has changed from blue to pink.
6. If the filter paper is not kept moist during incubation, the reaction can be falsely negative.

Disclaimer Message: This is a **CONTROLLED** document for internal use only. Any documents appearing in paper form are not controlled and should be checked against the electronic file version prior to use.

CROSS-REFERENCES:

- MIC10510-Referral of Category B Specimens to *DynaLIFE* and Alberta Precision Laboratories

REFERENCES:

1. remel. (2010-07). *A.L.A. Disk* package insert

APPROVAL:

Date

REVISION HISTORY:

REVISION	DATE	Description of Change	REQUESTED BY
1.0	08 Apr 19	Initial Release	L. Steven
2.0	30 Jun 21	Procedure reviewed and added to NTHSSA policy template	L. Steven

Disclaimer Message: This is a **CONTROLLED** document for internal use only. Any documents appearing in paper form are not controlled and should be checked against the electronic file version prior to use.