PROGRAM Standard Operating Procedure – Laboratory Services			
Title: MIC32300 – Respiratory Culture	Policy Number:		
Program Name: Laboratory Services			
Applicable Domain: Lab, DI and Pharmacy Services			
Additional Domain(s):			
Effective Date:	Next Review Date:		
Issuing Authority:	Date Approved:		
Director of Health Services			
Accreditation Canada Applicable Standard: N/A			

GUIDING PRINCIPLE:

Pneumonia may be categorized as: i) Community acquired pneumonia (CAP), ii) Nosocomial or Hospital acquired pneumonia (NAP / HAP), iii) Aspiration pneumonia and iv) Pneumonia in immunocompromised patients. The most common organisms to cause CAP include *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Legionella pneumophila*. HAP is more commonly due to aerobic gram-negative bacilli, anaerobes, *Staphylococcus aureus*, *Streptococcus pneumoniae* and others. Aspiration pneumonia may be due to a mixture of oral aerobes and anaerobes.

PURPOSE/RATIONALE:

This standard operating procedure describes how to determine the significance of growth in respiratory specimens.

SCOPE/APPLICABILITY:

This procedure applies to Medical Laboratory Technologists (MLTs) processing specimens for respiratory culture.

Туре	Sterile container	
Source	 Sputum Endotracheal aspirate (ETT) and Auger suction Bronchial aspirates and Bronchoalveolar lavage (BAL) 	
Stability	 If the sample is received in the laboratory and processed greater than 72 hours from collection: Add specimen quality comment: "Delayed transport may adversely affect pathogen recovery" 	
Storage Requirements	Refrigerated	

SAMPLE INFORMATION:

Title: MIC32300-Respiratory Culture	Type: Laboratory Services Program SOP
Issuing Authority: Director of Health Services	Policy Number:
Next Review Date:	Date Approved:

Criteria for rejection	 Unlabeled/mislabeled specimen Specimen container label does not match patient identification on requisition Duplicate specimens obtained with same collection method within 24 hours Leaking specimens Improperly collected, labeled, transported, or handled bronchial aspirate, BAL specimens, lung aspirates and lung biopsy specimens should be processed. Waiver of responsibility form SCM-40110 needs to be filled out by the responsible nurse
---------------------------	--

REAGENTS and/or MEDIA:

- Blood agar (BA), Chocolate agar (CHO) and MacConkey agar (MAC)
- Identification reagents: catalase, oxidase, Staph latex test, Strep latex test, etc.

SUPPLIES:

- Disposable inoculation needles
- Microscope slides
- Wooden sticks

EQUIPMENT:

- Biosafety cabinet
- 35° ambient air and 35° CO₂ incubators
- Vitek 2 and supplies

SPECIAL SAFETY PRECAUTIONS:

Containment Level 2 facilities, equipment, and operational practices for work involving infectious or potentially infectious materials or cultures:

- Ensure that appropriate hand hygiene practices be used
- Lab gown must be worn when performing activities with potential pathogens
- Gloves must be worn when direct skin contact with infected materials is unavoidable
- Eye protection must be used when there is a known or potential risk of exposure of splashes
- All procedures that may produce aerosols, or involve high concentrations or large volumes should be conducted in a biological safety cabinet (BSC)
- The use of needles, syringes and other sharp objects should be strictly limited

All patient specimens are assumed to be potentially infectious. Routine Practices must be followed. Since viable micro-organisms are used, all cultures must be handled with appropriate precautions. All equipment in contact with cultures should be decontaminated by appropriate methods.

QUALITY CONTROL:

Refer to Test Manual for reagent quality control procedures

PROCEDURE INSTRUCTIONS:

Step	Action		
Processing specimens for respiratory culture			
1	 In the biosafety cabinet: Use a sterile swab to inoculate BA, CHO, and MAC from the specimen. Select the most purulent or most blood-tinged portion Streak for isolated growth using a disposable inoculation needle Prepare a smear by rolling the swab gently across the slide to avoid destruction of cellular elements and disruption of bacterial arrangements 		
2	 Incubate all media: Place BA and CHO in the CO₂ incubator Place MAC in the O₂ incubator 		
3	Allow smear to dry and perform gram stain. Gram stain must be read before culture plates. Refer to MIC20115-Gram Stain Procedure.		
4	Ensure the quality of the specimen has been evaluated and is considered acceptable for culture. Refer to MIC20300-Gram stain resulting in LIS- Respiratory Cultures. NOTE: Bronchial wash and bronchoalveolar lavage specimens are processed regardless of specimen quality		

Probable Pathogens			
 Bacillus anthracis*+ Brucella spp.*+ Burkholderia mallei/pseudomallei*+ Cryptococcus neoformans/gattii Fungi and Molds 	 Neisseria gonorrhoeae Nocardia spp. Streptococcus pyogenes Streptococcus agalactiae in newborn Yersinia pestis*+ 		
Potential	Pathogens		
 Acinetobacter spp. β-hemolytic strep Burkholderia spp. Corynebacterium pseudodiphtheriticum Fastidious Gram-negative bacilli Haemophilus influenzae Moraxella catarrhalis 			
Commensal Flora			
 Anaerobes Capnocytophaga spp. Coagulase negative Staphylococci Eikenella spp. Eikenella spp. Eikenella spp. Eikenella spp. 	hilus spp. nogenic Yeast spp.		
Risk group 3 organism. If suspected, refer to Policy B-0160: "Specimens Containing Suspected Risk Group 3 Pathogens" for Primary Specimen Handling Flow			

Chart +All work should be performed in the BSC

Step		Action		
		correlates with gram stain results. If		
	discordant results are found between the gram stain and growth:			
1	Re-examine smear and cultu	re plates		
-		Check for anaerobic growth		
	Re-incubate media to resolve			
	Consider re-smearing or re-planting specimen			
2	Observe BA and CHO plates a	at 24 hours and 48 hours		
2	Observe MAC plate at 24 hours			
	Significant growth is defined as	bacterial morphotypes that are:		
-	 Moderate to beavy growth in the second or greater guadr 			
3		of the plate provided there is little or no		
	other normal respiratory flora			
		ort the following probable pathogens:		
	Bacillus anthracis	Fungi and Molds		
	• Brucella spp.	Neisseria gonorrhoeae		
	Burkholderia	Nocardia spp.		
	mallei/pseudomallei	Streptococcus pyogenes		
	Cryptococcus	 Streptococcus agalactiae in newborns 		
	neoformans/gattii	Yersinia pestis		
	Report the following potential pathogens, but do not make an effort to find low numbers, unless seen in smear:			
	Streptococcus pneumoniae	Haemophilus influenzae		
	Report the following potential pathogens if present in significar amounts, even if not predominant:			
	amounts, even if not predom	inant:		
	Moraxella catarrhalis	inant:Neisseria meningitidis		
	Moraxella catarrhalis Report the following potentia	• Neisseria meningitidis al pathogens if present in significant		
4	Moraxella catarrhalis Report the following potentia amounts, even if not predom	• Neisseria meningitidis al pathogens if present in significant inant for inpatients only:		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: Single morphotype of enteric 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred β-hemolytic strep 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant iominant organism in the culture: Single morphotype of enteric Gram-negative bacilli 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred β-hemolytic strep Corynebacterium pseudodiphtheriticum Fungi and Molds 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: Single morphotype of enteric Gram-negative bacilli Staphylococcus aureus 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred β-hemolytic strep Corynebacterium pseudodiphtheriticum 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: Single morphotype of enteric Gram-negative bacilli Staphylococcus aureus 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred β-hemolytic strep Corynebacterium pseudodiphtheriticum Fungi and Molds Report as "commensal flora" 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: Single morphotype of enteric Gram-negative bacilli Staphylococcus aureus 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred β-hemolytic strep Corynebacterium pseudodiphtheriticum Fungi and Molds Report as "commensal flora" Note: If Enterococci, coagulase- 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: Single morphotype of enteric Gram-negative bacilli Staphylococcus aureus regative Staphylococci or Candida spp. 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred β-hemolytic strep Corynebacterium pseudodiphtheriticum Fungi and Molds Report as "commensal flora" 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: Single morphotype of enteric Gram-negative bacilli Staphylococcus aureus regative Staphylococci or Candida spp. list with minimal identification 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred β-hemolytic strep Corynebacterium pseudodiphtheriticum Fungi and Molds Report as "commensal flora" Note: If Enterococci, coagulase- are the only organisms present, Anaerobes 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: Single morphotype of enteric Gram-negative bacilli Staphylococcus aureus regative Staphylococci or Candida spp. list with minimal identification Enterococci spp. 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred β-hemolytic strep Corynebacterium pseudodiphtheriticum Fungi and Molds Report as "commensal flora" Note: If Enterococci, coagulase- are the only organisms present, Anaerobes Capnocytophaga spp. 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: Single morphotype of enteric Gram-negative bacilli Staphylococcus aureus regative Staphylococci or Candida spp. list with minimal identification Enterococci spp. Haemophilus spp. not H.influenzae 		
4	 Moraxella catarrhalis Report the following potentia amounts, even if not predom Acinetobacter spp. Burkholderia spp. Report the following potentia amounts AND if it is the pred β-hemolytic strep Corynebacterium pseudodiphtheriticum Fungi and Molds Report as "commensal flora" Note: If Enterococci, coagulase- are the only organisms present, Anaerobes 	 Neisseria meningitidis al pathogens if present in significant inant for inpatients only: Pseudomonas aeruginosa Stenotrophomonas maltophilia al pathogens if present in significant ominant organism in the culture: Single morphotype of enteric Gram-negative bacilli Staphylococcus aureus regative Staphylococci or Candida spp. list with minimal identification Enterococci spp. 		

REPORTING INSTRUCTIONS:

IF	REPORT
No growth after 1 day	 PRELIM: Report: "No Growth after 1 Day" Report: "Further report to follow"
No growth after 2 days	FINAL:Report: "No Growth after 2 Days"
Growth of probable pathogen	 Report organism full identification List quantitation Report susceptibility testing as per ASTM
Growth of potential pathogens that meets criteria for reporting	 Report organism full identification List quantitation Report susceptibility testing as per ASTM
Growth of potential pathogen that does NOT meet criteria for reporting	 Report: "Commensal flora" List quantitation
Growth of commensal flora	 Report: "Commensal flora" List quantitation
Mix of enteric Gram-negative bacilli	 Report: "Mixture of coliform organisms" List quantitation

NOTE:

- Refer to Reportable Diseases Public Health Act as of September 2009 for reporting to OCPHO (HPU1)
- Refer to LQM70620-Laboratory Critical Results List-Microbiology for results that need to be phoned to ordering location
- Refer to MIC36100-Nosocomial Infection Notification Job Aid to determine if organism needs to be copied to Infection Prevention and Control
- Refer to MIC36200-Referral of Category A Specimens to APL for sending category A isolates to APL
- Refer to MIC36300-Referral of Category B Specimens to APL for sending isolates to APL
- Refer to MIC36400-Referral of Category B Specimens to DL for sending isolates to DynaLIFE

LIMITATIONS:

- 1. A positive culture with *Streptococcus pneumoniae* or *Haemophilus influenzae* generally indicates an infection, although carriage may lead to false-positive results.
- 2. False-negative cultures can result from improper collection, delayed transport, contamination of the specimen with normal oral microbiota, low organism levels or from prior antimicrobial therapy.
- 3. False-positive cultures can result from contamination of the specimen by normal respiratory flora and its subsequent growth on culture and over interpretation by the laboratory.

Title: MIC32300-Respiratory Culture	Type: Laboratory Services Program SOP
Issuing Authority: Director of Health Services	Policy Number:
Next Review Date:	Date Approved:

CROSS-REFERENCES:

- Policy B-0160: "Specimens Containing Suspected Risk Group 3 Pathogens"
- LQM70620-Laboratory Critical Results List-Microbiology
- SCM40110-Waiver of Responsibility
- MIC20115-Gram Stain Procedure
- MIC20300-Gram stain resulting in LIS-Respiratory Cultures
- MIC36100-Nosocomial Infection Notification Job Aid
- MIC36200-Referral of Category A Specimens to APL
- MIC36300-Referral of Category B Specimens to APL
- MIC36400-Referral of Category B Specimens to DL

REFERENCES:

- 1. Leber, A. (2016). *Clinical microbiology procedures handbook.* (4thed.) Washington, D.C.: ASM Press
- Jorgensen J.H., Pfaller M.A., Carroll K.C., Funke G., Landry M.L., Richter S.S., Warnock D.W. (2015). *Manual of Clinical Microbiology*, 11th edition. Washington, D.C: ASM Press

APPROVAL:

Date

REVISION HISTORY:

REVISION	DATE	Description of Change	REQUESTED BY
1.0	28 May 18	Initial Release	L. Steven
2.0	22 Feb 21	Procedure reviewed and added to NTHSSA policy template	L. Steven
3.0	27 Feb 23	Procedure reviewed	L. Steven