

Get the facts about Biotin

What is biotin?

Biotin is a water-soluble B-vitamin (B7) that is integral to energy and metabolism (gluconeogenesis, fatty acid synthesis, and carbohydrate utilization).¹

Why has biotin intake increased?

Biotin has gained increasing popularity as an over-the-counter supplement and is commonly included in multivitamins and beauty products marketed for hair and nails.²⁻⁶ Many of these hair-and-nail vitamins include biotin at concentrations up to 100 times the Dietary Reference Intake (e.g., 3000 µg); some report as much as 10,000 µg.² In addition, clinicians may prescribe biotin supplementation to help prevent biotin deficiency in pregnancy⁷⁻⁸ or to reduce leg cramps in dialysis patients.⁹ Clinicians may also prescribe high doses of biotin for multiple sclerosis, inborn metabolic disorders, and mitochondrial energy disorders.^{9,10-15}

What is the Dietary Reference Intake for biotin?

The Dietary Reference Intake (DRI), or adequate intake in the case of biotin, is age-dependent. In adults, the adequate intake is 30–70 µg/day.^{1,16-18} This corresponds to an adult reference range of approximately 0.12–0.54 ng/mL,¹⁹ depending on the population from which the reference interval was derived.

In what foods is biotin found?

Meats and Poultry

- Beef Liver
- Hamburger
- Pork Chop
- Egg

Fish

- Salmon
- Tuna

Dairy

- Milk (2%)
- Yogurt (Plain)

Fruits and Vegetables

- Sweet Potato
- Spinach
- Broccoli
- Banana
- Apple

Nuts and Grains

- Sunflower Seeds
- Almonds
- Oatmeal
- Bread (Whole Wheat)

What is the potential risk of biotin interference with clinical laboratory tests?

- Multiple manufacturers use a streptavidin biotin complex in many of their immunoassays. Advantages of this complex include its high binding affinity, adaptability for binding antigen or antibody, and ability to readily attach to a solid phase (such as a microbead).
- Supra-physiological doses of biotin ingested for either cosmetic or pharmacologic use can result in serum concentrations as high as 1160 ng/mL ($\mu\text{g/L}$) 1 hour after a single oral biotin dose of 300 mg.²⁰
- Biotin interference can cause either falsely depressed or falsely elevated patient test results.


How long does it take after biotin use is discontinued for a patient's biotin to reach a level that does not impact results?

The time after last use required for a patient's biotin to reach a level that does not impact results depends on a variety of factors including, but not limited to, dose, duration of use, clinical conditions, age, and the half-life of biotin in the serum.

- For a single oral biotin dose of approximately 600 μg , which is greater than DRI, the half-life has been reported as less than 2 hours.²¹
- More recently, the half-life for single oral biotin doses between 100–300 mg (100,000–300,000 μg) has been shown to vary between 8–19 hours.¹²⁻¹³
- For individuals ingesting mega-doses of biotin, up to 300 mg/day, serum concentrations as high as 1160 ng/mL have been observed.¹⁴

References:

1. Zempleni J, et al. *Expert Rev Endocrinol Metab.* 2008;3:715-24.
2. Palsdottir H. 7 Health Benefits of Biotin. September 11, 2016. Authoritative Nutrition. Available from: <https://authoritativnutrition.com/biotin-benefits/>.
3. Paxton A. CAP Today. 2016. Available from: <http://www.captodayonline.com/beauty-fads-ugly-downsidetest-interference>.
4. Boccaletti V, et al. *Pediatr Dermatol.* 2007;24:E14-6.
5. Bolander FF. *Curr Opin Investig Drugs.* 2006;7:912-5.
6. Seymons K, et al. *Pediatr Dermatol.* 2004;21:231-5.
7. Mock DM, et al. *J Nutr.* 1997;127:710-6.
8. Perry CA, et al. *J Nutr.* 2014;144:1977-84.
9. Oguma S, et al. *Tohoku J Exp Med.* 2012;227:217-23.
10. de Parscau L, et al. *Pediatric.* 1989;44:383-6.
11. Kummer S, et al. *N Engl J Med.* 2016;375:704-6.
12. Peyro Saint Paul L, et al. *Expert Opin Drug Metab Toxicol.* 2016;12:327-44.
13. Sedel F, et al. *Neuropharmacology.* 2016;110:644-53.
14. Tabarki B, et al. *Neurology.* 2013;80:261-7.
15. Tourbah A, et al. *Mult Scler.* 2016;22:1719-31.
16. 2016 Institute of Medicine report; Dietary Reference Intakes (DRIs): Vitamins.
17. Samarasinghe S, et al. *Endocr Pract.* 2017.
18. Zempleni J, Mock DM. *Am J Clin Nutr.* 1999;69:504-8.
19. Burtis CA, et al. *Tietz Textbook of Clinical Chemistry and Molecular Diagnostics.* 5th ed. Elsevier Health Sciences; c2012.
20. Piketty ML, et al. *Clin Chem Lab Med.* 2017;55:817-25.
21. Bitsch R, et al. *Int J Vitam Nutr Res.* 1989;59:65-71.



Ask your patients:

- Are you currently taking a multivitamin that includes biotin?
- Are you currently taking a hair, skin, and nail supplement?
- Are you currently taking biotin as part of a therapeutic regimen?

Ask your clinical laboratory provider:

- What other interferents may impact patient results?

Siemens Healthineers Headquarters

Siemens Healthcare GmbH
Henkestr. 127
91052 Erlangen,
Germany
Phone: +49 9131 84-0
siemens.com/healthineers

Local Contact Information

Siemens Healthcare Diagnostics Inc.
Laboratory Diagnostics
511 Benedict Avenue
Tarrytown, NY 10591-5005
USA
Phone: +1 914-631-8000
siemens.com/healthineers