

---

**Janssen Research & Development \***

**Clinical Protocol**

---

**A Phase 2, Multicohort Open-Label Study of JNJ-68284528, a Chimeric Antigen Receptor T cell (CAR-T) Therapy Directed Against BCMA in Subjects with Multiple Myeloma**

**CARTITUDE-2**

---

**Protocol 68284528MMY2003; Phase 2  
AMENDMENT 3/USA-1**

JNJ-68284528 (ciltacabtagene autoleucel)

\*Janssen Research & Development is a global organization that operates through different legal entities in various countries. Therefore, the legal entity acting as the sponsor for Janssen Research & Development studies may vary, such as, but not limited to Janssen Biotech, Inc.; Janssen Products, LP; Janssen Biologics, BV; Janssen-Cilag International NV; Janssen Pharmaceutica NV; Janssen, Inc; Janssen Sciences Ireland UC; or Janssen Research & Development, LLC. The term “sponsor” is used throughout the protocol to represent these various legal entities; the sponsor is identified on the Contact Information page that accompanies the protocol.

US sites of this study will be conducted under US Food & Drug Administration IND regulations (21 CFR Part 312).

**EudraCT NUMBER:** 2018-004124-10

**Status:** Approved

**Date:** 14 November 2020

**Prepared by:** Janssen Research & Development, LLC

**EDMS number:** EDMS-RIM-222691, 1.0

Local Amendment to: EDMS-ERI-164244214, 4.0

**GCP Compliance:** This study will be conducted in compliance with Good Clinical Practice, and applicable regulatory requirements.

---

**Confidentiality Statement**

The information provided herein contains Company trade secrets, commercial or financial information that the Company customarily holds close and treats as confidential. The information is being provided under the assurance that the recipient will maintain the confidentiality of the information under applicable statutes, regulations, rules, protective orders or otherwise.

**TABLE OF CONTENTS**

<b>TABLE OF CONTENTS .....</b>	<b>2</b>
<b>LIST OF ATTACHMENTS .....</b>	<b>5</b>
<b>LIST OF IN-TEXT TABLES AND FIGURES .....</b>	<b>6</b>
<b>PROTOCOL AMENDMENTS .....</b>	<b>8</b>
<b>DOCUMENT HISTORY .....</b>	<b>8</b>
<b>SYNOPSIS .....</b>	<b>28</b>
<b>TIME AND EVENTS SCHEDULES: COHORT A, COHORT B, AND COHORT C (TABLE 1 AND TABLE 2) .....</b>	<b>33</b>
<b>TIME AND EVENTS SCHEDULES: COHORT D (TABLE 3 AND TABLE 4) .....</b>	<b>44</b>
<b>TIME AND EVENTS SCHEDULES: COHORT E (TABLE 5, TABLE 6, TABLE 7, AND TABLE 8) .....</b>	<b>56</b>
<b>ABBREVIATIONS .....</b>	<b>72</b>
<b>1. INTRODUCTION .....</b>	<b>75</b>
1.1. Background .....	75
1.1.1. Multiple Myeloma .....	75
1.1.2. B cell Maturation Antigen .....	75
1.1.3. CAR-T Therapy .....	76
1.1.4. JNJ-68284528 .....	76
1.1.5. Clinical Studies .....	77
1.2. Overall Rationale for the Study .....	80
1.3. Potential Safety Risks and Mitigation Strategies .....	81
<b>2. OBJECTIVES, ENDPOINTS, AND HYPOTHESIS .....</b>	<b>83</b>
2.1. Objectives and Endpoints .....	83
2.2. Hypotheses .....	85
<b>3. STUDY DESIGN AND RATIONALE .....</b>	<b>85</b>
3.1. Overview of Study Design .....	85
3.1.1. Cohort A Study Design .....	86
3.1.2. Cohort B Study Design .....	89
3.1.3. Cohort C Study Design .....	90
3.1.4. Cohort D Study Design .....	93
3.1.5. Cohort E Study Design .....	96
3.1.6. Retreatment with JNJ-68284528 .....	98
3.2. Rationale of Dose and Administration Schedule Selection .....	98
3.3. Study Design Rationale .....	99
<b>4. SUBJECT POPULATION .....</b>	<b>103</b>
4.1. Cohort A Eligibility Criteria .....	104
4.1.1. Cohort A Inclusion Criteria .....	104
4.1.2. Cohort A Exclusion Criteria .....	106
4.2. Cohort B Eligibility Criteria .....	110
4.2.1. Cohort B Inclusion Criteria .....	110
4.2.2. Cohort B Exclusion Criteria .....	113
4.3. Cohort C Eligibility Criteria .....	116
4.3.1. Cohort C Inclusion Criteria .....	116
4.3.2. Cohort C Exclusion Criteria .....	118

4.4.	Cohort D Eligibility Criteria .....	121
4.4.1.	Cohort D Inclusion Criteria.....	121
4.4.2.	Cohort D Exclusion Criteria .....	124
4.5.	Cohort E Eligibility Criteria .....	127
4.5.1.	Cohort E Inclusion Criteria.....	127
4.5.2.	Cohort E Exclusion Criteria.....	129
4.6.	Prohibitions and Restrictions .....	132
<b>5.</b>	<b>TREATMENT ALLOCATION AND BLINDING .....</b>	<b>132</b>
<b>6.</b>	<b>DOSAGE AND ADMINISTRATION .....</b>	<b>133</b>
6.1.	Study Treatment Administration.....	133
6.1.1.	Criteria for Apheresis (all cohorts unless otherwise specified).....	133
6.1.2.	Administration of Conditioning Regimen (Cyclophosphamide and Fludarabine) (All Cohorts).....	134
6.1.2.1.	Criteria for Conditioning Regimen (Cyclophosphamide and Fludarabine Dosing) (All Cohorts).....	134
6.1.3.	JNJ-68284528 Administration (All Cohorts) .....	135
6.1.3.1.	Exceptional Release Criteria .....	136
6.1.3.2.	Evaluation Prior to Administration of JNJ-68284528 (All Cohorts).....	137
6.1.3.3.	Pre-JNJ-68284528 Infusion Supportive Therapy (All Cohorts).....	137
6.1.4.	Lenalidomide (Cohort D).....	137
6.1.5.	Cohort E: Induction and Post JNJ-68284528 Therapy.....	140
6.1.5.1.	Cohort E: D-VRd Induction Treatment .....	140
6.1.5.1.1.	Daratumumab.....	141
6.1.5.1.1.1.	Daratumumab Subcutaneous Preparation .....	141
6.1.5.1.1.2.	Daratumumab Administration .....	141
6.1.5.1.2.	Bortezomib .....	143
6.1.5.1.3.	Lenalidomide .....	143
6.1.5.1.4.	Dexamethasone .....	144
6.1.5.1.5.	Dose Modification Guidelines for Bortezomib, Lenalidomide, and Dexamethasone .....	144
6.1.5.2.	Cohort E: D+R Consolidation Treatment.....	151
6.1.5.3.	Cohort E: Medications Related to Daratumumab Dosing .....	153
6.2.	Management Guidelines for Potential Risks (All Cohorts).....	153
6.2.1.	Management of Cytokine Release Syndrome .....	153
6.2.2.	Neurologic Toxicities.....	156
6.2.2.1.	CAR-T Cell-related Neurotoxicity (Immune Effector Cell-Associated Neurotoxicity Syndrome [ICANS]) .....	156
6.2.2.2.	Other Neurotoxicities .....	159
6.2.3.	Tumor Lysis Syndrome.....	160
6.2.4.	Second Primary Malignancy .....	160
6.2.5.	Cytopenia.....	160
6.2.6.	Hypogammaglobulinemia .....	161
6.2.7.	Infections .....	161
6.2.8.	Hypersensitivity Reactions.....	162
6.2.9.	Cohort E: Management of Infusion-related Reactions and Local Injection-site Reactions of Daratumumab SC .....	162
6.2.9.1.	Infusion-related Reactions.....	162
6.2.9.2.	Injection-site Reactions .....	163
<b>7.</b>	<b>TREATMENT COMPLIANCE.....</b>	<b>163</b>
<b>8.</b>	<b>PRESTUDY AND CONCOMITANT THERAPY .....</b>	<b>163</b>
8.1.	Prophylaxis for Herpes Zoster Reactivation.....	165
8.2.	Permitted Medications.....	165
8.3.	Prohibited Therapies .....	165
8.3.1.	Additional Prohibited Therapies for Cohort E .....	166
8.4.	Subsequent Anticancer Therapy.....	167

<b>9. STUDY EVALUATIONS .....</b>	<b>167</b>
9.1. Study Procedures.....	167
9.1.1. Overview.....	167
9.1.2. Screening Phase .....	167
9.1.3. Apheresis .....	168
9.1.4. Cyclophosphamide and Fludarabine Conditioning Regimen .....	168
9.1.5. JNJ-68284528 Administration .....	168
9.1.6. Post-treatment Phase.....	168
9.1.6.1. Post-infusion Period .....	168
9.1.6.2. Post-treatment Period (Cohorts A, B, and C).....	169
9.1.6.3. Lenalidomide Treatment Period after JNJ-68284528 Infusion (Cohort D).....	169
9.1.6.4. D+R Consolidation Treatment Period after JNJ-68284528 Infusion (Cohort E) .....	170
9.1.6.5. Long-term Follow-up.....	171
9.2. Efficacy Evaluations .....	171
9.2.1. Bone Marrow Examination for MRD Assessment .....	172
9.2.2. Myeloma Protein Measurements in Serum and Urine.....	172
9.2.3. Serum Calcium Corrected for Albumin .....	173
9.2.4. Bone Marrow Examination.....	173
9.2.5. Skeletal Survey.....	173
9.2.6. Documentation of Extramedullary Disease/ Extramedullary Plasmacytomas.....	174
9.2.7. Local Laboratory Assessments .....	174
9.3. Pharmacokinetics and Immunogenicity .....	175
9.3.1. Evaluations .....	175
9.3.2. Analytical Procedures .....	175
9.3.3. Pharmacokinetic Parameters .....	175
9.3.4. Immunogenicity Assessments/Antibodies to JNJ-68284528.....	176
9.4. Pharmacokinetic/Pharmacodynamic Evaluations .....	176
9.5. Biomarkers Evaluations .....	176
9.5.1. Pharmacodynamic/Predictive Markers .....	177
9.6. Patient-Reported Outcome Assessments.....	177
9.7. Safety Evaluations .....	179
9.8. Sample Collection and Handling.....	184
9.9. Medical Resource Utilization.....	184
<b>10. SUBJECT COMPLETION/DISCONTINUATION OF STUDY TREATMENT/ WITHDRAWAL FROM THE STUDY .....</b>	<b>185</b>
10.1. Completion .....	185
10.2. Discontinuation of Study Treatment.....	185
10.3. Withdrawal from the Study.....	186
10.4. Withdrawal from the Use of Research Samples .....	186
<b>11. STATISTICAL METHODS.....</b>	<b>186</b>
11.1. Subject Information .....	187
11.2. Sample Size Determination .....	187
11.3. Efficacy Analyses .....	187
11.4. Pharmacokinetic Analyses.....	189
11.5. Immunogenicity Analyses .....	189
11.6. Pharmacokinetic/Pharmacodynamic Analyses.....	190
11.7. Safety Analyses .....	190
11.8. Patient-reported Outcome Assessments .....	191
<b>12. ADVERSE EVENT REPORTING .....</b>	<b>191</b>
12.1. Definitions .....	191
12.1.1. Adverse Event Definitions and Classifications .....	191
12.1.2. Attribution Definitions.....	193
12.1.3. Severity Criteria .....	193
12.2. Special Reporting Situations.....	193
12.3. Procedures.....	194

12.3.1.	All Adverse Events.....	194
12.3.2.	Serious Adverse Events .....	195
12.3.3.	Adverse Events of Special Interest.....	197
12.3.4.	Pregnancy.....	198
12.4.	Contacting Sponsor Regarding Safety.....	198
<b>13.</b>	<b>PRODUCT QUALITY COMPLAINT HANDLING.....</b>	<b>198</b>
13.1.	Procedures.....	198
13.2.	Contacting Sponsor Regarding Product Quality .....	199
<b>14.</b>	<b>STUDY TREATMENT INFORMATION .....</b>	<b>199</b>
14.1.	Physical Description of Study Treatment.....	199
14.2.	Packaging .....	199
14.3.	Labeling.....	199
14.4.	Preparation, Handling, and Storage.....	199
14.5.	Drug Accountability .....	200
<b>15.</b>	<b>STUDY-SPECIFIC MATERIALS.....</b>	<b>200</b>
<b>16.</b>	<b>ETHICAL ASPECTS .....</b>	<b>201</b>
16.1.	Study-specific Design Considerations .....	201
16.2.	Regulatory Ethics Compliance .....	201
16.2.1.	Investigator Responsibilities .....	201
16.2.2.	Independent Ethics Committee or Institutional Review Board .....	202
16.2.3.	Informed Consent .....	203
16.2.4.	Privacy of Personal Data .....	204
16.2.5.	Long-term Retention of Samples for Additional Future Research.....	205
16.2.6.	Country Selection .....	205
<b>17.</b>	<b>ADMINISTRATIVE REQUIREMENTS .....</b>	<b>205</b>
17.1.	Protocol Amendments.....	205
17.2.	Regulatory Documentation .....	206
17.2.1.	Regulatory Approval/Notification .....	206
17.2.2.	Required Prestudy Documentation.....	206
17.3.	Subject Identification, Enrollment, and Screening Logs .....	207
17.4.	Source Documentation.....	207
17.5.	Case Report Form Completion .....	208
17.6.	Data Quality Assurance/Quality Control .....	208
17.7.	Record Retention .....	209
17.8.	Monitoring .....	209
17.9.	Study Completion/Termination.....	210
17.9.1.	Study Completion/End of Cohort .....	210
17.9.2.	Study Termination.....	210
17.10.	On-site Audits.....	211
17.11.	Use of Information and Publication .....	211
<b>REFERENCES.....</b>		<b>213</b>
<b>INVESTIGATOR AGREEMENT .....</b>		<b>261</b>
 <b>LIST OF ATTACHMENTS</b>		
Attachment 1:	Criteria for Response to Multiple Myeloma Treatment.....	217
Attachment 2:	Cytokine Release Syndrome ASBMT (ASTCT) Consensus Grading System .....	219
Attachment 3:	Immune Effector Cell-associated Encephalopathy (ICE) Tool.....	220
Attachment 4:	Immune Effector Cell-associated Neurotoxicity Syndrome (ICANS) ASBMT (ASTCT) Consensus Grading System <sup>a,b</sup> .....	221
Attachment 5:	International Myeloma Working Group Diagnostic Criteria .....	222

Attachment 6:	Prior Cancer Therapy for Multiple Myeloma .....	223
Attachment 7:	Eastern Cooperative Oncology Group Performance Status Grade .....	224
Attachment 8:	Formulas for Estimating Glomerular Filtration Rate .....	225
Attachment 9:	Serum Calcium Corrected for Albumin .....	226
Attachment 10:	Hepatitis B Virus Screening .....	227
Attachment 11:	Patient Reported Outcomes (PRO) Measures .....	228
Attachment 12:	Conversion Table for Glucocorticosteroid Dose .....	240
Attachment 13:	Anticipated Adverse Events .....	241
Attachment 14:	JNJ-68284528 Outpatient Administration Guidelines .....	242
Attachment 15:	Monitoring for Subjects Receiving JNJ-68284528 as Outpatient .....	244
Attachment 16:	Contraceptive and Barrier Guidance and Collection of Pregnancy Information .....	245
Attachment 17:	Handwriting Adverse Event Toxicity Grading Criteria .....	248
Attachment 18:	Anti-microbial Prophylaxis Recommendations .....	249
Attachment 19:	International Staging System (ISS) .....	250
Attachment 20:	COVID-19 Guidance on Study Conduct .....	251
Attachment 21:	Myeloma Frailty Score Calculator .....	253
Attachment 22:	Characterization of Imaging Response .....	254
Attachment 23:	Recommendations for the Reporting of Imaging Results .....	255
Attachment 24:	Asthma Guidelines .....	256
Attachment 25:	Antihistamines That May Be Used Pre-dose .....	260

## LIST OF IN-TEXT TABLES AND FIGURES

### TABLES

Table 1:	Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Study Procedures/Assessments .....	33
Table 2:	Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling .....	41
Table 3:	Cohort D: Time and Events Schedule for Study Procedures/Assessments .....	44
Table 4:	Cohort D: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling .....	52
Table 5:	Cohort E: Time and Events Schedule for Study Procedures/Assessments (Screening to Conditioning Regimen) .....	56
Table 6:	Cohort E: Time and Events Schedule for Study Procedures/Assessments (JNJ-68284528 Infusion to Post-treatment) .....	62
Table 7:	Cohort E: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling (Screening to Conditioning Regimen) .....	68
Table 8:	Cohort E: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling (JNJ-68284528 Infusion to Disease Progression or Study Completion) .....	69
Table 9:	Risks Associated with JNJ-68284528 and Mitigation Strategies .....	81
Table 10:	JNJ-68284528 Administration .....	135
Table 11:	Pre-infusion Medications .....	137
Table 12:	Criteria for Lenalidomide Administration after JNJ-68284528 Infusion – Cohort D .....	138
Table 13:	Dose Modification Guidelines for Lenalidomide .....	139
Table 14:	Lenalidomide Dose Reduction Steps .....	140
Table 15:	Lenalidomide Starting Dose After JNJ-68284528 Infusion for Subjects with Renal Impairment .....	140
Table 16:	D-VRd Induction Therapy .....	141
Table 17:	Dose modification Guidelines for Bortezomib, Lenalidomide, and Dexamethasone .....	145
Table 18:	Dose Modifications for Bortezomib .....	150
Table 19:	Dose Modifications for Lenalidomide .....	151
Table 20:	Dose Modifications for Dexamethasone .....	151
Table 21:	Criteria for Lenalidomide Administration after JNJ-68284528 Infusion – Cohort E .....	151
Table 22:	D+R Consolidation Therapy .....	152
Table 23:	Guidelines for the Management of Cytokine Release Syndrome .....	155
Table 24:	Guidelines for the Management of CAR-T Cell-related Neurotoxicity (eg, ICANS) .....	157
Table 25:	Guidelines for the Management of Raised ICP / Cerebral Edema <sup>a</sup> .....	159

**FIGURES**

Figure 1: LCAR-B38M Coding Region.....	76
Figure 2: Schematic Overview of the Study, Cohort A.....	88
Figure 3: Schematic Overview of the Study, Cohort B.....	90
Figure 4: Schematic Overview of the Study, Cohort C.....	92
Figure 5: Schematic Overview of the Study, Cohort D.....	95
Figure 6: Schematic Overview of the Study, Cohort E.....	97
Figure 7: Progression-free Survival According to MRD Status $10^{-5}$ in Studies 54767414MMY3003 and 54767414MMY3004.....	103

**PROTOCOL AMENDMENTS**

<b>DOCUMENT HISTORY</b>	
<b>Document</b>	<b>Date</b>
Amendment 3/USA-1	14 November 2020
Amendment 3	13 July 2020
Amendment 2	23 March 2020
Amendment 1	31 October 2019
Original Protocol	07 June 2019

**Amendment 3/USA-1 (14 November 2020)**

**Overall Rationale for the Amendment:** The overall reason for the amendment is to increase the sample size in Cohort A to obtain safety and efficacy data in subjects receiving ciltacabtagene autoleucl (cilta-cel) manufactured with the commercial process.

<b>Section number and Name</b>	<b>Description of Change</b>	<b>Brief Rationale</b>
Synopsis (Overview of Study Design); 1.2. Overall Rationale for the Study; 3.1. Overview of Study Design; 3.1.1. Cohort A Study Design  1.1.4. JNJ-68284528	The number of subjects enrolled in Cohort A was increased from 20 to 40. The size of all other cohorts remains unchanged.  Added that a similar manufacturing process will be used to generate commercial product in Cohort A of the study.	The sample size of Cohort A was increased to obtain safety and efficacy data in subjects receiving cilta-cel manufactured using the commercial process.
Time and Events Schedule (Table 7)	Deletion of footnotes	Correction of error. Footnotes are not applicable to assessments prior to cilta-cel infusion
Time and Events Schedule (Table 8)	Deletion of footnotes 'd' and 'e' from CyTOF/ PBMC/ Plasma (whole blood) at Study Completion assessment.	Correction of error

**Amendment 3 (13 July 2020)**

**Overall Rationale for the Amendment:** The overall reason for the amendment is to add a new cohort of subjects with multiple myeloma representing an additional patient population of unmet medical need.

<b>Section Number and Name</b>	<b>Description of Change</b>	<b>Brief Rationale</b>
Title page; Synopsis; 1. Introduction	Added generic name (ciltacabtagene autoleucl) to the document.	Added generic name
Synopsis (Overview of Study Design); 1.2. Overall Rationale for the	Added description of the study design for Cohort E	Addition of Cohort E



Section Number and Name	Description of Change	Brief Rationale
Study; 3.1. Overview of Study Design; 3.1.5. Cohort E Study Design		
Synopsis (Dosage and Administration); 6. Dosage and Administration; 6.1.5. Cohort E: Induction and Post JNJ-68284528 Therapy	Dosing information provided for D-VRd induction therapy and post JNJ-68284528 administration of daratumumab, lenalidomide, and dexamethasone.	
Time and Events Schedule	Time and Events Tables 5 to 8 added for Cohort E	
3.3. Study Design Rationale	Added a rationale for the design of Cohort E	
4.5. Cohort E Eligibility Criteria	Added inclusion and exclusion criteria for subjects to be enrolled in Cohort E	
6.1.1. Criteria for Apheresis (all cohorts unless otherwise specified)	Added timing of apheresis with respect to D-VRd induction treatment	
6.2.9. Cohort E: Management of Infusion-related Reactions and Local Injection-site Reactions of Daratumumab SC	Added management guidelines specific to daratumumab	
7. Treatment Compliance	Added D-VRd and D+R	
8. Prestudy and Concomitant Therapy	Added concomitant medication recording period for Cohort E	
8.3.1. Additional Prohibited Therapies for Cohort E	Added prohibited therapies with the use of bortezomib	
9.1.2. Screening Phase	Added that subjects in Cohort E will receive induction treatment with D-VRd after screening and prior to apheresis	
9.1.6.4. D+R Consolidation Treatment Period after JNJ-68284528 (Cohort E)	Added section for Cohort E post-treatment period	
9.5.1 Pharmacodynamic/Predictive Markers	Added that data review by the DMC will determine if there are any potential pharmacodynamic interactions between D-VRd and JNJ-68284528.	
9.7. Safety Evaluations	Added a subsection for Indirect Antiglobulin Test (IAT) to be performed before the first dose of daratumumab in Cohort E.	
10.2. Discontinuation of Study Treatment	Added discontinuation of study treatment information for Cohort E	
11.1. Subject Information	Added description of Cohort E analysis population	
Attachment 19: International Staging System (ISS)	Added Revised International Staging System (R-ISS)	
Attachment 21: Myeloma Frailty Score Calculator	Addition of Myeloma Frailty Score Calculator	
Attachment 24: Asthma Guidelines	Added asthma guidelines for daratumumab	
Attachment 25: Antihistamine That May Be Used Predose	Added antihistamines that may be used predose for daratumumab	

Section Number and Name	Description of Change	Brief Rationale
Synopsis (Overview of Study Design); 3.1. Overview of Study Design; 9.1.6.3. Lenalidomide Treatment Period after JNJ-68284528 Infusion (Cohort D) 17.9.1. Study Completion/End of Cohort	Cohort D (and Cohort E) will be considered complete <b>2 ½ years</b> after the last subject receives their initial dose of JNJ-68284528.	Clarification on definition of study completion
Synopsis (Statistical Methods)	Sample size language revised to be consistent with Section 11.2.	Revision for consistency
Synopsis (Dosage and Administration); 3.1.4. Cohort D Study Design (Figure 5)	One <b>or more cycles</b> of lenalidomide bridging therapy are permitted.	Revision made for clarity
Synopsis (Dosage and Administration); 3.1.4. Cohort D Study Design; 6.1.4. Lenalidomide (Cohort D)	Clarified that initiation of lenalidomide post-JNJ-68284528 infusion requires resolution of CRS or neurotoxicities <b>per ASTCT criteria</b> .	
Time and Events Schedule (Table 1)	Added (under screening assessments) PET/CT or MRI whole body scan from screening to apheresis for subjects without measurable disease in serum or urine. Moved from MRD assessment by imaging (optional)	
Time and Events Schedule (Table 2 and Table 4)	Aligned window for collection of biomarker samples at apheresis with hematology and chemistry sample collection ( $\leq 72$ hour window)	
Time and Events Schedule (Table 3)	Moved details of hematology collection in lenalidomide treatment period into footnote (w).	
Time and Events Schedule (Table 3)	Clarified that chemistry laboratory assessment should be performed at the start of each lenalidomide cycle during bridging.	
Time and Events Schedule (Table 3)	Added window for collection of hematology and chemistry on Day 1 with Cohorts A, B, and C.  Added window for bone marrow aspirate and extramedullary plasmacytomas at conditioning regimen.  Added MRD assessment by imaging at screening for subjects without measurable disease in urine or serum at baseline  Added window for assessment of extramedullary plasmacytomas	
3.1. Overview of Study Design	Added the following: For all cohorts, at select sites <del>within the US</del> and at investigator discretion with sponsor approval, study visits in the post-treatment part of the study, as early as after Day 100 after JNJ-68284528 infusion, may be performed remotely via telemedicine technology that connects study subjects to their research coordinators and	

Section Number and Name	Description of Change	Brief Rationale
	investigators <b>if not using the sponsor's telemedicine solution.</b>	
6.1.4. Lenalidomide (Cohort D)	Labeled Table 12 as Cohort D. Added cross reference to Table 15 in footnote.	
6.1.4. Lenalidomide (Cohort D)	Added that lenalidomide pill counts should be performed every cycle.	
14.5. Drug Accountability	Clarified that the investigator is responsible for ensuring that all <b>sponsor-provided</b> study treatment received at the site is inventoried and accounted for throughout the study.	
Attachment 1: Criteria for Response to Multiple Myeloma treatment	In footnote 'a' added the following: CR in such subjects indicates a normal FLC ratio of 0.26 to 1.65 ( <b>or reference range in testing laboratory</b> ) in addition to CR criteria listed above.	
Synopsis (Evaluations)	Added: For subjects with disease, but neither serum nor urine measurable disease, baseline PET/CT, or whole body MRI may be used to satisfy the measurable disease criteria.	Revisions for allow for PET/CT or whole body MRI to satisfy the measurable disease criteria.
Time and Events Schedule (Table 1)	For subjects with baseline measurable disease by PET/CT, added MRD assessment by imaging at screening, apheresis, and at 6 months, 12 months, and then yearly after infusion	
4.1.1. Cohort A Inclusion Criteria; 4.2.1. Cohort B Inclusion Criteria; 4.3.1. Cohort C Inclusion Criteria	Revised inclusion criterion to state that: For subjects with neither serum nor urine measurable disease, baseline PET/CT, or whole body MRI may be used to satisfy the measurable disease criteria	
9.2. Efficacy Evaluations	Added: Subjects who did not meet the definition of measurable disease per inclusion criterion and were enrolled in the study based on disease that is followed by PET/CT or whole body MRI must also follow the efficacy assessments for all other disease compartments (ie, serum, urine, bone marrow, skeletal (bone) lesions) as described in the Time and Events Schedule.	
9.2.6. Documentation of Extramedullary Disease/ Extramedullary Plasmacytomas	Added: These response criteria are applicable only to patients screened with measurable disease in serum or urine. For patients screened without measurable disease in serum or urine, via PET/CT or whole body MRI refer to Attachment 22 for response characterization.	
10.2. Discontinuation of Study Treatment	Revised bullet #4 as follows: Confirmed disease progression per IMWG criteria (Attachment 1) either between the time of conditioning therapy and infusion of JNJ-68284528, or during the post-JNJ-68284528 infusion <del>lenalidomide</del> treatment phase of Cohort D <b>with lenalidomide (as applicable) and Cohort E with daratumumab and lenalidomide. For subjects in Cohorts A, B, and C without measurable disease in serum or urine at screening, progression on either PET/CT scan or Whole Body MRI (see Attachment 22 for imaging response criteria) will be used.</b>	

Section Number and Name	Description of Change	Brief Rationale
11.3. Efficacy Analyses	<p>Revised MRD negative rate definition for subjects without measurable disease in Cohorts A, B, and C.</p> <p>For ORR added: For subjects with neither serum nor urine measurable disease and using PET/CT or whole body MRI to satisfy the measurable disease criteria (Cohorts A, B, and C), see Attachment 22 for adjudication of response criteria and Attachment 23 for reporting considerations.</p> <p>For PFS added: For subjects in Cohorts A, B, and C without measurable disease in serum or urine at screening disease, PFS is defined as the time from the date of the initial infusion of JNJ-68284528 to the date of disease progression as identified by imaging response criteria in Attachment 22 or death due to any cause, whichever occurs first.</p>	
Attachment 22: Characterization of Imaging Response	Addition of Characterization of Imaging Response via PET/CT or Whole Body MRI	
Attachment 23: Recommendations for the Reporting of Imaging Results	Addition of Recommendations for the Reporting of Imaging Results	
Synopsis (Overview of Study Design); 1.2. Overall Rationale for the Study; 3.1. Overview of Study Design; 4.2.1. Cohort B Inclusion Criteria	For Cohort B: Disease progression per IMWG criteria $\leq 12$ months after ASCT or disease progression $\leq 12$ months from the start of anti-myeloma therapy for subjects who have not had an ASCT. Confirmation may be from either central or local testing.	Correction to window
Time and Events Schedule (Table 1, Table 3)	Revised timing of skeletal survey to allow it to be performed at screening or $\leq 14$ days prior to the first dose of conditioning regimen.	Change made to allow for flexibility to allow skeletal survey to be performed in parallel with other disease evaluations.
Time and Events Schedule (Table 2)	Footnote 'g': deleted "For subjects in US" as PK and biomarker sample collection at the subject's home is not limited to only the US.	Correction
Time and Events Schedule (Table 3)	For Cohort D, added MRD assessment by imaging (optional) at screening.	Revision to align with common medical practice
Time and Events Schedule (Table 3)	Deleted duplicate information stating lenalidomide dosing is dependent on adequate hematologic recovery	Deletion of redundant information
4.3.1. Cohort C Inclusion Criteria	Criterion 7c revised to allow for transfusion within 3 days (reduced from 7 days) before hemoglobin and platelet testing.	Subjects in this cohort may have poor bone marrow reserve and require additional support
4.4.1. Cohort D Inclusion Criteria	Criterion 7d revised to state that women of childbearing potential must follow the contraception criteria outlined in the global REVLIMID® pregnancy prevention program <b>or equivalent local REMS as applicable in their region, whichever is more stringent.</b>	Clarification to inclusion criterion

Section Number and Name	Description of Change	Brief Rationale
4.1.2. Cohort A Exclusion Criteria; 4.2.2. Cohort B Exclusion Criteria; 4.3.2. Cohort C Exclusion Criteria; 4.4.2. Cohort D Exclusion Criteria	Added 'Contraindications' to exclusion criterion 16a (Cohort A), exclusion criterion 16b (Cohort B), exclusion criterion 15c (Cohort C), and exclusion criterion 14d (Cohort D).	Clarification to exclusion criteria
6. Dosage and Administration	For Cohort D, lenalidomide <b>given post JNJ-68284528</b> is also considered a study treatment.	Clarification on the definition of study treatment for Cohort D
6.1.4. Lenalidomide (Cohort D)	Consistent with the synopsis and Section 3.1.4, revised to state that subjects will receive lenalidomide at a dose of 10 mg per day upon adequate hematologic recovery from ASCT (absolute neutrophil count [ANC] $\geq 1 \times 10^9/L$ and platelet count $\geq 75 \times 10^9/L$ )	Clarification of lenalidomide dose after apheresis and prior to conditioning regimen
6.1.4. Lenalidomide (Cohort D)	In Table 14, Dose level -1; corrected starting dose (5mg) as Days 1-21 every <b>28 day cycle</b> .	Correction of error
12.1.1. Adverse Event Definitions and Classifications	For determination of the expectedness of an adverse event, the daratumumab Investigator's Brochure should be used in regions without marketing authorization. For dexamethasone, bortezomib, and lenalidomide with marketing authorization, expectedness of an adverse event will be determined by the prescribing information.	Revisions include daratumumab, lenalidomide, bortezomib, and dexamethasone
12.3.1. All Adverse Events	Revised language for exceptions	Revised for consistency with current template language
12.3.1. All Adverse Events; 12.3.3. Adverse Events of Special Interest	The sponsor assumes responsibility for appropriate reporting of RCL positive test results to regulatory authorities and will notify the investigator of positive RCL test results in a timely manner. RCL positive test results will not be collected as an adverse event in the eCRF.	Revision to clarify testing and reporting of replication competent lentivirus (RCL)
12.3.3. Adverse Events of Special Interest; 12.3.4. Pregnancy	Revised text to clarify that all subjects must adhere to the global REVLIMID® pregnancy prevention program or equivalent local lenalidomide pregnancy prevention program applicable in their region, whichever is more stringent	Revisions to provide additional guidance on lenalidomide Global Pregnancy Prevention Plan
Attachment 16: Contraceptive and Barrier Guidance and Collection of Pregnancy Information	Renamed attachment from "Definition of Woman of Childbearing Potential" to "Contraceptive and Barrier Guidance and Collection of Pregnancy Information". Added examples of contraceptives.	
12.3.1. All Adverse Events; Attachment 13: Anticipated Events	Deleted as anticipated events are considered not applicable for non-randomized studies	Correction in response to latest recommendations
Attachment 20: COVID-19 Guidance on Study Conduct	Added guidance on study conduct during the COVID-19 pandemic	COVID-19 Appendix
Throughout the protocol	Minor grammatical, formatting, or spelling changes were made.	Minor errors were noted

**Amendment 2** (23 March 2020)

**Overall Rationale for the Amendment:** The overall reason for the amendment is to add other neurotoxicities as a safety risk and implement additional monitoring and risk minimization measures for JNJ-68284528.

Section number and Name	Description of Change	Brief Rationale
Time and Events Schedule: Cohort A, Cohort B, and Cohort C (Table 1) Time and Events Schedule: Cohort D (Table 3)	Collection of a handwriting sample and frequency of testing was added to the study procedures. Study Day 35 was added to the table as additional assessment days for collection of handwriting samples.  Added that collection of medical history should include neurologic history  Added that ICE neurologic test should be performed at least until <b>ICANS is resolved</b> .  Added: Reporting of neurologic adverse events or exacerbation of existing neurologic adverse events will be extended to up to 12 months after JNJ-8284528 infusion	Update of safety information to include other neurotoxicities
1.1.5. Clinical Studies	Added to summary of data for Study 68284528MMY2001 a description of 6 cases of other neurotoxicities.	
1.3. Potential Safety Risks and Mitigation Strategies	Added other neurotoxicities to Table 5 (Risks Associated with JNJ-68284528 and Mitigation Strategies) as a subsection of neurologic toxicities	
2. Objectives and Endpoints	Added exploratory objective and endpoints to characterize potential early predictive markers for neurotoxicity	
4.1.2. Cohort A Exclusion Criteria (criterion 17a); 4.2.2. Cohort B Exclusion Criteria (criterion 17b); 4.3.2. Cohort C Exclusion Criteria (criterion 16c); 4.4.2. Cohort D Exclusion Criteria (criterion 17d)	Added the following to the list of serious underlying medical conditions:  Any history of Parkinson's disease or other neurodegenerative disorder	
6.1.3. JNJ-68284528 Administration	Table 6 – Added: 'Hospitalization is required for Grade 2, 3, or 4 CAR-T cell-related neurotoxicity (eg, ICANS) <b>temporally associated to CRS.</b> '  <b>Hospitalization for neurotoxicity that is not temporally associated with CRS, or any other neurologic adverse events, is at the discretion of the investigator.</b>	
6.2.2. Neurologic Adverse events	Divided section into overall neurologic adverse events with sub-sections of CAR-T cell related neurotoxicity (ICANS) (Section 6.2.2.1) and other neurotoxicities (Section 6.2.2.2).  Added that subjects should be monitored for neurotoxicity for 1 year post JNJ-68284528 infusion.	

Section number and Name	Description of Change	Brief Rationale
6.2.2.1. CAR-T Cell-related Neurotoxicity (ICANS)	<p>Added “Consider performing neuroimaging (eg, MRI)...”</p> <p>Clarified that hospitalization is required for Grade 2, 3, or 4 CAR-T cell-related neurotoxicity temporally associated to CRS.</p> <p>Added recommendation to rule out alternative etiologies at first sign of neurotoxicity.</p> <p>Table 13 (Guidelines for Management of ICANS) – added to row 1 (ICE score 7-9) when no concurrent CRS dexamethasone can be considered</p>	
6.2.2.2. Other Neurotoxicities	New section with guidance to monitor for other neurotoxicities	
9.7. Safety Evaluations	<p>Text updated to include additional guidance regarding the reporting of neurotoxicity events.</p> <p>To neurologic examination subsection, Added the following: For subjects with prior pertinent neurologic disease (eg, stroke, encephalitis) consider baseline MRI of the brain and an EEG.</p> <p>Added a section describing the assessment for qualitative changes in handwriting.</p>	
12.3.1. All Adverse Events	Added that changes in handwriting using criteria in Attachment 17 should be reported as an adverse event in the eCRF.	
Attachment 17: Handwriting Adverse Event Toxicity Grading Criteria	Table for added for qualitative grading of handwriting sample	
Time and Events Schedule: Cohort A, Cohort B, and Cohort C (Table 1) Time and Events Schedule: Cohort D (Table 3)	<p>Table 1 (footnote g) and Table 3 (footnote g), Added that Serology results performed as standard of care within 28 days prior to apheresis are acceptable</p> <p>Hepatitis B recommendation revised as follows: HBsAg, anti-HBc, anti-HBs, HBV DNA quantification (for subjects who are anti-HBs positive without history of vaccination or for subjects who are <b>anti-HBc positive with or without</b> anti-HBs positive <del>and anti-HBe positive</del>).</p>	Revisions to enrich the clinical trial population by allowing inclusion of subjects who are anti-HBc positive with or without anti-HBs positive
9.7. Safety Evaluations	Revisions made to Hepatitis B serology: Hepatitis B: HBsAg, anti-HBc, anti-HBs, HBV DNA quantification (for subjects who are anti-HBs positive without history of vaccination or for subjects who are <b>anti-HBc positive with or without</b> anti-HBs positive <del>and anti-HBe positive</del> )	

Section number and Name	Description of Change	Brief Rationale
Attachment 10: Hepatitis B Virus Screening	Revised hepatitis B screening guide to allow subjects who are anti-HBc positive only to participate in the study Added "If there is evidence of HBV reactivation initiate treatment as appropriate per institutional guidance".	
Synopsis	Subheadings added to dosage and administration to separate individual cohorts Repeated dosage information in Cohort D rather than cross referencing to another cohort to group information for clarity	Revisions made for clarity
Time and Events Schedule (Table 1, Table 3)	Added window to clarify timing of collection of hematology samples at apheresis For infectious disease testing, merged screening and apheresis columns and added collection window Clarified timing of sample collections for MRD MySim-Q changed from optional to 'unless translation not available'	
Time and Events Schedule (Table 1, Table 3)	For concomitant medications, added that collection is continuous '...until <b>at least 100</b> days...'. Also added "Thereafter, continue to report concomitant therapy given for any Adverse Events considered related to study drug until the end of the study".	
4.1.1. Cohort A Inclusion Criteria	Clarified criteria for lenalidomide refractory includes patients with non-responsive disease while on lenalidomide for at least 1 cycle	
4.3.2. Cohort C Exclusion Criteria	Revised prior antitumor therapy as follows: Targeted therapy, epigenetic therapy, or treatment with an investigational drug or used an invasive investigational medical device within 14 days <del>or at least 5 half-lives, whichever is longer</del>	
7. Treatment Compliance	Added that site specific methods to ensure compliance with outpatient medication is permissible.	
9.5. Biomarkers Evaluations	Defined that additional biomarker samples include but are not limited to serum or PBMCs from whole blood. Clarified that a tumor sample should be collected from subjects diagnosed with a SPM.	
9.5. Biomarkers Evaluations	Added that if a subject dies and an autopsy is preformed, specimens may be requested by the sponsor for analysis.	
9.9. Medical Resource Utilization	Added that costs associated with the medical encounters will be collected separately from the eCRF. All health economic data will be used only in a de-identified manner.	
9.7. Safety Evaluations; 12.1.3. Safety Criteria	Defined grading criteria for ICANS and CRS.	
11.3. Efficacy Analyses	Added: A sensitivity analysis of overall MRD negative rate will be performed based on the subjects in the mITT analysis set who received the JNJ-68284528 product that met all of the pre-specified release criteria	



Section number and Name	Description of Change	Brief Rationale
Time and Events Schedule (Table 1)	Revised collection of disease characteristics from screening phase, to occur between screening and the conditioning regimen.	Change made to avoid additional bone marrow collections
Time and Events Schedule (Table 1, Table 3)	Day 21 visit window changed from $\pm 1$ day to $\pm 2$ days for consistency with the PK/biomarker assessments in Tables 2 and 4.	Revised visit window for consistency
Time and Events Schedule (Table 3)	Revised language for pregnancy testing during lenalidomide treatment: to be 'repeated' <del>and repeated at a maximum of</del> at least every 28 days.	Revised pregnancy testing guidelines to be consistent with lenalidomide REMS
4.4.1. Cohort D Inclusion Criteria	Criterion 7d revised to state that subject must agree to practice 2 methods of reliable birth control simultaneously from <b>4 weeks prior to initiating treatment with lenalidomide</b> <del>the time of signing the ICF</del> until 1 year after receiving JNJ-68284528 infusion	
Attachment 16: Definition of Woman of Childbearing Potential	Postmenopausal state definition changed from 12 months with no menses to 24 months with no menses	
Time and Events Schedule (Table 3)	Added serum FLC and serum/urine immunofixation, MRD (bone marrow aspirate) collections and assess extramedullary plasmacytomas at conditioning regimen	Change for consistency with other cohorts
4.3.1. Cohort C Exclusion Criteria	Criterion 3c modified as follows: Targeted therapy, epigenetic therapy, or treatment with an investigational drug or used an invasive investigational medical device within 14 days <del>or at least 5 half lives, whichever is longer.</del>	
Time and Events Schedule (Table 4)	Added collection of PBMCs for functional in vitro assays	Correction of error
4.1.2. Cohort A Exclusion Criteria; 4.2.2. Cohort B Exclusion Criteria; 4.3.2. Cohort C Exclusion Criteria 4.4.2. Cohort D Exclusion Criteria	Localized prostate cancer with a Gleason score of 6 corrected to "Gleason score of $\leq 6$ ".	
Time and Events Schedule (Cohort D, Table 4)	Added collection of PBMCs for functional in vitro assays (omitted in error in Amendment 1)	
9.1.6.2. Post-treatment Period (Cohorts A, B, and C)	Paper PROs will be used in this study. Deleted the following: "PRO assessments may be captured via smart phone based application and electronic patient reported outcomes (ePRO) instruments".	
1.1.5. Clinical Studies	Added an update to Study 68284528MMY2002	Updated study information added
1.3. Potential Safety Risks and Mitigation Strategies (Table 5)	Added that sponsor should be notified if a subject is experiencing Grade 2 or higher CRS  Added that sponsor should be notified if a subject is experiencing any grade ICANS	Revised risk language

Section number and Name	Description of Change	Brief Rationale
1.3. Potential Safety Risks and Mitigation Strategies (Table 5); 6.2.7. Infections	Added that JNJ-68284528 should not be administered to subjects with active infections Added recommendation for extended use of anti-microbial therapies	Revised guidance for infections
Attachment 18	Added table with anti-microbial prophylaxis recommendations	
3.1. Overview of Study Design	Data Monitoring Committee (DMC) added for periodic review of all safety data.	Data Monitoring Committee added
3.1. Overview of Study Design; 9.1.6.2. Post-treatment Period (Cohorts A, B, and C)	Added that 'with sponsor approval' study visits in the post-treatment part of the study 'as early as' after Day 100 may be performed remotely via telemedicine technology.	Revisions to allow telemedicine visits to occur earlier for some subjects
3.1.1. Cohort A Study Design; 3.1.2. Cohort B Study Design; 3.1.3. Cohort C Study Design; 3.1.4. Cohort D Study Design	Revised bridging therapy options to include therapies to which a subject has not been previously exposed.	Revisions to bridging therapy recommendations
Time and Events Schedule (Cohort D, Table 3); Synopsis (Overview of Study Design); 3.1.4. Cohort D Study Design	Added that alternative bridging therapy instead of, or in addition to, lenalidomide is permissible with sponsor approval.	
Time and Events Schedule (Cohort D, Table 3);	Added that Cycle 2 of bridging therapy is permitted with sponsor approval.	
3.1.5. Retreatment with JNJ-68284528	Bridging therapy is allowed for subjects receiving retreatment with permission from the sponsor.	
Synopsis (Dosage and Administration); 3.1.4. Cohort D Study Design; 6.1.4. Lenalidomide (Cohort D)	The first 5 subjects in Cohort D will be treated with JNJ-68284528 only, there will be no lenalidomide after JNJ-68284528. The DMC will review the data from these 5 subjects and recommend if treating with lenalidomide should proceed. Once dosing of subjects with lenalidomide is initiated, a stagger of at least 4 weeks between subjects will continue for the next 5 subjects.	Revised staggered dosing strategy for Cohort D
6.1.4. Lenalidomide (Cohort D)	Added: <b>Initiation and continuation of lenalidomide treatment is dependent on no additional safety concerns by investigator or sponsor</b>	
6.1.3. JNJ-68284528 Administration (All Cohorts)	Target dose statement in Table 6 revised as follows: The target dose will be the RP2D <del>which is anticipated to be</del> of $0.75 \times 10^6$ CAR-positive viable T cells/kg (range: $0.5-1.0 \times 10^6$ CAR-positive viable T cells/kg) as described in Section 3.3	Clarification that RP2D is known (no longer anticipated to be)

Section number and Name	Description of Change	Brief Rationale
1.3. Potential Safety Risks and Mitigation Strategies (Table 5); 6.2.5. Cytopenia	Added consideration of parvovirus B19 monitoring	Additional guidance for cytopenia and hypogammaglobulinemia
1.3. Potential Safety Risks and Mitigation Strategies (Table 5); 6.2.6. Hypogammaglobulinemia	Added that subjects IgG <400 mg/dL and recurrent infections may receive prophylactic IVIG	
9.1.6.3. Lenalidomide Treatment Period after JNJ-68284528 Infusion (Cohort D)	Lenalidomide treatment period lasts until study completion defined as: <del>2 years</del> after the last subject in Cohort D has discontinued <b>lenalidomide for 4 weeks or has had 2 years of follow-up after receiving</b> <del>received</del> his or her initial dose of JNJ-68284528, <b>whichever is later</b>	Revised definition of lenalidomide treatment period.
9.2.7. Local Laboratory Assessments;  9.2. Efficacy Evaluations  Time and Events Schedule (Table 1 and Table 3),	Added a section to allow for local laboratory data to be collected if central laboratory data is not available.  Added footnote 's' (Table 1) and footnote 'u' (Table 3)	Addition to allow for flexibility in collecting laboratory data
Attachment 14 (JNJ-68284528 Outpatient Administration Guidelines)	Added that sponsor approval is required when evaluating for the suitability of outpatient administration	Revised for consistency
Throughout the protocol	Minor grammatical, formatting, or spelling changes were made.	Minor errors were noted

### Amendment 1 (31 October 2019)

**Overall Rationale for the Amendment:** The overall reason for the amendment is to add 2 new cohorts of subjects with multiple myeloma representing different populations of unmet medical need.

Section number and Name	Description of Change	Brief Rationale
Synopsis, Overview of Study Design; 1.2. Overall Rationale for the Study; 3.1. Overview of Study Design; 3.1.3. Cohort C Study Design	Added description of the study design for Cohort C	Addition of Cohort C, which consists of subjects with relapsed refractory disease previously treated with a PI, IMiD, anti-CD38 monoclonal antibody, and BCMA-directed therapy.
Time and Events Schedule	Cohort C assessments to be collected per Table 1 and Table 2	
3.3. Study Design Rationale	Added rationale for the design of Cohort C	
4.3. Cohort C Eligibility Criteria	Added inclusion and exclusion criteria for subjects to be enrolled in Cohort C	

<b>Section number and Name</b>	<b>Description of Change</b>	<b>Brief Rationale</b>
Synopsis, Overview of Study Design; 1.2. Overall Rationale for the Study; 3.1. Overview of Study Design; 3.1.4. Cohort D Study Design	Added description of the study design for Cohort D	Addition of Cohort D, which consists of subjects with recently diagnosed multiple myeloma who did not achieve CR after 4-8 total cycles of initial therapy, including induction, high-dose chemotherapy, and ASCT with or without consolidation.
Synopsis, Dosage and Administration;	Subjects in Cohort D will receive 1 cycle of lenalidomide after apheresis and before conditioning regimen. At least 21 days post infusion of JNJ-68284528, and after adequate hematologic recovery, subjects will receive lenalidomide treatment.	
Time and Events Schedule	Time and Events Tables 3 and 4 added for Cohort D	
3.1.5. Retreatment with JNJ-68284528	Added "Subjects in Cohort D will not be allowed to receive retreatment".	
3.3. Study Design Rationale	Added rationale for the design of Cohort D	
4.4. Cohort D Eligibility Criteria	Added inclusion and exclusion criteria for subjects to be enrolled in Cohort D	
6.1.4. Lenalidomide (Cohort D)	Section added to provide dosing information for lenalidomide.	
7. Treatment Compliance	Added that pill counts will be used to assess compliance to lenalidomide	
8.2. Permitted Medications	Added cross reference to lenalidomide prescribing information for subjects in Cohort D	
8.3. Prohibited Therapies	Added cross reference to lenalidomide prescribing information for subjects in Cohort D	
9.1.7.2. Post-treatment Period (Cohort D)	Added separate description for the post-treatment period for Cohort D.	
10.2. Discontinuation of Study Treatment	Added that criteria apply to discontinuation of lenalidomide.	
12.3.3. Adverse Events of Special Interest	Added reference to the REMS program for lenalidomide with regards to the embryo-fetal risks to ensure compliance with the guidance on subject education	
Synopsis, Dosage and Administration; 3.1.1. Cohort A Study Design; 3.1.2. Cohort B Study Design; 3.1.3. Cohort C Study Design; 3.1.4. Cohort D Study Design; 3.2. Rationale of Dose and Administration Schedule Selection; 6.1.2. Administration of Conditioning Regimen; 9.1.4. Cyclophosphamide and Fludarabine Conditioning Regimen	Added "The dose of fludarabine should be reduced to 24 mg/m <sup>2</sup> for subjects with an estimated glomerular filtration rate of 30 to 70 mL/min/1.73m <sup>2</sup> ."	

<b>Section number and Name</b>	<b>Description of Change</b>	<b>Brief Rationale</b>
Synopsis, Statistical Methods; 11.2. Sample Size Determination	Deleted “For Cohort A and Cohort B, an MRD negative rate of less than 10% is deemed not clinically meaningful. With 20 subjects, the lower bound of 90% CI will be $\geq 10\%$ if more than 4 subjects achieve MRD negativity.”	Removed language referring to specific MRD negative rates in individual cohorts as sample size is not driven by statistics.
Synopsis, Study Hypotheses; 2.2. Hypotheses	Removed specific statistical language for MRD negative rate.	Aligned hypothesis statement with changes to statistical methods section
Time and Events Schedule (Table 1); 9.7. Safety Evaluations	Added an ocular examination for subjects in Cohort C who received prior ADC treatment Footnote ‘r’ specifies requirements of ocular examination.	Addition for safety monitoring
Time and Events Schedule (Table 1); 6.1.3. JNJ-68284528 Administration (All Cohorts); 9.1.7. Post-treatment Phase; 16.1. Study-specific Design Considerations; Attachment 14; Attachment 15	Added guidance for screening subjects for outpatient suitability and monitoring for subjects receiving JNJ-68284528 as an outpatient	Addition of guidance to evaluate subjects for outpatient suitability
Time and Events Schedule (Table 1); 2.1. Objectives and Endpoints; 9.6. Patient Reported Outcomes; 11.8. Patient-reported Outcome Assessments; Attachment 11	The EQ-5D-5L measure was replaced with the Multiple Myeloma Symptom and Impact Questionnaire (MySIm-Q). The MySIm-Q is an optional assessment.  Added endpoint for worsening of symptoms using MySIm-Q	Change made in response to evolving PRO strategy
Time and Events Schedule (Table 1); 2.1. Objectives and Endpoints; 9.9. Medical Resource Utilization	MRU assessments will be collected for subjects in all cohorts	Addition for consistency with other studies in the program
Time and Events Schedule (Table 1)	Added $\leq 24$ -hour window to hematology and chemistry assessments collected pre-dose (JNJ-68284528 infusion) and for assessment criteria to be collected prior to JNJ-68284528 infusion.  Windows added for assessment of extramedullary plasmacytomas  Footnote ‘i’ (vital signs): added window of $\pm 10$ minutes	Addition for consistency with other studies in the program
Time and Events Schedule (Table 2)	Added a window of $\leq 4$ hours for pre-dose PK and biomarker samples not previously defined with collection window	

<b>Section number and Name</b>	<b>Description of Change</b>	<b>Brief Rationale</b>
Time and Events Schedule (Table 1)	<p>Frequency of serology testing expressed in weeks instead of months</p> <p>Added that adverse events considered related to study drug need to be reported until the end of study</p> <p>Footnote 'g': added additional guidance around testing for subjects who are positive for antibodies to hepatitis B and subjects with HBV vaccination.</p>	Revisions incorporated to harmonize language within this protocol and across studies
4.1.1. Cohort A Inclusion Criteria	Criterion 1a: Revised language to clarify lenalidomide refractory requirement	
4.1.1. Cohort A Inclusion Criteria; 4.2.1. Cohort B Inclusion Criteria	Criterion 7a and 7b: Replaced laboratory criteria of 'creatinine clearance' with 'estimated glomerular filtration rate'	
4.1.2. Cohort A Exclusion Criteria; 4.2.2. Cohort B Exclusion Criteria	<p>Revised criterion 15a and 15b regarding requirement of supplemental oxygen</p> <p>Criterion 17a and 17b: clarified that bacterial infection 'requiring systemic antimicrobial therapy' as a serious underlying medical condition.</p>	
6.1.1. Criteria for Apheresis; 6.1.2.1 Criteria for Conditioning Regimen	Harmonized language across JNJ-68284528 protocols has been incorporated	
9.7. Safety Evaluations	<p>Added "Beyond the adverse event reporting period, adverse events that are considered related to study drug need to be reported until the end of the study"</p> <p>Added guidance regarding collection of follow-up data for Grade 3 or higher laboratory abnormalities</p>	
Time and Events Schedule (Table 1)	Day 56 MRD evaluation added	To harmonize with bone marrow collections for PK and biomarkers
Time and Events Schedule (Table 1)	During the JNJ-68284528 infusion period, post-infusion period, and post-treatment phase added that pregnancy testing should be performed as clinically indicated 'or as mandated by local regulations, whichever is more stringent'	Revision to pregnancy testing to accommodate local regulations
Time and Events Schedule (Table 1)	Timing of the MRD (bone marrow aspirate) assessment changed from Day 28 to Day 56	Change made to more accurately capture the depth of response
Time and Events Schedule (Table 1)	Added PRO assessments (EORTC QLQ-C30, MySiM-Q, and PGIS) at apheresis	Assessment at apheresis added in order to obtain 2 timepoints before subjects receive treatment.
Time and Events Schedule (Table 2)	Footnote 'h' (whole blood immunophenotyping) and 'i' (flow cytometry, bone marrow) to specify when additional samples may be collected at the discretion of the investigator.	Footnotes added for guidance regarding additional sample collections

Section number and Name	Description of Change	Brief Rationale
Synopsis, Dosage and Administration; 6.1.4. Lenalidomide (Cohort D)	Added that subjects will continue to receive lenalidomide for 2 years post JNJ-68284528 infusion	Duration of lenalidomide treatment added.
Synopsis, Overview of Study Design; 1.2. Overall Rationale for Study Design	Added that subjects who have received prior therapy that is targeted to BCMA are excluded from Cohorts A and B; and that subjects who met eligibility criteria for Cohort A and Cohort B must be enrolled in Cohort B.	Additions for clarity
Time and Events Schedule (Table 1)	Table 1, added entry for criteria for JNJ-68284528 administration	
Time and Events Schedule (Table 1)	Added footnote 'e' to clarify the collection window for disease characteristics cytogenetics.	
Time and Events Schedule (Table 1)	The placement of footnote 'k' for quantitative immunoglobulins was updated as it applies to the entire duration of the study.  Added footnote 'q' to indicate that additional immunoglobulin samples may be collected as clinically indicated for safety	
Time and Events Schedule (Table 1)	The description of extramedullary Plasmacytomas was updated to: 'By physical examination (if applicable): measurable sites at D28, D56, D78, D100 and then every 4 weeks. By radiologic imaging: D78, D156 then every 12 weeks (for all subjects with a history of plasmacytomas or as clinically indicated for others)' <del>Measurable sites Day 28, Day 56, Day 78, Day 100 then every 4 weeks for physical examination (if applicable) and Day 78 and Day 156 then every 12 weeks for radiologic assessment (for subjects with a history of plasmacytomas or as clinically indicated for others).</del>	
Time and Events Schedule (Table 1); 1.3. Potential Safety Risks and Mitigation Strategies; 6.2.4. Second Primary Malignancy; 9.1.7.3. Long-term Follow-up	Added language to indicate that a tumor sample should be collected for any second primary malignancies in text and footnote 'p'.	
Time and Events Schedule (Table 1); 9.1.7.1. Post-treatment Period (Cohorts A, B, and C)	Added to survival follow-up that survival status will also be collected prior to the any planned efficacy analysis	
1.3. Potential Safety Risks and Mitigations	Added cross references from Table 5 to management guidelines in the document	
3.1.1. Cohort A Study Design; 3.1.2. Cohort B Study Design	Removed 'anticipated to be' from the recommended Phase 2 dose as the dose has now been established.	

Section number and Name	Description of Change	Brief Rationale
4.1.2. Cohort A Exclusion Criteria; 4.2.2. Cohort B Exclusion Criteria	Criterion 5a and 5b modified to state “Ongoing” toxicity from previous anticancer therapy...	
6.1.3.1. Exceptional Release Criteria	Added text stating that the investigator will discuss with the subject if product does not meet pre-specified release criteria	
6.2.7. Infections	Clarified that HBV reactivation has occurred “in subjects receiving other CAR-T products”	
6.2.8. Hypersensitivity Reactions	Added reference to lenalidomide prescribing information for subjects in Cohort D	
8.3. Prohibited Therapies	Added that ‘systemic’ corticosteroids’ should be avoided.	
9.2.1. Bone Marrow Examination for MRD Assessment	Added “In case the myeloma clone is not be identified successfully from the baseline fresh bone marrow aspirate, the sponsor will ask for non-decalcified diagnostic tissue.”	
9.7. Safety Evaluations	Added for Cohort D: adverse events will be collected 100 days after JNJ-68284528 infusion or 30 days after the last dose of lenalidomide (whichever is later).	
12.1.1. Adverse Event Definitions and Classifications	Clarified definition of a serious adverse event as inpatient hospitalization <b>that was not required by the protocol.</b>	
12.3.1. All Adverse Events; 12.3.3. Adverse Events of Special Interest	Clarified the time period for reporting adverse events and serious adverse events as 100 days after infusion of JNJ-68284528 (Cohorts A, B, and C), and 100 days after infusion of JNJ-68284528 or 30 days after the last dose of lenalidomide, whichever is later (for Cohort D)	
12.3.2. Serious Adverse Events	Added, for clarity: Events that require an escalation of care when the subject is already hospitalized should be recorded as an SAE. Examples of such events were added in text.	
Throughout the document	Revised reference to ‘Cohort A and Cohort B’ to state ‘all Cohorts’ where applicable or reference to specific individual Cohorts, as applicable.	
Time and Events Schedule (Table 1)	PRO-CTCAE is no longer optional.	Updates to PRO-CTCAE
Time and Events Schedule (Table 1); 9.7. Safety Evaluations; 12.1.3. Severity Criteria; Attachment 2; Attachment 4	Updated reference for ASBMT criteria and added incorporated change in name (to ASTCT).	Update to ASBMT criteria
Time and Events Schedule (Table 2)	Added footnote ‘g’ to state that PK and biomarker collections may be performed at the subject’s home by mobile study personnel	Revisions to include the capability for nursing home visits



<b>Section number and Name</b>	<b>Description of Change</b>	<b>Brief Rationale</b>
3.1. Overview of Study Design; 9.1.7.1. Post-treatment period (Cohorts A, B, and C)	Added that, at the discretion of the investigator, study visits in the post treatment phase may be performed remotely via telemedicine technology and that blood sample collection may be performed in the subject's home by mobile study personnel. PRO assessments may be captured via smart-phone based application and ePRO instruments.	
1.1.5. Clinical Studies	Updated summary of preliminary data for Study 68284528MMY2001  Added a brief summary of preliminary data for Study 68284528MMY2002  Consolidated summary of data from the Legend-2 study	Updated data provided for ongoing studies
1.3. Potential Safety Risks and Mitigation Strategies	In Table 5, additional guidance added for CRS, SPM, cytopenias, and hypogammaglobulinemia	Updates in response to evolving safety information
3.1.1 Cohort A Study Design; 3.1.2 Cohort B Study Design	Guidance provided on dose reduction for fludarabine in subjects with renal dysfunction.	
6.2.5. Cytopenia	Added text stating that "severe thrombocytopenia may increase the risk of bleeding"  Added precaution for neutropenia and thrombocytopenia for subjects in Cohort D upon receiving lenalidomide after JNJ-68284528	
6.2.6. Hypogammaglobulinemia	Added that vaccination with live virus is not permitted for at least 4 weeks prior to the start of conditioning regimen "and for 100 days after infusion of JNJ-68284528"	
3.1.2. Cohort B Study Design	Deleted text stating that the medical monitor must be contact before administration of bridging therapy and replaced with more specific guidance regarding when bridging therapy will be allowed	Guidance added for use of bridging therapy in Cohort B
3.1.5. Retreatment with JNJ-68284528	Revised criteria to state that to be eligible for retreatment the subject must have no ongoing Grade 2 non-hematologic toxicity (except for nausea, vomiting, hair loss) and no ongoing Grade 3 or higher hematologic toxicity.	Refined retreatment criteria
3.2. Rationale of Dose and Administration Schedule Selection	Added brief summary of data from Study 68284528MMY2001 used to form the decision of the Safety Evaluation Team	Recommended Phase 2 dose has been determined

Section number and Name	Description of Change	Brief Rationale
3.3. Study Design Rationale	Deleted “Further, the sponsor will assess whether a fixed-timepoint at 1 year for evaluation of MRD negativity is as predictive of long-term outcome as the overall MRD negative rate.”	Correction to MRD analysis
4.1.1. Cohort A Inclusion Criteria	Criterion 1a – deleted text for smoldering myeloma Criterion 2a – revised evidence of progressive disease within 6 months of the last regimen Criterion 8a – correction to timing of pregnancy testing (deleted prior to first dose of conditioning regimen)	Revision for consistency in language with protocols from other JNJ-68284528 studies
4.2.1. Cohort B Inclusion Criteria	Criterion 1b - removed mention of smoldering myeloma Criterion 8b – correction to timing of pregnancy testing (deleted prior to first dose of conditioning regimen)	
4.1.1. Cohort A Inclusion Criterion; 4.1.2. Cohort A Exclusion Criteria; 4.2.1. Cohort B Inclusion Criterion; 4.2.2. Cohort B Exclusion Criterion	The amount of time that a subject must agree to remain on a highly effective method of contraception, planning to become pregnant, or father a child, changed from ‘at least 100 days’ to ‘1 year’ after receiving an infusion of JNJ-68284528	Updates to the length of time subjects must remain on a highly effective method of contraception
4.1.2. Cohort A Exclusion Criteria; 4.2.2. Cohort B Exclusion Criteria	Criteria 3a revised with updated list of exceptions to active malignancies.	Revision to be consistent with current template language
6.1.3.1. Exceptional Release Criteria	Added that notification and approval for use of the product will be obtained “in compliance with local regulations Changed “investigators <b>should</b> inform the study subject” to “investigators <b>will</b> inform the study subject” Deleted text stating that subjects must sign an ICF to consent to receive product as that is no longer a requirement	Corrections to exceptional release criteria
8. Prestudy and Concomitant Therapy	Added requirements for Cohorts C and D	Revision to accommodate the addition of new cohorts
9.1.1 Overview	Updated volume of blood required for all cohorts	
9.1.6.2. Post-treatment Period	Specified in the section header that language in this section is applicable to Cohorts A, B, and C	
8. Prestudy and Concomitant Therapy	A list of medications given for adverse events (to be recorded) has been added	Added guidance to concomitant medications that should be recorded
9.1.7.1. Post-treatment Phase	Removed text that is repetitive with other sections of the protocol	Consolidated text to remove redundancy

<b>Section number and Name</b>	<b>Description of Change</b>	<b>Brief Rationale</b>
11.3. Efficacy Analyses	Added "For time-to-event endpoints, such as DOR, PFS, and OS, the distributions will be provided using Kaplan-Meier estimates. Detailed analysis methods will be provided in the Statistical Analysis Plan."	Added methods for analysis of time-to-event endpoints
12.3.1. All Adverse Events	Added that all deaths at any time not related to disease progression occurring at any time of the study should be reported to the sponsor following expedited reporting procedures.	Revised in response to health authority feedback
12.3.3. Adverse Events of Special Interest	Added that the following events must be reported to the sponsor using the Serious Adverse Event Form within 24-hours of awareness: $\geq$ Grade 3 CRS, $\geq$ Grade 3 neurotoxicity, $\geq$ Grade 3 tumor lysis syndrome second primary malignancies	
Attachment 16	Woman of Childbearing Potential and Woman of Not of Childbearing Potential has been defined in detail	Provided a definition of women of childbearing potential
Throughout the protocol	Minor grammatical, formatting, or spelling changes were made.	Minor errors were noted

## SYNOPSIS

A Phase 2, Multicohort Open-Label Study of JNJ-68284528, a Chimeric Antigen Receptor T cell (CAR-T) Therapy Directed Against BCMA in Subjects with Multiple Myeloma

JNJ-68284528 (ciltacabtagene autoleucl) is an autologous chimeric antigen receptor T cell (CAR-T) therapy that targets B cell maturation antigen (BCMA), a molecule expressed on the surface of mature B-lymphocytes and malignant plasma cells. The JNJ-68284528 drug product used in this study and the LCAR-B38M CAR-T cell drug product used in the first-in-human Legend-2 study express an identical CAR protein. The JNJ-68284528 drug product will be produced using modified manufacturing and scale-up processes. Results from the Phase 1b portion of Study 68284528MMY2001 and the Legend-2 study indicate that JNJ-68284528 and LCAR-B38M CAR-T cells have significant anti-myeloma activity and a safety profile consistent with the known mechanism of action of the product.

## PRIMARY OBJECTIVE, ENDPOINT

Objective	Endpoint
<b>Primary</b>	
<ul style="list-style-type: none"> <li>To evaluate the overall minimal residual disease (MRD) negative rate of subjects who receive JNJ-68284528</li> </ul>	<ul style="list-style-type: none"> <li>MRD negative rate (<math>10^{-5}</math> threshold) as defined by the International Myeloma Working Group (IMWG) criteria using next generation sequencing (NGS)</li> </ul>

## STUDY HYPOTHESES

The primary hypothesis is that JNJ-68284528 will induce a deep-response, measured by MRD negative rate in the clinical settings investigated.

## OVERVIEW OF STUDY DESIGN

This is a Phase 2, multicohort, open-label, multicenter study to determine whether treatment with JNJ-68284528 (alone or with other treatment regimens) results in MRD negativity in adult subjects with multiple myeloma. Multiple patient populations of unmet medical need will be studied. Approximately 40 subjects will be enrolled in Cohort A, and approximately 20 subjects each will be enrolled in all other cohorts. The primary endpoint for all cohorts will be overall MRD negative rate at a  $10^{-5}$  threshold.

Subjects will be enrolled into one of the following cohorts based on eligibility criteria.

- Cohort A: progressive disease after 1 to 3 prior lines of therapy, including a proteasome inhibitor (PI) and immunomodulatory drug (IMiD) either individually or in combination. Subjects are required to be refractory to lenalidomide. Subjects who have received prior therapy that is targeted to BCMA are excluded from this cohort.
- Cohort B: one line of previous therapy containing a PI and an IMiD and early relapse defined as disease progression  $\leq 12$  months after an autologous stem cell transplantation (ASCT) or  $\leq 12$  months after the start of front-line therapy for subjects who have not had an ASCT. Subjects who have received prior therapy that is targeted to BCMA are excluded from this cohort.
- Cohort C: relapsed or refractory disease in subjects previously treated with a PI, IMiD, anti-CD38 monoclonal antibody, and BCMA-directed therapy (excluding cellular immunotherapy).
- Cohort D (JNJ-68284528 plus lenalidomide): multiple myeloma without complete response after 4 to 8 total cycles of initial therapy, including induction, high-dose chemotherapy and ASCT with or without consolidation

- Cohort E (daratumumab, bortezomib, lenalidomide, and dexamethasone [D-VRd] induction, JNJ-68284528, then daratumumab and lenalidomide [D+R]): high risk newly diagnosed and untreated multiple myeloma (hr-NDMM) subjects for whom stem cell transplant is not planned as initial therapy.

Subjects who meet the eligibility criteria for Cohort A and Cohort B, must be enrolled in Cohort B.

All enrolled subjects will undergo apheresis to acquire peripheral blood mononuclear cells (PBMCs). JNJ-68284528 will be generated from the subject's T cells selected from the apheresis product. After JNJ-68284528 production and product release, subjects will receive a conditioning regimen of cyclophosphamide and fludarabine. JNJ-68284528 will be administered 5 days to 7 days after the start of the conditioning regimen. The primary analysis for each cohort will occur approximately 1 year after the last subject in each cohort has received his or her initial dose of JNJ-68284528.

Cohorts A, B, and C will each be considered complete after the last subject has had 2 years of follow-up after the initial dose of JNJ-68284528.

Cohort D and Cohort E will be considered complete 2 ½ years after the last subject receives their initial dose of JNJ-68284528.

## DOSAGE AND ADMINISTRATION

### *Cohort A, Cohort B, and Cohort C*

All subjects in Cohorts A, B, and C will receive a conditioning regimen consisting of cyclophosphamide 300 mg/m<sup>2</sup> intravenously (IV) daily and fludarabine 30 mg/m<sup>2</sup> IV daily for 3 days. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an estimated glomerular filtration rate (eGFR) of 30 to 70 mL/min/1.73m<sup>2</sup>. JNJ-68284528 IV infusion will take place 5 to 7 days after the start of the conditioning regimen. The target dose is 0.75 x 10<sup>6</sup> CAR-positive viable T cells/kg (range: 0.5-1.0 x 10<sup>6</sup> CAR-positive viable T cells/kg).

### *Cohort D*

Subjects in Cohort D will receive the following:

- After apheresis and prior to administration of cyclophosphamide and fludarabine (conditioning regimen prior to JNJ-68284528 infusion): 1 or more cycles of lenalidomide at a dose of 10 mg per day upon adequate hematologic recovery from ASCT (absolute neutrophil count [ANC] ≥ 1 x 10<sup>9</sup>/L and platelet count ≥ 75 x 10<sup>9</sup>/L). Alternative bridging therapy regimen is permissible with sponsor approval.
- After JNJ-68284528 production and product release, subjects will receive a conditioning regimen consisting of cyclophosphamide 300 mg/m<sup>2</sup> IV daily and fludarabine 30 mg/m<sup>2</sup> IV daily for 3 days. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an estimated glomerular filtration rate (eGFR) of 30 to 70 mL/min/1.73m<sup>2</sup>. JNJ-68284528 IV infusion will take place 5 to 7 days after the start of the conditioning regimen. The target dose is 0.75 x 10<sup>6</sup> CAR-positive viable T cells/kg (range: 0.5-1.0 x 10<sup>6</sup> CAR-positive viable T cells/kg).
- A strategy of staggered dosing with JNJ-68284528 will be applied at the start of enrollment to Cohort D. There must be an observation period of at least 4 weeks between administration of JNJ-68284528 to the first 5 subjects to allow for subject 28-day safety review prior to next subject dosing. In addition, the first 5 subjects will not receive lenalidomide after JNJ-68284528 therapy. After the first 5 subjects are dosed with JNJ-68284528, the Data Monitoring Committee (DMC) will convene to review safety and any other relevant data.

- Based on recommendation of the DMC, subsequent subjects in this cohort will be eligible to receive lenalidomide after JNJ-68284528. The same strategy of staggered dosing will also be applied to the first 5 subjects receiving lenalidomide after JNJ-68284528 (ie, the 6<sup>th</sup> through 10<sup>th</sup> subject enrolled in Cohort D). There must be an observation period of at least 4 weeks between administration of JNJ-68284528 (followed by lenalidomide) to these 5 subjects. The DMC will review safety and any other relevant data from the first 5 subjects who receive JNJ-68284528 followed by lenalidomide before a decision is made regarding the treatment plan for further subjects enrolled in Cohort D.

JNJ-68284528 plus lenalidomide treatment: Subjects will initiate lenalidomide maintenance therapy at a minimum of 21 days post JNJ-68284528 infusion and after resolution of any cytokine release syndrome (CRS) or neurologic toxicities per ASTCT criteria. Subjects will continue to receive lenalidomide until confirmed PD, unacceptable toxicity, or for 2 years post JNJ-68284528 infusion, whichever occurs first. The initial dose of lenalidomide post-JNJ-68284528 infusion will depend on the level of hematologic recovery.

- Criteria for lenalidomide administration after JNJ-68284528 infusion depend on the level of hematologic recovery and are summarized in the table below.

Hematologic Parameter		Starting Dose of Lenalidomide
ANC	Platelet Count	
$\geq 1.0 \times 10^9/L$	AND $\geq 75 \times 10^9/L$	10 mg daily
$0.75 \times 10^9/L$ to $< 1.0 \times 10^9/L$	AND $\geq 50 \times 10^9/L$	Start 5 mg daily, increase to 10mg per day when ANC is $\geq 1.0 \times 10^9/L$ and the platelet count is $\geq 75 \times 10^9/L$
$\geq 0.75 \times 10^9/L$	AND $50 \times 10^9/L$ to $< 75 \times 10^9/L$	
$< 0.75 \times 10^9/L$	OR $< 50 \times 10^9/L$	Lenalidomide <u>must</u> be held if <u>either one</u> of these criteria are present

- If well tolerated after 3 cycles of lenalidomide treatment at 10 mg, the dose of lenalidomide may be increased to 15 mg per day at the discretion of the investigator.
- For subjects with an eGFR  $< 60$  mL/min/1.73m<sup>2</sup>, the lenalidomide dose should be reduced to 5 mg per day. Other dose adjustments should be performed based on local prescribing information consistent with the protocol guidance for dose reductions.

### Cohort E

Subjects in Cohort E will receive the following:

- Eligible subjects will receive 4 cycles of D-VRd induction therapy as tolerated. The apheresis will be performed after Cycle 1 or Cycle 2.
- After JNJ-68284528 production and product release, subjects will receive a conditioning regimen consisting of cyclophosphamide 300 mg/m<sup>2</sup> intravenously (IV) daily and fludarabine 30 mg/m<sup>2</sup> IV daily for 3 days. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an estimated glomerular filtration rate (eGFR) of 30 to 70 mL/min/1.73m<sup>2</sup>. The conditioning regimen will be initiated a minimum of 21 days after the last dose of D-VRd. JNJ-68284528 IV infusion will take place 5 to 7 days after the start of the conditioning regimen. The target dose is  $0.75 \times 10^6$  CAR-positive viable T cells/kg (range: 0.5-1.0 x 10<sup>6</sup> CAR-positive viable T cells/kg).
- After infusion of JNJ-68284528: All subjects will receive treatment with D+R consolidation therapy until confirmed PD, unacceptable toxicity, or for 2 years post JNJ-68284528 infusion, whichever occurs first. D+R should be started when the following conditions are met:

- Lenalidomide will be initiated at a minimum of 21 days post JNJ-68284528 infusion, and will only start after resolution of any grade cytokine release syndrome (CRS) or neurological toxicities per ASTCT criteria. The initial dose of lenalidomide will depend on the level of hematologic recovery.

Hematologic Parameter			Starting Dose of Lenalidomide
ANC		Platelet Count	
$\geq 1.0 \times 10^9/L$	AND	$\geq 75 \times 10^9/L$	10 mg daily
$0.75 \times 10^9/L$ to $< 1.0 \times 10^9/L$	AND	$\geq 50 \times 10^9/L$	Start 5mg daily, increase to 10mg per day when ANC is $\geq 1.0 \times 10^9/L$ <u>and</u> the platelet count is $\geq 75 \times 10^9/L$
$\geq 0.75 \times 10^9/L$	AND	$50 \times 10^9/L$ to $< 75 \times 10^9/L$	
$< 0.75 \times 10^9/L$	OR	$< 50 \times 10^9/L$	Lenalidomide <u>must</u> be held if <u>either one</u> of these criteria are present

- o For subjects with an eGFR  $< 60$  mL/min/1.73m<sup>2</sup>, dose reduction of lenalidomide should be performed based on local prescribing information and in alignment with the guidance provided in the protocol
- o If well tolerated after 3 cycles of 10 mg lenalidomide treatment, the dose of lenalidomide may be increased to 15 mg per day at the discretion of the investigator.
- o Further up-titration by 5 mg increments may proceed after 6 cycles until a maximum dose of 25 mg daily (eg, lenalidomide Cycle 1: 10mg, Cycle 2: 10mg, Cycle 3: 10mg, Cycle 4: 15mg, Cycle 5: 15mg, Cycle 6: 15mg, Cycle 7: 20mg, Cycle 8: 25mg).
- Daratumumab will be initiated approximately 60 to 100 days post JNJ-68284528 infusion, upon adequate recovery from ongoing toxicities. Initiate daratumumab once platelet and absolute neutrophil count (ANC) are  $\geq 50 \times 10^9/L$  and  $\geq 1.0 \times 10^9/L$ , respectively. The initiation of daratumumab should coincide with the beginning of a lenalidomide cycle. Subjects must complete 1 cycle of lenalidomide before starting daratumumab, even if hematologic recovery occurs sooner.

## EVALUATIONS

Disease status will be evaluated according to the IMWG consensus recommendations for multiple myeloma. Efficacy evaluations will include measurements of tumor burden/residual disease, myeloma proteins, bone marrow examinations, skeletal surveys, extramedullary plasmacytomas, and serum calcium corrected for albumin. For subjects in Cohorts A, B, and C with neither serum nor urine measurable disease, baseline PET/CT, or whole body MRI may be used to satisfy the measurable disease criteria.

Blood and serum samples will be collected for assessment of JNJ-68284528 pharmacokinetics, immunogenicity (antibodies to JNJ-68284528), and predictive biomarkers of response or resistance to JNJ-68284528.

Data regarding subjects' health-related quality of life (HRQoL), symptoms, functioning, and general well-being will be captured using patient-reported outcome (PRO) measures.

Safety will be evaluated by adverse events, laboratory test results, vital sign measurements, physical examination findings (including neurologic examination), assessments of cardiac function, Immune Effector Cell-associated Encephalopathy (ICE) score. Performance status will be assessed with the ECOG scale.

## **STATISTICAL METHODS**

No formal statistical hypothesis testing will be performed. The sample size of each cohort is selected to collect necessary data on preliminary efficacy and safety information.



**TIME AND EVENTS SCHEDULES: COHORT A, COHORT B, AND COHORT C (TABLE 1 AND TABLE 2)**

**Table 1: Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Post-treatment (Day 101 and up to End of Cohort) <sup>c</sup>
					Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	
<b>Screening Assessments</b>																
Informed consent <sup>a</sup>	X (Before the 1 <sup>st</sup> study related procedure)															
Eligibility criteria (See Section 4)	X															
Demography, Medical History (including neurologic history)	X															
Disease Characteristics <sup>e</sup>	X	Disease characteristics to be collected prior to start of conditioning regimen														
PET/CT or MRI whole body scan (subjects without measurable disease in serum or urine)	X	X <sup>f</sup> (≤72 hour window)			Perform at 6 months, 12 months, and then yearly (±16 days) after infusion of JNJ-68284528)											
ECOG performance status	X		Prior to 1 <sup>st</sup> dose only	X										X		X
12-lead ECG	X		As clinically indicated													
Physical Examination	X		A symptom-directed physical examination should be performed as clinically indicated													
Height	X															
Echocardiogram or MUGA scan	X (≤8 weeks of apheresis)		For subjects who receive bridging therapy that includes agents with known cardiac toxicity (per prescribing information), assessment of cardiac function should be repeated within 7 days prior to the start of the conditioning regimen then again as clinically indicated if the subject develops signs/symptoms of heart failure													

**Table 1: Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Post-treatment (Day 101 and up to End of Cohort) <sup>e</sup>	
					Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>		
ICE neurologic test	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Day -5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5	Day 1 (Infusion)	ICE test must be repeated at any incidence of suspected CAR-T cell-related neurotoxicity (eg, ICANS). Perform at least daily until ICANS is resolved.											(every 28 days up to 12 months then every 56 days) <sup>d</sup> (± 7 days)	
Handwriting sample				X (≤24 hours prior to infusion) <sup>f</sup>	X	X	X	X	X	X	X	X	X	X	X	X	Perform monthly up to Day 184
Ocular Exam; subjects who received prior ADC (Cohort C Only) <sup>f</sup>	X				As clinically indicated												
<b>Outpatient Administration:</b> In consultation with and approval of the sponsor. See <a href="#">Attachment 15</a> for outpatient monitoring																	
Evaluation for outpatient suitability (See <a href="#">Attachment 14</a> )		X		X (predose)													
All Subjects with hospital discharge on Day 10								Daily phone calls during business hours from site staff, Days 11-14									
<b>Assessments Prior to Apheresis and Conditioning Regimen</b>																	
Criteria for Apheresis (See Section 6.1.1)		X															
Criteria for Conditioning Regimen (See Section 6.1.2.1)			≤72 hours of the 1 <sup>st</sup> dose only														

**Table 1: Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Post-treatment (Day 101 and up to End of Cohort) <sup>e</sup>
					Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	
	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Day -5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5	Day 1 (Infusion)												(every 28 days up to 12 months then every 56 days) <sup>d</sup> (± 7 days)
Criteria for JNJ-68284528 Administration (See Section 6.1.3.2)				X (predose) ≤24-hour window												
<b>Laboratory Assessments (See Section 9.7).</b> To be performed by the local laboratory except for the calcium and albumin-adjusted calcium, which will be performed at the central laboratory (local labs may be used to assess eligibility). Blood samples collection may be performed at the participant’s home by mobile study personnel (i.e. nurses and mobile phlebotomist) in the post-treatment period, after the Day 100.																
Hematology	X	X (≤72 hours prior to apheresis)	Prior to 1 <sup>st</sup> dose only (≤72 hour window)	X (predose) ≤24-hour window	X	X	X	X	X	X		X	X	X	X	
Chemistry	X	X (≤72 hour window)	Prior to 1 <sup>st</sup> dose only (≤72 hour window)	X (predose) ≤24-hour window	X	X	X	X	X	X		X	X	X	X	
Serology <sup>g</sup>	X				For subjects at risk for HBV activation monitor HBV DNA and AST/ALT every 12 weeks (± 7 days) for 12 months post JNJ-68284528 dose <sup>g</sup>											
Coagulation (PT/INR, aPTT, fibrinogen, D-dimer)	X				As clinically indicated for subjects who have fever or other signs of potential CRS											
Serum Pregnancy test (in subjects with childbearing potential)	X	X (≤72 hour window)	Prior to 1 <sup>st</sup> dose only (≤72 hour window)		As clinically indicated or as mandated by local regulations, whichever is more stringent											
Infectious Disease Testing <sup>h</sup>	X (within 60 days prior to apheresis, as applicable per local regulations)															

**Table 1: Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Post-treatment (Day 101 and up to End of Cohort) <sup>e</sup>
					Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	
	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Day -5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5	Day 1 (Infusion)												(every 28 days up to 12 months then every 56 days) <sup>d</sup> (± 7 days)
<b>Study Intervention Administration</b>																
Weight	X	X (for JNJ-68284528 dose calculation)	Prior to 1 <sup>st</sup> dose only	X												
Vital signs, including oxygen saturation	X	X	X	X <sup>i</sup>	X	X	X	X	X	X			X			
Temperature					Measure at least twice a day <sup>j</sup>											
Apheresis		X														
Cyclophosphamide and fludarabine			X													
Pre-infusion medication (see Section 6.1.3.3 for requirements prior to dosing with JNJ-68284528)				X												
JNJ-68284528 (See SIPP and IPPI for administration of JNJ-68284528)				X												
<b>Serum and Urine Disease Evaluations</b> (See Section 9.2 for efficacy assessments. Blood and 24-hour urine: to be sent to the central laboratory <sup>8</sup> . Disease evaluation should continue to be performed until confirmed disease progression, death, start of a new anticancer treatment, withdrawal of consent for study participation, or study completion, whichever occurs first.) Subjects with disease progression who receive retreatment with JNJ-68284528 must continue with disease evaluation visits. For subjects at select sites, blood samples collection may be performed at the participant’s home by mobile study personnel (i.e. nurses and mobile phlebotomist) in the post-treatment period, after the Day 100.																
Serum β2-microglobulin			X (prior to first dose [≤7 days])													
Quantitative Immunoglobulins <sup>k, q</sup>	X		X (prior to first dose [≤7 days])							X			X	X	X	X

**Table 1: Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Post-treatment (Day 101 and up to End of Cohort) <sup>e</sup>
					Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	
	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Day -5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5													(every 28 days up to 12 months then every 56 days) <sup>d</sup> (± 7 days)
Serum M-protein quantitation by electrophoresis	X		X (prior to first dose [≤7 days])							X			X	X	X	X
24-hour urine protein electrophoresis sample	X <sup>1</sup>		X (prior to first dose [≤7 days])							X			X	X	X	X
Serum calcium corrected for albumin	X		X (prior to first dose [≤7 days])							X			X	X	X	X
Serum FLC and serum/urine immunofixation	X		Serum FLC and serum/urine immunofixation are to be performed prior to the start of conditioning regimen (Day -5 [≤7 days]) and when CR is suspected or maintained; for subjects with measurable disease only by light chain criteria serum FLC will also be performed at every assessment when an SPEP is performed													
<b>Other Disease Evaluations</b>																
MRD (bone marrow aspirate) <sup>m</sup>			X (prior to first dose [≤7 days])	Sample should be collected: <ul style="list-style-type: none"> <li>For all dosed subjects at Day 56, and at 6 month, 12 month, 18 month (Day 520), 24 (Day 744) months (± 16 days) regardless of the status of disease measured in blood and urine</li> <li>For subjects with suspected CR, at the time of CR and then yearly for subjects that remain on study up to disease progression.</li> </ul>												
Bone marrow aspirate and core biopsy for disease evaluation			X (prior to first dose [≤7 days])	To confirm CR, sCR, and at disease progression (immunohistochemistry or immunofluorescence). Can be taken at the same time as the MRD sample if needed.												
Skeletal Survey <sup>n</sup>	X (at screening or ≤14 days prior to first dose of conditioning regimen)			As clinically indicated to document disease progression or response.												
Assess extramedullary Plasmacytomas <sup>o</sup>			X (≤14 days prior to first dose)	By physical examination (if applicable): measurable sites at D28, D56, D78, D100 and then every 4 weeks. By radiologic imaging: D78, D156 then every 12 weeks (for all subjects with a history of plasmacytomas or as clinically indicated for others) Day 28 to 100 (±2 day window), Day 101 and later (±7 day window).												
MRD assessment by imaging (optional)				If performed, must be recorded in the CRF												
Biomarker evaluations	See Biomarker Time & Events Schedule ( <a href="#">Table 2</a> )															

**Table 1: Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Post-treatment (Day 101 and up to End of Cohort) <sup>e</sup>
					Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	
	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Day -5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5	Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	(every 28 days up to 12 months then every 56 days) <sup>d</sup> (± 7 days)
<b>Patient Reported Outcomes (PRO) and Medical Resource Utilization (MRU):</b> PRO assessments to be completed before any clinical tests or procedures scheduled for the same day as the PRO assessments that would influence the subject’s perceptions of their current health																
EORTC QLQ-C30	X	X (≤72 hour window)				X				X			X	X	X	X; every 112 days (±7 days)
MySLm-Q (Unless translation not available)	X	X (≤72 hour window)				X				X			X	X	X	X; every 112 days (±7 days)
PGIS	X	X (≤72 hour window)				X				X			X	X	X	X; every 112 days (±7 days)
PGIC										X			X	X	X	
PRO CTCAE	X					X				X			X	X	X	
MRU				X				X		X			X		X	X; every 28 days until Day 180
<b>Ongoing Subject Review</b>																
Adverse Events	Continuous from the time of signing of ICF until 100 days after last administration of any study treatment. Beyond the adverse event reporting period, adverse events that are considered related to study drug need to be reported until the end of the study. Report new neurologic adverse events or exacerbation of existing neurologic adverse events up to 12 months after JNJ-68284528 infusion. Second primary malignancies should be followed from the time of signing of ICF signing to study completion <sup>p</sup> CRS should be evaluated according to the ASBMT (ASTCT) consensus grading (Lee 2019) (Attachment 2) CAR-T cell-related neurotoxicity (eg, ICANS) should be graded according to the ASBMT (ASTCT) consensus grading (Attachment 4) Events of HBV reactivations should be reported during the first year post-dosing of JNJ-68284528															
Concomitant medication	Continuous from the time of signing of ICF until at least 100 days after last administration of any study treatment. Thereafter, continue to report concomitant therapy given for any Adverse Events considered related to study drug until the end of the study.															
Survival Follow-up	After disease progression is documented, survival status will be obtained every 16 weeks until study completion. Survival status will also be collected prior to any planned efficacy analysis.															
Subsequent Anticancer Therapy	After disease progression is confirmed, subsequent anticancer therapy will be obtained every 16 weeks until study completion															

Abbreviations: ADC=antibody-drug conjugate; aPTT=activated partial thromboplastin time; ASBMT=American Society for Blood and Bone Marrow Transplantation; ASTCT=American Society for Transplantation and Cellular Therapy; CAR-T=chimeric antigen receptor T (cells); CR=complete response; sCR=stringent complete response; CRS= cytokine release syndrome; CT=computed tomography; CTCAE=Common Terminology Criteria for Adverse Events; D=Day; ECOG=Eastern Cooperative Oncology Group; EORTC-QLQ=European Organization for Research and Treatment of Cancer Quality of Life Questionnaire; FISH=fluorescence in situ hybridization; FLC=free light chain; HBV=hepatitis B virus; ICANS=Immune-effector Cell-associated Neurotoxicity Syndrome; ICE=Immune effector Cell-associated Encephalopathy; ICF=informed consent form; INR=international normalized ratio; IPP=investigational product preparation instructions; MRD=minimal residual disease; MRI=magnetic resonance imaging; MRU=Medical Resource Utilization; MUGA=multiple-gated acquisition; MySim-Q=Multiple Myeloma Symptom and Impact Questionnaire; PGIC=Patient Global Impression of Change; PGIS=Patient Global Impression of Severity; PRO=patient reported outcome; PT=prothrombin time; SIPP=site investigational product procedures manual; SPEP=serum protein electrophoresis; UPEP=urine protein electrophoresis.

- <sup>a</sup> ICF must be signed before any study-related procedures are performed and remains in effect even if the screening evaluation is not performed within the 28-day Screening Phase. Evaluations for eligibility determination performed outside the screening window may need to be repeated. For subjects who require a repeat apheresis see Section 9.1.3 for assessments that should be collected before the second apheresis. If the second apheresis falls outside of the 28-day window, all screening assessments (except bone marrow collection) must be repeated.
- <sup>b</sup> Assessments may be conducted  $\leq 72$  hours predose unless otherwise noted.
- <sup>c</sup> For subjects who discontinue the study before Day 100, the Day 100 assessments should be performed prior to withdrawal if feasible. Subjects who discontinue after Day 100 but before study completion should have urine and serum disease assessments performed prior to withdrawal if feasible at the time of discontinuation, unless performed within 14 days prior to discontinuation. The end of each study cohort will be defined as 2 years after the last subject in that particular cohort received his or her initial dose of JNJ-68284528
- <sup>d</sup> Post-treatment assessments to be obtained until progressive disease is documented or the start of subsequent anticancer therapy, with the exception of survival status and subsequent anticancer therapy, which are to be collected until study completion. PRO assessments are collected until study completion (continued after disease progression or subsequent anticancer therapy).
- <sup>e</sup> Disease characteristics cytogenetics (full karyotyping or FISH as well as molecular genetics [if applicable], both of which may originate from a bone marrow assessment performed prior to or during the screening period until the start of the conditioning regimen) and information on extra-medullary disease, as applicable. A pathologist/cytogeneticist should complete the cytogenetics data collection worksheet.
- <sup>f</sup> Pre-infusion ICE test and handwriting sample should be performed before pre-medication with diphenhydramine
- <sup>g</sup> Serology results performed as standard of care within 28 days prior to apheresis are acceptable. Hepatitis B: HBsAg, anti-HBc, anti-HBs, HBV DNA quantification [for subjects who are anti-HBs positive without a history of vaccination or for subjects who are anti-HBc positive with or without anti-HBs positive; if HBV-DNA is negative: monitor HBV-DNA, AST/ALT every 12 weeks ( $\pm 7$  days) for 12 months post-dose of JNJ-68284528]. Hepatitis C: HCV antibody, HCV RNA (for subjects who are anti HCV positive); HIV serology. See [Attachment 10](#)
- <sup>h</sup> HIV, hepatitis B, hepatitis C, HTLV, and other infectious diseases will be performed within 60 days of apheresis or as applicable per local regulations, whichever is more stringent
- <sup>i</sup> Immediately before the start of infusion, at the end of infusion, and 0.5, 1, 2 hours after end of infusion (window  $\pm 10$  minutes). Monitor until normalized after a CRS event.
- <sup>j</sup> Temperature will be checked at least twice a day up to Day 28. Subjects will be provided with a thermometer and instructed on the use of the thermometer and entering 2 temperatures including their maximum daily temperature in a diary. Diary will be reviewed at each visit, then collected on Day 28 and stored with patient source documents.
- <sup>k</sup> All subjects will be evaluated for IgG, IgA, IgM. Testing for IgD and IgE will only be performed for subjects with IgD and IgE-type myeloma.
- <sup>l</sup> UPEP sample collected as part of the standard of care and prior to the subject signing ICF may be used for analysis at the central laboratory.
- <sup>m</sup> Bone marrow morphology from an aspirate and core biopsy to be assessed locally at all time points. Additional bone marrow aspirate samples will be collected for biomarkers (see [Table 2](#)). Bone marrow aspirate for MRD should be taken from first or second aspiration attempt, if feasible. If for any reason a bone marrow aspirate is not performed at pre-dose, or if a baseline clone cannot be established from the pre-dose bone marrow aspirate collection, then non-decalcified diagnostic tissue will be requested.
- <sup>n</sup> Results from skeletal survey performed as routine follow-up within 42 days before start of apheresis may be used without these tests being repeated. Additional imaging (X-ray, CT, or MRI) will be performed as clinically indicated (eg, to document response or progression) (Section 9.2.5).
- <sup>o</sup> Results from radiologic plasmacytoma assessments performed  $\leq 14$  days prior to the first dose of the conditioning as routine follow-up for subject's disease may be used. Extramedullary plasmacytomas should be assessed for all subjects with a history of plasmacytomas or if clinically indicated at screening, by clinical examination or radiologic imaging (Section 9.2.6).
- <sup>p</sup> A tumor sample should be collected and sent to the sponsor for DNA, RNA, or protein analysis to investigate the presence of lentiviral elements
- <sup>q</sup> Additional immunoglobulin samples may be collected as clinically indicated for safety

- <sup>r</sup> For Cohort C, subjects who have received prior antibody-drug conjugate (ADC), ocular exam is to include best-corrected visual acuity (BCVA), slit lamp examination (with special focus on cornea), intraocular pressure, and dilated fundoscopic examination.
- <sup>s</sup> Local laboratory assessments may be used under specified circumstances (See Section [9.2.7](#))
- <sup>t</sup> Only if screening assessment was performed using whole body MRI. Not required if PET/CT imaging is utilized



**Table 2: Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling**

	Screening Phase	Apheresis	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (any subject who received infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup> and Post-treatment (Day 101 up to study completion) <sup>b</sup>													At PD	At Study Completion for subjects without PD
					Day 1 (Infusion)	Day 2 (± 2 hour)	Day 3 (± 4 hour)	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days)		
<b>Pharmacokinetics</b>																			
PK CAR transgene levels blood sample <sup>b</sup>			X (prior to first dose [≤7 days])	Pre-dose (≤4 hr window); Post EOI (within 30 minutes)	24-hour post-EOI	X	X	X	X	X	X		X	X	X	X; then every 8 weeks up to 1 year		X	X
Soluble serum BCMA sample			X (prior to first dose [≤7 days])	Pre-dose (≤4 hr window); Post EOI (within 30 minutes)	24-hour post-EOI	X	X	X	X	X	X		X	X	X	X; then every 8 weeks up to 1 year		X	X
PK CAR transgene levels bone marrow sample			X (prior to first dose [≤7 days])								X			X			X		
ADA sample (serum) <sup>b,c</sup>				Pre-dose ≤4 hour window					X		X			X	X	X	X	X	X
<b>Biomarker Sampling</b>																			
Immuno-phenotyping (whole blood) <sup>d, f</sup>		X (≤72 hour window)	X (prior to first dose [≤7 days])	Pre-dose ≤4 hour window	24-hour post EOI		X	X	X	X	X	X	X	X	X	X; then every 8 weeks up to 1 year <sup>h</sup>		X	X <sup>d</sup>

**Table 2: Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling**

	Screening Phase	Apheresis	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (any subject who received infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup> and Post-treatment (Day 101 up to study completion) <sup>b</sup>													At PD	At Study Completion for subjects without PD			
					Day 1 (Infusion)	Day 2 (± 2 hour)	Day 3 (± 4 hour)	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days)			Day 184 (± 7 days)		
Flow cytometry, (aspirate) (bone marrow) <sup>d,i</sup>	≤28 days prior to apheresis	Upon enrollment	Day -5,* -4, -3 *Window of Day -7 to Day -5	Day 1 (Infusion)								X			X			X	X	X		
CyTOF (aspirate) (bone marrow) <sup>d,e</sup>			X (prior to first dose [≤7 days])									X			X			X	X	X		
CyTOF/PBMC/Plasma (whole blood) <sup>d,e</sup>		X (≤72 hour window)					X	X	X	X	X	X	X	X			X	X	X	X		
PBMCs for functional in vitro assays		X (≤72 hour window)						X				X			X		X;	then every 8 weeks up to 1 year				
Cytogenetics (bone marrow)			X (prior to first dose [≤7 days])																X			
Replication Competent Lentivirus (RCL) (whole blood)			X Prior to 1 <sup>st</sup> dose only (≤7 day window)	Pre-dose (≤4 hour window)		At approximately 3, 6, and 12 months; then yearly for 15 years post infusion																
Serum protein analysis			X (prior to first dose [≤7 days])	Pre-dose (≤4 hour window)	Additional serum protein sample will be taken at each SIFE/SPEP evaluation, as well as at each MRD sample collection																	

**Table 2: Cohort A, Cohort B, and Cohort C: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling**

	Screening Phase	Apheresis	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (any subject who received infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup> and Post-treatment (Day 101 up to study completion) <sup>g</sup>													At PD	At Study Completion for subjects without PD
					Day 2 (± 2 hour)	Day 3 (± 4 hour)	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days)	Day 184 (± 7 days)		
Cytokine profiling <sup>f</sup> (serum)	≤28 days prior to apheresis	Upon enrollment	Day -5,* -4, -3 *Window of Day -7 to Day -5	Day 1 (Infusion)  Pre-dose; 2hrs Post-dose (±10 minutes)	X	X	X	X	X	X	X		X	X	X	X			

Abbreviations: ADA=anti-drug antibody; BCMA=B-cell maturation antigen; CAR=chimeric antigen receptor; CR = complete response; CRS=cytokine release syndrome; CyTOF=cytometry by time of flight; EOI=end of infusion; MRD=minimal residual disease; PBMC=peripheral blood mononuclear cell; PD=progressive disease; PK=pharmacokinetic; sCR=stringent complete response

- <sup>a</sup> For subjects who discontinue the study before Day 100, the Day 100 assessments should be performed if feasible.
- <sup>b</sup> Collect additional samples when any of the following are suspected or reported: 1) CRS or CAR-T cell-related neurotoxicity (eg, ICANS) Grade ≥2 (at onset of the event, and 24 and 72 hours after) or as clinically indicated; and 2) as indicated based on emerging data
- <sup>c</sup> ADA sample should be collected if a subject withdraws from the study after JNJ-68284528 administration but prior to disease progression or study completion.
- <sup>d</sup> Sample should be collected at suspected CR.
- <sup>e</sup> Sample should be collected at 12 months, relative to Day 1, for subjects that achieve CR/sCR and remain on study
- <sup>f</sup> Collect additional samples when any of the following are suspected or reported: 1) CRS or CAR-T cell-related neurotoxicity (eg, ICANS) (any grade) (at onset of the event, and then every 24 hours until CRS or ICANS event has stabilized or is resolving at which time additional collections should occur at 24, 48, and 72 hours) or as clinically indicated; and 2) as indicated based on emerging data.
- <sup>g</sup> Pharmacokinetic and biomarker samples collection (serum samples only) may be performed at the subject’s home by mobile study personnel (i.e. nurses and mobile phlebotomist) in the post-treatment period, after the Day 100.
- <sup>h</sup> If the 1-year immunophenotyping sample has detectable levels of CAR+ T cells, additional samples may be collected for central laboratory assessment at the discretion of the investigator
- <sup>i</sup> If the last post-D100 immunophenotyping samples has detectable levels of CAR+ T cells an additional bone marrow sample may be collected at the discretion of the investigator for central laboratory assessment

**TIME AND EVENTS SCHEDULES: COHORT D (TABLE 3 AND TABLE 4)**

**Table 3: Cohort D: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Lenalidomide (upon adequate hematologic recovery from ASCT) <sup>y</sup>	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Lenalidomide Treatment Period (As early as Day 22 and up to End of Cohort) <sup>e</sup>
						Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	
<b>Screening Assessments</b>																	
Informed consent <sup>a</sup>	X (Before the 1 <sup>st</sup> study related procedure)																
Eligibility criteria (See Section 4)	X																
Demography, Medical History (including neurologic history)	X																
Disease Characteristics <sup>e</sup>	X																
ECOG performance status	X			Prior to 1 <sup>st</sup> dose only	X									X		X	
12-lead ECG	X		As clinically indicated														
Physical Examination	X		A symptom-directed physical examination should be performed as clinically indicated														
Height	X																
Echocardiogram or MUGA scan	X (≤8 weeks of apheresis)		For subjects who receive bridging therapy that includes agents with known cardiac toxicity (per prescribing information), assessment of cardiac function should be repeated within 7 days prior to the start of the conditioning regimen then again as clinically indicated if the subject develops signs/symptoms of heart failure														
ICE neurologic test					X (≤24 hours prior to infusion) <sup>f</sup>	ICE test must be repeated at any incidence of suspected CAR-T cell-related neurotoxicity (eg, ICANS). Perform at least daily until ICANS is resolved.											

**Table 3: Cohort D: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Lenalidomide (upon adequate hematologic recovery from ASCT) <sup>y</sup>	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Lenalidomide Treatment Period (As early as Day 22 and up to End of Cohort) <sup>c</sup>
						Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	
	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Cycle 1 (Additional Cycles permitted with Sponsor approval)	Day - 5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5	Day 1 (Infusion)												(every 28 days up to 12 months then every 56 days) <sup>d,t</sup> (± 7 days)
Handwriting sample					X (≤24 hours prior to infusion) <sup>f</sup>	X	X	X	X	X	X	X	X	X	X	X	Perform monthly up to Day 184
<b>Outpatient Administration:</b> In consultation with and approval of the sponsor. See <a href="#">Attachment 15</a> for outpatient monitoring																	
Evaluation for outpatient suitability (See <a href="#">Attachment 14</a> )		X			X (predose)												
All Subjects with hospital discharge on Day 10									Daily phone calls during business hours from site staff, Days 11-14								
<b>Assessments Prior to Apheresis and Conditioning Regimen</b>																	
Criteria for Apheresis (See <a href="#">Section 6.1.1</a> )		X															
Criteria for Conditioning Regimen (See <a href="#">Section 6.1.2.1</a> )				≤72 hours of the 1 <sup>st</sup> dose only													
Criteria for JNJ-68284528 Administration (See <a href="#">Section 6.1.3.2</a> )					X (predose)												

**Table 3: Cohort D: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Lenalidomide (upon adequate hematologic recovery from ASCT) <sup>v</sup>	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Lenalidomide Treatment Period (As early as Day 22 and up to End of Cohort) <sup>c</sup>
						Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	
	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Cycle 1 (Additional Cycles permitted with Sponsor approval)	Day - 5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5	Day 1 (Infusion)												(every 28 days up to 12 months then every 56 days) <sup>d,t</sup> (± 7 days)
<b>Laboratory Assessments (See Section 9.7).</b> To be performed by the local laboratory except for the calcium and albumin-adjusted calcium, which will be performed at the central laboratory (local labs may be used to assess eligibility). Blood samples collection may be performed at the participant’s home by mobile study personnel (i.e. nurses and mobile phlebotomist) in the post-treatment period, after the Day 100.																	
Hematology <sup>s</sup>	X	X (≤72 hours prior to apheresis)	X <sup>w</sup> (within 24 hrs prior to 1st dose)	Prior to 1 <sup>st</sup> dose only (≤72 hour window)	X (predose) (≤24 hour window)	X	X	X	X	X	X		X	X	X	X	X <sup>w</sup>
Chemistry <sup>s</sup>	X	X (≤72 hour window)	X (within 24 hrs prior to 1st dose and at the start of each Len cycle)	Prior to 1 <sup>st</sup> dose only (≤72 hour window)	X (predose) (≤24 hour window)	X	X	X	X	X	X		X	X	X	X	X (at start of each Len cycle)
Serology <sup>g</sup>	X					For subjects at risk for HBV activation monitor HBV DNA and AST/ALT every 12 weeks (±7 days) for 12 months post JNJ-68284528 dose <sup>g</sup>											
Coagulation (PT/INR, aPTT, fibrinogen, D-dimer)	X					As clinically indicated for subjects who have fever or other signs of potential CRS											
Thyroid assessment (via local laboratory) <sup>f</sup>	X					Every 6 months or more frequently, as clinically necessary											
Serum Pregnancy test (in subjects with childbearing potential)	X	X (≤72 hour window)	10-14 d prior to 1st dose and within 24 hrs prior to 1st dose	Prior to 1 <sup>st</sup> dose only (≤72 hour window)	Within 24 hours prior to start of lenalidomide, every week for the first 4 weeks, then repeated at least every 28 days Pregnancy testing must be performed 4 weeks after stopping lenalidomide. Additional pregnancy testing is done as clinically indicated or consistent with any country specific requirements listed in the local prescribing information for lenalidomide, whichever is more stringent												
Infectious Disease Testing <sup>h</sup>	X (within 60 days prior to apheresis, as applicable per local regulations)																

**Table 3: Cohort D: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Lenalidomide (upon adequate hematologic recovery from ASCT) <sup>y</sup>	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Lenalidomide Treatment Period (As early as Day 22 and up to End of Cohort) <sup>c</sup>
						Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	
	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Cycle 1 (Additional Cycles permitted with Sponsor approval)	Day -5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5	Day 1 (Infusion)												(every 28 days up to 12 months then every 56 days) <sup>d,t</sup> (± 7 days)
<b>Study Intervention Administration</b>																	
Weight	X	X (for JNJ-68284528 dose calculation)		Prior to 1 <sup>st</sup> dose only	X												
Vital signs, including oxygen saturation	X	X		X	X <sup>i</sup>	X	X	X	X	X	X			X			
Temperature <sup>i</sup>						Measure at least twice a day <sup>j</sup>											
Apheresis		X															
Cyclophosphamide and fludarabine				X													
Pre-infusion medication (see Section 6.1.3.3 for requirements prior to dosing with JNJ-68284528)					X												
JNJ-68284528 (See SIPP and IPPI for administration of JNJ-68284528)					X												
Lenalidomide			X			Initiate lenalidomide a minimum of 21 days post JNJ-68284528 after resolution of associated CRS or neurologic toxicities. Refer to <a href="#">Table 12</a> and Section 6.1.4 for additional dosing considerations Lenalidomide will be taken daily (continuously) in 28-day cycles, until PD											
<b>Accountability/Exposure Check</b>																	
Pill count for Lenalidomide <sup>s</sup>			X								X			X	X	X	X

**Table 3: Cohort D: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Lenalidomide (upon adequate hematologic recovery from ASCT) <sup>y</sup>	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Lenalidomide Treatment Period (As early as Day 22 and up to End of Cohort) <sup>c</sup>
						Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	
	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Cycle 1 (Additional Cycles permitted with Sponsor approval)	Day - 5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5	Day 1 (Infusion)												(every 28 days up to 12 months then every 56 days) <sup>d,†</sup> (± 7 days)
<b>Serum and Urine Disease Evaluations</b> (See Section 9.2 for efficacy assessments. Blood and 24-hour urine: to be sent to the central laboratory <sup>u</sup> . Disease evaluation should continue to be performed until confirmed disease progression, death, start of a new anticancer treatment, withdrawal of consent for study participation, or study completion, whichever occurs first.). Disease evaluation for subjects that discontinue lenalidomide treatment due to adverse events should continue until disease progression. For subjects at select sites within the US, blood samples collection may be performed at the participant’s home by mobile study personnel (i.e. nurses and mobile phlebotomist) in the post-treatment period, after the Day 100																	
Serum β2-microglobulin				X (prior to first dose [≤7 days])													
Quantitative Immunoglobulins <sup>k,q</sup>	X			X (prior to first dose [≤7 days])						X			X	X	X		X
Serum M-protein quantitation by electrophoresis	X			X (prior to first dose [≤7 days])						X			X	X	X		X
24-hour urine protein electrophoresis sample	X <sup>1</sup>			X (prior to first dose [≤7 days])						X			X	X	X		X
Serum calcium corrected for albumin	X			X (prior to first dose [≤7 days])						X			X	X	X		X
Serum FLC and serum/urine immunofixation	X			Serum FLC and serum/urine immunofixation are to be performed prior to the start of conditioning regimen (Day -5 [≤7 days]) and when CR is suspected or maintained; for subjects with measurable disease only by light chain criteria serum FLC will also be performed at every assessment when an SPEP is performed													
<b>Other Disease Evaluations</b>																	
MRD (bone marrow aspirate) <sup>m</sup>	X				Sample should be collected: <ul style="list-style-type: none"> <li>For all dosed subjects at Day 56, and at 6 month, 12 month, 18 month (Day 520), 24 (Day 744) months (± 16 days) regardless of the status of disease measured in blood and urine</li> <li>For subjects with suspected CR at the time of CR and then yearly for subjects that remain on study up to disease progression.</li> </ul>												



**Table 3: Cohort D: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Lenalidomide (upon adequate hematologic recovery from ASCT) <sup>y</sup>	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Lenalidomide Treatment Period (As early as Day 22 and up to End of Cohort) <sup>c</sup>
						Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	
Bone marrow aspirate and core biopsy for disease evaluation	X			X (prior to first dose [≤7 days])	To confirm CR, sCR, and at disease progression (immunohistochemistry or immunofluorescence). Can be taken at the same time as MRD sample if needed.												
Skeletal Survey <sup>n</sup>	X (at screening or ≤14 days prior to the first dose of conditioning regimen)				As clinically indicated to document disease progression or response.												
Assess extramedullary Plasmacytomas <sup>o</sup>	X			X (≤14 days prior to first dose)	Measurable sites Day 28, Day 56, Day 78, Day 100 then every 4 weeks for physical examination (if applicable) and Day 78 and Day 156 then every 12 weeks for radiologic assessment (for subjects with a history of plasmacytomas or as clinically indicated for others). Day 28 to 100 (±2 day window), Day 101 and later (±7 day window).												
MRD assessment by imaging (optional)	X				If performed, must be recorded in the CRF												
Biomarker evaluations	See Biomarker Time & Events Schedule (Table 2)																
<b>Patient Reported Outcomes (PRO) and Medical Resource Utilization (MRU):</b> PRO assessments to be completed before any clinical tests or procedures scheduled for the same day as the PRO assessments that would influence the subject's perceptions of their current health																	
EORTC QLQ-C30	X	X (≤72 hour window)				X					X			X	X	X	X; every 112 days (±7 days)
MySI-m-Q (Unless translation not available)	X	X (≤2 hour window)				X					X			X	X	X	X; every 112 days (±7 days)
PGIS	X	X (≤72 hour window)				X					X			X	X	X	X; every 112 days (±7 days)
PGIC											X			X	X	X	
PRO CTCAE	X					X					X			X	X	X	
MRU					X				X		X			X		X	X; Every 28 days until Day 180

**Table 3: Cohort D: Time and Events Schedule for Study Procedures/Assessments**

	Screening Phase	Apheresis	Lenalidomide (upon adequate hematologic recovery from ASCT) <sup>y</sup>	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>c</sup>											Lenalidomide Treatment Period (As early as Day 22 and up to End of Cohort) <sup>c</sup>
						Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	
	≤28 days prior to apheresis <sup>a</sup>	Upon enrollment	Cycle 1 (Additional Cycles permitted with Sponsor approval)	Day - 5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5	Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>c</sup>	(every 28 days up to 12 months then every 56 days) <sup>d,†</sup> (± 7 days)
<b>Ongoing Subject Review</b>																	
Adverse Events	Continuous from the time of signing of ICF until 100 days after administration of JNJ-68284528, or 30 days after last dose of lenalidomide, whichever is later. Beyond the adverse event reporting period, adverse events that are considered related to study drug need to be reported until the end of the study. Report new neurologic adverse events or exacerbation of existing neurologic adverse events up to 12 months after JNJ-68284528 infusion. Second primary malignancies should be followed from the time of signing of ICF signing to study completion <sup>p</sup> CRS should be evaluated according to the ASBMT (ASTCT) consensus grading ( <a href="#">Lee 2019</a> ) ( <a href="#">Attachment 2</a> ) CAR-T cell-related neurotoxicity (eg, ICANS) should be graded according to the ASBMT (ASTCT) consensus grading ( <a href="#">Attachment 4</a> ) Events of HBV reactivations should be reported during the first year post-dosing of JNJ-68284528																
Concomitant medication	Continuous from the time of signing of ICF until at least 100 days after administration of JNJ-68284528, or 30 days after last dose of lenalidomide, whichever is later. Thereafter, continue to report concomitant therapy given for any Adverse Events considered related to study drug until the end of the study.																
Survival Follow-up	After disease progression is documented, survival status will be obtained every 16 weeks until study completion. Survival status will also be collected prior to any planned efficacy analysis.																
Subsequent Anticancer Therapy	After disease progression is confirmed, subsequent anticancer therapy will be obtained every 16 weeks until study completion																

Abbreviations: ANC=absolute neutrophil count; aPTT=activated partial thromboplastin time; ASBMT=American Society for Blood and Bone Marrow Transplantation; ASTCT=American Society for Transplantation and Cellular Therapy; CAR-T=chimeric antigen receptor T (cells); CR=complete response; sCR=stringent complete response; CRS=cytokine release syndrome; CT=computed tomography; CTCAE=Common Terminology Criteria for Adverse Events; D=Day; ECOG=Eastern Cooperative Oncology Group; EORTC-QLQ=European Organization for Research and Treatment of Cancer Quality of Life Questionnaire; FISH=fluorescence in situ hybridization; FLC=free light chain; HBV=hepatitis B virus; ICANS=Immune-effector Cell-Associated Neurotoxicity Syndrome; ICE=Immune effector Cell-associated Encephalopathy; ICF=informed consent form; INR=international normalized ratio; IPPI=investigational product preparation instructions; Len=lenalidomide; MRD=minimal residual disease; MRI=magnetic resonance imaging; MUGA=multiple-gated acquisition; MySim-Q=Multiple Myeloma Symptom and Impact Questionnaire; PGIC=Patient Global Impression of Change; PGIS=Patient Global Impression of Severity; PRO=patient reported outcome; PT=prothrombin time; SIPP=site investigational product procedures manual; SPEP=serum protein electrophoresis; UPEP=urine protein electrophoresis.

<sup>a</sup> ICF must be signed before any study-related procedures are performed and remains in effect even if the screening evaluation is not performed within the 28-day Screening Phase. Evaluations for eligibility determination performed outside the screening window may need to be repeated. For subjects who require a repeat apheresis see Section 9.1.3 for assessments that should be collected before the second apheresis. If the second apheresis falls outside of the 28 day window, all screening assessments (except bone marrow collection) must be repeated.

<sup>b</sup> Assessments may be conducted ≤72 hours predose unless otherwise noted. Window for the start of the conditioning regimen is Day -7 to Day -5.

<sup>c</sup> For subjects who discontinue the study before Day 100, the Day 100 assessments should be performed prior to withdrawal if feasible. Subjects who discontinue after Day 100 but before study completion should have urine and serum disease assessments performed prior to withdrawal if feasible at the time of discontinuation, unless performed within 14 days prior to discontinuation. Cohort D will be considered complete 2 ½ years after the last subject receives their initial dose of JNJ-68284528.

- d Post-treatment assessments to be obtained until progressive disease is documented or the start of subsequent anticancer therapy, with the exception of survival status and subsequent anticancer therapy, which are to be collected until study completion. PRO assessments are collected until study completion (continued after disease progression or subsequent anticancer therapy).
- e Disease characteristics cytogenetics (full karyotyping or FISH as well as molecular genetics [if applicable]), both of which may originate from prior to or during the screening period and information on extra-medullary disease, as applicable. A pathologist/cytogeneticist should complete the cytogenetics data collection worksheet.
- f Pre-infusion ICE test and handwriting sample should be performed before pre-medication with diphenhydramine
- g Serology results performed as standard of care within 28 days prior to apheresis are acceptable. Hepatitis B: HBsAg, anti-HBc, anti-HBs, HBV DNA quantification [for subjects who are anti-HBs positive without a history of vaccination or for subjects who are anti-HBc positive with or without anti-HBs positive ]; if HBV-DNA is negative: monitor HBV-DNA, AST/ALT every 12 weeks ( $\pm 7$  days) for one year post-dose of JNJ-68284528]. Hepatitis C: HCV antibody, HCV RNA (for subjects who are anti HCV positive); HIV serology. See [Attachment 10](#)
- h HIV, hepatitis B, hepatitis C, HTLV, and other infectious diseases will be performed within 60 days of apheresis or as applicable per local regulations, whichever is more stringent
- i Immediately before the start of infusion, at the end of infusion, and 0.5, 1, 2 hours after end of infusion (window  $\pm 10$  minutes). Monitor until normalized after a CRS event.
- j Temperature will be checked at least twice a day up to Day 28. Subjects will be provided with a thermometer and instructed on the use of the thermometer and entering 2 temperatures including their maximum daily temperature in a diary. Diary will be reviewed at each visit, then collected on Day 28 and stored with patient source documents.
- k All subjects will be evaluated for IgG, IgA, IgM. Testing for IgD and IgE will only be performed for subjects with IgD and IgE-type myeloma.
- l UPEP sample collected as part of the standard of care and prior to the subject signing ICF may be used for analysis at the central laboratory.
- m Bone marrow morphology from an aspirate and core biopsy to be assessed locally at all time points. Additional bone marrow aspirate samples will be collected for biomarkers (see [Table 2](#)). Bone marrow aspirate for MRD should be taken from first or second aspiration attempt, if feasible. If for any reason a bone marrow aspirate is not performed at pre-dose, or if a baseline clone cannot be established from the pre-dose bone marrow aspirate collection, then non-decalcified diagnostic tissue will be requested.
- n Results from skeletal survey performed as routine follow-up within 42 days before start of apheresis may be used without these tests being repeated. Additional imaging (X-ray, CT, or MRI) will be performed as clinically indicated (eg, to document response or progression) (Section [9.2.5](#)).
- o Results from radiologic plasmacytoma assessments performed  $\leq 14$  days prior to the first dose of the conditioning as routine follow-up for subject's disease may be used. Extramedullary plasmacytomas should be assessed for all subjects with a history of plasmacytomas or if clinically indicated at screening, by clinical examination or radiologic imaging (Section [9.2.6](#)).
- p A tumor sample should be collected and DNA, RNA, or protein analysis may be performed to investigate the presence of lentiviral elements
- q Additional immunoglobulin samples may be collected as clinically indicated for safety
- r Thyroid assessment: TSH only, if TSH is  $< \text{LLN}$  or  $> \text{ULN}$  need free T3 and free T4 testing
- s Hematology, chemistry and pill count are to be done every 28 days past 12 months; all other assessments will be collected every 56 days or as otherwise noted.
- t Hematology and chemistry samples will continue to be collected every 28 days for subjects receiving lenalidomide beyond 12 months.
- u Local laboratory assessments may be used under specified circumstances (see Section [9.2.7](#))
- v If an alternative bridging therapy is used, assessments specified for lenalidomide toxicity do not apply
- w Weekly for the first 2 cycles, twice a month for Cycle 3 then once a month at start of each lenalidomide cycle

**Table 4: Cohort D: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling**

	Screening Phase	Apheresis	Lenalido- mide	Cyclophos- phamide and fludarabine conditioning regimen	JNJ- 68284528 Infusion	Post Infusion (any subject who received infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup> and Post-treatment (Day 101 up to study completion) <sup>#</sup>														At PD	At Study Com- ple- tion for sub- jec- ts with- out PD
						Day 2 (± 2 hour)	Day 3 (± 4 hour)	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days)	Day 184 (±7 days)			
<b>Pharmacokinetics</b>																					
PK CAR transgene levels blood sample <sup>b</sup>				X (prior to first dose [≤7 days])	Pre-dose (≤4 hr window); Post EOI (within 30 minutes)	24- hour post- EOI	X	X	X	X	X	X		X	X	X	X; then every 8 weeks up to 1 year		X	X	
Soluble serum BCMA sample				X (prior to first dose [≤7 days])	Pre-dose (≤4 hr window); Post EOI (within 30 minutes)	24- hour post- EOI	X	X	X	X	X	X		X	X	X	X; then every 8 weeks up to 1 year		X	X	
PK CAR transgene levels bone marrow sample				X (prior to first dose [≤7 days])								X			X			X			
ADA sample (serum) <sup>b,c</sup>					Pre-dose (≤4 hour window)					X		X			X	X	X	X	X	X	

**Table 4: Cohort D: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling**

	Screening Phase	Apheresis	Lenalido- mide	Cyclophos- phamide and fludarabine conditioning regimen	JNJ- 68284528 Infusion	Post Infusion (any subject who received infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup> and Post-treatment (Day 101 up to study completion) <sup>a</sup>													At PD	At Study Com- ple- tion for sub- jec- ts with- out PD			
						Day 2 (± 2 hour)	Day 3 (± 4 hour)	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days)	Day 184 (±7 days)					
	≤28 days prior to apheresis	Upon enroll- ment	Cycle 1 (Additional Cycles permitted with Sponsor approval)	Day -5,* -4, -3 *Window of Day -7 to Day -5	Day 1 (Infusion)																		
<b>Biomarker Sampling</b>																							
Immuno- phenotyping (whole blood) <sup>d,f</sup> Includes the flow PK		X (≤72 hour window)		X (prior to first dose [≤7 days])	Pre-dose (≤4 hour window)	24 hour post EOI		X	X	X	X	X	X	X	X	X	X	X	X	X; then every 8 weeks up to 1 year <sup>h</sup>		X	X <sup>e</sup>
Flow cytometry, (aspirate) (bone marrow) <sup>d,i</sup>				X (prior to first dose [≤7 days])								X			X					X	X	X	
CytoF (aspirate) (bone marrow) <sup>d,e</sup>				X (prior to first dose [≤7 days])								X			X					X	X	X	
CytoF/ PBMC/ Plasma (whole blood) <sup>d,e</sup>		X (≤72 hour window)						X	X	X	X	X	X	X	X			X	X	X			X <sup>d,e</sup>
PBMCs for functional in vitro assays		X (≤72 hour window)							X			X			X			X		X; then every 8 weeks up to 1 year			

**Table 4: Cohort D: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling**

	Screening Phase	Apheresis	Lenalidomide	Cyclophosphamide and fludarabine conditioning regimen	JNJ-68284528 Infusion	Post Infusion (any subject who received infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup> and Post-treatment (Day 101 up to study completion) <sup>a</sup>														At PD	At Study Completion for subjects without PD	
						Day 2 (± 2 hour)	Day 3 (± 4 hour)	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days)	Day 184 (± 7 days)				
Cytogenetics (bone marrow)	≤28 days prior to apheresis	Upon enrollment	Cycle 1 (Additional Cycles permitted with Sponsor approval)	Day -5,* -4, -3 *Window of Day -7 to Day -5	Day 1 (Infusion)															X		
Replication Competent Lentivirus (RCL) (whole blood)				X Prior to 1 <sup>st</sup> dose only (≤7 day window)	Pre-dose (≤4 hour window)	At approximately 3, 6, and 12 months; then yearly for 15 years post infusion																
Serum protein analysis				X (prior to first dose [≤7 days])	Pre-dose (≤4 hour window)	Additional serum protein sample will be taken at each SIFE/SPEP evaluation, as well as at each MRD sample collection																
Cytokine profiling <sup>f</sup> (serum)				X Prior to 1 <sup>st</sup> dose only (≤7 day window)	Pre-dose; 2hrs Post-dose (±10 minutes)	X	X	X	X	X	X	X		X	X	X	X					

Abbreviations: ADA=anti-drug antibody; BCMA=B-cell maturation antigen; CAR=chimeric antigen receptor; CR = complete response; CRS=cytokine release syndrome; CyTOF=cytometry by time of flight; EOI=end of infusion; MRD=minimal residual disease; PBMC=peripheral blood mononuclear cell; PD=progressive disease; PK=pharmacokinetic; sCR=stringent complete response

<sup>a</sup> For subjects who discontinue the study before Day 100, the Day 100 assessments should be performed if feasible.  
<sup>b</sup> Collect additional samples when any of the following are suspected or reported: 1) CRS or CAR-T cell-related neurotoxicity (eg, ICANS) Grade ≥2 (at onset of the event, and 24 and 72 hours after) or as clinically indicated; and 2) as indicated based on emerging data  
<sup>c</sup> ADA sample should be collected if a subject withdraws from the study after JNJ-68284528 administration but prior to disease progression or study completion.  
<sup>d</sup> Sample should be collected at suspected CR.  
<sup>e</sup> Sample should be collected at 12 months, relative to Day 1, for subjects that achieve CR/sCR and remain on study  
<sup>f</sup> Collect additional samples when any of the following are suspected or reported: 1) CRS or CAR-T cell-related neurotoxicity (eg, ICANS) (any grade) (at onset of the event, and then every 24 hours until CRS or ICANS event has stabilized or is resolving at which time additional collections should occur at 24, 48, and 72 hours) or as clinically indicated; and 2) as indicated based on emerging data.

- <sup>g</sup> Pharmacokinetic and biomarker samples collection (serum samples only) may be performed at the participant's home by mobile study personnel (i.e. nurses and mobile phlebotomist) in the post-treatment period, after the Day 100.
- <sup>h</sup> If the 1-year immunophenotyping sample has detectable CAR+ T cells, additional samples may be collected for central laboratory assessment at the discretion of the investigator
- <sup>i</sup> If the last post-D100 immunophenotyping samples has detectable levels of CAR+ T cells and additional bone marrow sample may be collected at the discretion of the investigator for central laboratory assessment

**TIME AND EVENTS SCHEDULES: COHORT E (TABLE 5, TABLE 6, TABLE 7, AND TABLE 8)**

**Table 5: Cohort E: Time and Events Schedule for Study Procedures/Assessments (Screening to Conditioning Regimen)**

	Screening Phase	Induction Treatment D-VRd (Cycles 1-4, 21-Day Cycle) Upon enrollment			Apheresis <i>Note: Occurs after Cycle 1 or 2, before Cycle 3</i> Minimum of 21 days after last dose of D-VRd	Peripheral Stem Cell Mobilization and Harvesting (Optional) <sup>c</sup>	Cyclophosphamide and fludarabine conditioning regimen Upon recovery from C4 Induction therapy; Minimum of 21 days after last dose of D-VRd
		D1	D8 (±2 days)	D15 (±2 days)			
	≤28 days prior to Induction Treatment <sup>a</sup>					After apheresis (Post-Induction Cycle 2, 3 or 4)	<b>Day -5,* -4, -3<sup>b</sup></b> <b>*Window of Day -7 to Day -5</b>
<b>Screening Assessments</b>							
Informed consent <sup>a</sup>	X Before the 1 <sup>st</sup> study related procedure						
Eligibility criteria	X						
Demography, Medical History (including neurological history)	X						
Disease Characteristics	X						
ECOG performance status	X	C3 only					Prior to 1 <sup>st</sup> dose only
Myeloma Frailty Index	X						
12-lead ECG	X	As clinically indicated					
Physical examination	X	Symptom and disease directed examination as clinically indicated					
Height	X						
Spirometry (subjects with known COPD only)	X						
Echocardiogram or MUGA scan	X (≤8 weeks before induction treatment)						As clinically indicated if the subject develops signs/symptoms of heart failure
<b>Outpatient Administration</b> In consultation with and approval of the sponsor. See <a href="#">Attachment 15</a> for outpatient monitoring							
Evaluation for outpatient suitability (See <a href="#">Attachment 14</a> )					X		
<b>Assessments Prior to Apheresis and Conditioning Regimen</b>							
Criteria for Apheresis (See Section 6.1.1)					X; After recovery from 1 or 2 cycles of D-VRd induction and ≤72 hours prior to apheresis		
Criteria for Conditioning Regimen (See Section 6.1.2.1)							≤ 72 hours of the 1 <sup>st</sup> dose only



**Table 5: Cohort E: Time and Events Schedule for Study Procedures/Assessments (Screening to Conditioning Regimen)**

	Screening Phase	Induction Treatment <b>D-VRd</b> (Cycles 1-4, 21-Day Cycle) Upon enrollment			Apheresis <i>Note: Occurs after Cycle 1 or 2, before Cycle 3</i> Minimum of 21 days after last dose of D-VRd	Peripheral Stem Cell Mobilization and Harvesting (Optional) <sup>c</sup>	Cyclophosphamide and fludarabine <b>conditioning regimen</b> Upon recovery from C4 Induction therapy; Minimum of 21 days after last dose of D-VRd
		D1	D8 (±2 days)	D15 (±2 days)			
	≤28 days prior to <b>Induction Treatment<sup>a</sup></b>					After apheresis (Post-Induction Cycle 2, 3 or 4)	<b>Day -5,* -4, -3<sup>b</sup></b> <b>*Window of Day -7 to Day -5</b>
<b>Study Intervention and Administration</b>							
Weight	X	X			X (for JNJ-68284528 dose calculation)		Prior to 1 <sup>st</sup> dose only
Vital signs, including oxygen saturation	X	X	X	X	X	As clinically indicated; per investigator discretion or local guidelines	X
Apheresis					X		
Cyclophosphamide and G-CSF <sup>c</sup>						For stem cell mobilization per local standard of care	
Cyclophosphamide and fludarabine							X
<b>D-VRd Induction Regimen</b>							
<b>Pre-Daratumumab Medications (administer 1-3 hours before daratumumab)</b>							
Dexamethasone 20 mg <sup>d</sup>		X	C1-2 only	C1-2 only			
Diphenhydramine 25-50 mg (or equivalent)		X	C1-2 only	C1-2 only			
Paracetamol 650-1000 mg;		X	C1-2 only	C1-2 only			
Montelukast 10 mg (recommended prior to daratumumab C1D1 and is optional before all other doses)		C1 only					
<b>D-VRd</b>							
Dexamethasone 20 mg, oral or intravenous <sup>m</sup>		On Days 1, 2, 4, 5, 8, 9, 11, 12					

**Table 5: Cohort E: Time and Events Schedule for Study Procedures/Assessments (Screening to Conditioning Regimen)**

	Screening Phase	Induction Treatment D-VRd (Cycles 1-4, 21-Day Cycle)			Apheresis <i>Note: Occurs after Cycle 1 or 2, before Cycle 3</i> Minimum of 21 days after last dose of D-VRd	Peripheral Stem Cell Mobilization and Harvesting (Optional) <sup>c</sup>	Cyclophosphamide and fludarabine conditioning regimen Upon recovery from C4 Induction therapy; Minimum of 21 days after last dose of D-VRd
		Upon enrollment					
	≤28 days prior to Induction Treatment <sup>a</sup>	D1	D8 (±2 days)	D15 (±2 days)		After apheresis (Post-Induction Cycle 2, 3 or 4)	Day -5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5
Bortezomib 1.3 mg/m <sup>2</sup> ; administer by subcutaneous injection		On Days 1, 4, 8, 11 of each cycle					
Lenalidomide 25 mg, oral		On Day 1-14 of each cycle					
Daratumumab 1800 mg SC; Refer to SIPP or approved prescribing information for drug preparation and administration recommendations		On Days 1, 8, 15 for Cycles 1-2 [weekly] then Day 1 of Cycles 3-4					
Pill count for lenalidomide <sup>o</sup>		X					
Pill count for dexamethasone <sup>o</sup>		X					
<b>Laboratory Assessments (Section 9.7)</b> To be performed by the local laboratory except for the calcium and albumin-adjusted calcium, which will be performed at the central laboratory (local labs may be used to assess eligibility). Blood samples collection may be performed at the participant's home by mobile study personnel (i.e. nurses and mobile phlebotomist).							
Blood group and type and indirect antiglobulin test (IAT) results		Pre- dose CID1 <sup>c</sup>					
Hematology <sup>o</sup>	X	X	X	X	X (≤72 hours prior to apheresis )		Prior to 1st dose only (≤72 hour window)
Chemistry	X	X		X	X (≤72 hours window)		Prior to 1st dose only (≤72 hour window)
Serology <sup>f</sup>	X						
Coagulation (PT/INR, aPTT, fibrinogen, D-dimer)	X						
Serum Pregnancy test (in subjects with childbearing potential)	X	Within 24 hours prior to CID1	Weekly during Cycle 1 and then monthly in women with regular menstrual cycles or every 2 weeks in women with irregular menstrual cycles		X (≤72 hour window)		Prior to 1 <sup>st</sup> dose only (≤72 hour window)

**Table 5: Cohort E: Time and Events Schedule for Study Procedures/Assessments (Screening to Conditioning Regimen)**

	Screening Phase	Induction Treatment D-VRd (Cycles 1-4, 21-Day Cycle) Upon enrollment			Apheresis <i>Note: Occurs after Cycle 1 or 2, before Cycle 3</i> Minimum of 21 days after last dose of D-VRd	Peripheral Stem Cell Mobilization and Harvesting (Optional) <sup>c</sup>	Cyclophosphamide and fludarabine conditioning regimen Upon recovery from C4 Induction therapy; Minimum of 21 days after last dose of D-VRd
		D1	D8 (±2 days)	D15 (±2 days)			
	≤28 days prior to Induction Treatment <sup>a</sup>					After apheresis (Post-Induction Cycle 2, 3 or 4)	<b>Day -5,* -4, -3<sup>b</sup></b> <b>*Window of Day -7 to Day -5</b>
Infectious Disease Testing <sup>g</sup>					X (within 60 days of apheresis, as applicable per local regulations)		
<b>Serum and Urine Disease Evaluations</b> (See Section 9.2 for efficacy assessments. Blood and 24-hour urine: to be sent to the central laboratory. Disease evaluation should continue to be performed until confirmed disease progression, death, start of a new anticancer treatment, withdrawal of consent for study participation, or study completion, whichever occurs first.). Subjects with disease progression who receive retreatment with JNJ-68284528 must continue with disease evaluation visits.							
Serum β2-microglobulin	X						
Quantitative Immunoglobulins	X <sup>h</sup>						X (prior to first dose [≤7 days] <sup>h</sup> )
Serum M-protein quantitation by electrophoresis	X	X					X (prior to first dose [≤7 days])
24-hour urine protein electrophoresis sample	X <sup>i</sup>	X					X (prior to first dose [≤7 days])
Serum calcium corrected for albumin	X	X					X (prior to first dose [≤7 days])
Serum FLC and serum/urine immunofixation	X	X					Serum FLC and serum/urine immunofixation are to be performed prior to the start of conditioning regimen (Day -5 [≤7 days]) and when CR is suspected or maintained; for subjects with measurable disease only by light chain criteria serum FLC will also be performed at every assessment when an SPEP is performed
<b>Other Disease Evaluations</b>							
MRD (bone marrow aspirate) <sup>j</sup>	X (performed during screening)						X (prior to first dose [≤7 days])
Bone marrow aspirate and biopsy for disease evaluation	X						X (prior to first dose [≤7 days])
Skeletal Survey <sup>k</sup>	X (at screening, or within 14 days prior to the first dose of conditioning regimen)						
Assess extramedullary Plasmacytomas	X	If applicable, by physical exam Q4W, by radiologic exam (if required) Q12W					X (≤14 days prior to first dose) <sup>l</sup>

**Table 5: Cohort E: Time and Events Schedule for Study Procedures/Assessments (Screening to Conditioning Regimen)**

	Screening Phase	Induction Treatment D-VRd (Cycles 1-4, 21-Day Cycle)			Apheresis <i>Note: Occurs after Cycle 1 or 2, before Cycle 3</i> Minimum of 21 days after last dose of D-VRd	Peripheral Stem Cell Mobilization and Harvesting (Optional) <sup>c</sup>	Cyclophosphamide and fludarabine conditioning regimen Upon recovery from C4 Induction therapy; Minimum of 21 days after last dose of D-VRd
		Upon enrollment					
	≤28 days prior to Induction Treatment <sup>a</sup>	D1	D8 (±2 days)	D15 (±2 days)		After apheresis (Post-Induction Cycle 2, 3 or 4)	Day -5,* -4, -3 <sup>b</sup> *Window of Day -7 to Day -5
Biomarker evaluations	See Biomarker Time & Events Schedule (Table 7)						
<b>Patient Reported Outcomes (PRO):</b> To be completed before any clinical tests or procedures scheduled for the same day as the PRO assessments that would influence the subject's perceptions of their current health							
EORTC QLQ-C30	X				X (≤72 hour window)		
MySim-Q (Unless translation not available)	X				X (≤72 hour window)		
PGIS	X				X (≤72 hour window)		
PGIC							
PRO CTCAE Items	X				X (≤72 hour window)		
MRU		X					
<b>Ongoing Subject Review</b>							
Adverse Events	Continuous from the time of signing of ICF Second primary malignancies should be followed from the time of signing of ICF to study completion <sup>n</sup>						
Concomitant medication	Continuous from the time of signing of ICF						
Survival Follow-up	After disease progression is documented, survival status will be obtained every 16 weeks until study completion						
Subsequent Anticancer Therapy	After disease progression is confirmed, subsequent anticancer therapy will be obtained every 16 weeks until study completion						

Abbreviations: aPTT=activated partial thromboplastin time; CR=complete response; sCR=stringent complete response; CRS= cytokine release syndrome; CT=computed tomography; C=Cycle; D=Day; ECOG=Eastern Cooperative Oncology Group; ECG=electrocardiogram; EORTC-QLQ=European Organization for Research and Treatment of Cancer Quality of Life Questionnaire; FISH=fluorescence in situ hybridization; FLC=free light chain; IAT=indirect antiglobulin test; ICF=informed consent form; INR=international normalized ratio; IPPI=investigational product preparation instructions; MRD=minimal residual disease; MRI=magnetic resonance imaging; MUGA=multiple-gated acquisition; MySim-Q=Multiple Myeloma Symptom and Impact Questionnaire; PGIC=Patient Global Impression of Change; PGIS=Patient Global Impression of Severity; PRO=patient reported outcome; PT=prothrombin time; SC=subcutaneous; SIPP=site investigational product; w=weeks.

- <sup>a</sup> ICF must be signed before any study-related procedures are performed, and remains in effect even if the screening evaluation is not performed within the 28-day Screening Phase. Evaluations for eligibility determination performed outside the screening window may need to be repeated. For subjects who require a repeat apheresis see Section 9.1.3 for assessments should be collected before the second apheresis.
- <sup>b</sup> Window for the start of the conditioning regimen is Day -7 to Day -5.
- <sup>c</sup> Stem cell mobilization and harvesting is optional and per investigator preference. If chosen, stem cell mobilization may be performed using G-CSF with or without Plerixafor per local standard of care after Cycle 4 and stem cells will be harvested based on response to mobilization per institutional practice.
- <sup>d</sup> On daratumumab dosing days, dexamethasone pre-medication for daratumumab injection will replace the daily dose of dexamethasone (and should be administered 1 hour before the daratumumab infusion).
- <sup>e</sup> Blood type, Rh, and Indirect Antiglobulin Test (IAT) should be done before the first dose of daratumumab

- <sup>f</sup> Serology results performed as standard of care within 28 days prior to apheresis are acceptable. Hepatitis B: HBsAg, anti-HBc, anti-HBs, HBV DNA quantification (for subjects who are anti-HBs positive without a history of vaccination or for subjects who are anti-HBc positive with or without anti-HBs positive); Monitor HBV-DNA, AST/ALT every 3 months for one year post-dose of JNJ-68284528. Hepatitis C: HCV antibody, HCV RNA (for subjects who are anti HCV positive); HIV serology. See [Attachment 10](#).
- <sup>g</sup> HIV, hepatitis B, hepatitis C, HTLV, and other infectious diseases will be performed within 60 days of apheresis or as applicable per local regulations, whichever is more stringent
- <sup>h</sup> All subjects will be evaluated for IgG, IgA, IgM. Testing for IgD and IgE will only be performed for subjects with IgD and IgE-type myeloma.
- <sup>i</sup> UPEP sample collected as part of the standard of care and prior to the subject signing ICF may be used for analysis at the central laboratory.
- <sup>j</sup> Bone marrow morphology from an aspirate and core biopsy to be assessed locally at all time points. Additional bone marrow aspirate samples will be collected for biomarkers (See [Table 8](#))
- <sup>k</sup> Results from skeletal survey performed as routine follow-up within 42 days before start of induction therapy may be used without these tests being repeated (Section [9.2.5](#)). Additional imaging (X-ray, CT, or MRI) will be performed as clinically indicated (eg, to document response or progression).
- <sup>l</sup> Results from radiologic plasmacytoma assessments performed  $\leq 14$  days prior to the first dose of the conditioning as routine follow-up for subject's disease may be used. Extramedullary plasmacytomas should be assessed for all subjects with a history of plasmacytomas or if clinically indicated at screening, by clinical examination or radiologic imaging (Section [9.2.6](#)).
- <sup>m</sup> For subjects older than 75 years or underweight (BMI <18.5), the dexamethasone dose may be administered at a dose of 20 mg on days 1, 4, 8, and 11.
- <sup>n</sup> A tumor sample should be collected and DNA, RNA, or protein analysis may be performed to investigate the presence of lentiviral elements
- <sup>o</sup> Hematology, chemistry and pill count are to be done every 28 days past 12 months; all other assessments will be collected every 56 days or as otherwise noted.

**Table 6: Cohort E: Time and Events Schedule for Study Procedures/Assessments (JNJ-68284528 Infusion to Post-treatment)**

	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup>											Consolidation Treatment D+R (As early as day 21 and up to End of Cohort) <sup>a, b</sup>	
		Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>a</sup>	Lenalidomide
<b>Screening Assessments</b>														
ECOG performance status	X									X		X		
12-lead ECG	As clinically indicated													
Physical Examination	A symptom-directed physical examination should be performed as clinically indicated													
Echocardiogram or MUGA scan	As clinically indicated if the subject develops signs/symptoms of heart failure													
ICE neurologic test	X (≤24 hours prior to infusion) <sup>c</sup>	ICE test must be repeated at any incidence of suspected CAR-T cell-related neurotoxicity (eg, ICANS). Perform at least daily until resolved.												
Handwriting sample	X (≤24 hours prior to infusion) <sup>c</sup>	X	X	X	X	X	X	X	X	X	X	X	X	Perform monthly up to Day 184
<b>Outpatient Administration</b> In consultation with and approval of the sponsor. See <a href="#">Attachment 15</a> for outpatient monitoring														
Evaluation for outpatient suitability (See <a href="#">Attachment 14</a> )	X (predose)													
All Subjects with hospital discharge on Day 10				Daily phone calls during business hours from site staff, Days 11-14										
<b>Assessments Prior to Apheresis and Conditioning Regimen</b>														
Criteria for JNJ-68284528 Administration (See Section 6.1.3.2)	X (predose)													
<b>Study Intervention and Administration</b>														
Weight	X													
Vital signs, including oxygen saturation	X <sup>d</sup>	X	X	X	X	X	X			X			X	X
Temperature	Measure at least twice a day <sup>e</sup>													

**Table 6: Cohort E: Time and Events Schedule for Study Procedures/Assessments (JNJ-68284528 Infusion to Post-treatment)**

	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup>											Consolidation Treatment D+R (As early as day 21 and up to End of Cohort) <sup>a, b</sup>		
		Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>a</sup>	Lenalidomide	Daratumumab <sup>1</sup>
Pre-infusion medication (see Section 6.1.3.3 for requirements prior to dosing with JNJ-68284528)	X														
JNJ-68284528 (See SIPP and IPPI for administration of JNJ-68284528)	X														
<b>D+R Continued Treatment</b>															
<b>Pre-Daratumumab Medications (administer 1-3 hours before daratumumab)</b>															
Dexamethasone 20 mg															X
Diphenhydramine 25-50 mg (or equivalent);															X
Paracetamol 650-1000 mg;															X
Montelukast 10 mg (recommended prior to daratumumab C1D1 and is optional before all other doses)															X
<b>D+R</b>															
Lenalidomide						Initiate lenalidomide a minimum of 21 days post JNJ-68284528 after resolution of associated CRS or neurological toxicities. Refer to <a href="#">Table 21</a> and Section 6.1.5.2 for additional dosing considerations. Lenalidomide will be taken daily (21 days on, 7 days off) in 28-day cycles, until PD, unacceptable toxicities or until a maximum of 2 years.									

**Table 6: Cohort E: Time and Events Schedule for Study Procedures/Assessments (JNJ-68284528 Infusion to Post-treatment)**

	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup>											Consolidation Treatment D+R (As early as day 21 and up to End of Cohort) <sup>a, b</sup>				
		Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>a</sup>	Lenalidomide	Daratumumab <sup>j</sup>		
Daratumumab 1800 mg SC; Refer to SIPPm for drug preparation and administration recommendations															Initiate daratumumab approximately 60-100 days after JNJ-68284528 and upon adequate recovery from ongoing toxicities. Administer every 4 weeks until disease progression, unacceptable toxicities or end of study treatment		
Pill count for lenalidomide <sup>k</sup>														X			
<b>Laboratory Assessments (See Section 9.7).</b> To be performed by the local laboratory except for the calcium and albumin-adjusted calcium, which will be performed at the central laboratory (local labs may be used to assess eligibility). Blood samples collection may be performed at the participant’s home by mobile study personnel (i.e. nurses and mobile phlebotomist) in the post-treatment period, after the Day 100.																	
Hematology <sup>k</sup>	X (predose)	X	X	X	X	X	X		X	X	X	X	X	X (prior to each dose of daratumumab and prior to each cycle of lenalidomide)			
Chemistry <sup>k</sup>	X (predose)	X	X	X	X	X	X		X	X	X	X	X	X (prior to each cycle)			
Serology <sup>f</sup>		For subjects at risk for HBV activation monitor HBV DNA and AST/ALT every 12 weeks (±7 days) for 12 months post JNJ-68284528 dose <sup>f</sup>															
Coagulation (PT/INR, aPTT, fibrinogen, D-dimer)	As clinically indicated for subjects who have fever or other signs of potential CRS																
Serum Pregnancy test (in subjects with childbearing potential)						Within 24 hours prior to start of lenalidomide, every week for the first 4 weeks and repeated at a maximum of at least every 28 days (or every 14 days for women of child bearing potential with irregular menses) Pregnancy testing must be performed 4 weeks after stopping lenalidomide. Additional pregnancy testing is done as clinically indicated or consistent with any country specific requirements listed in the local prescribing information for lenalidomide, whichever is more stringent											
<b>Serum and Urine Disease Evaluations (See Section 9.2 for efficacy assessments.</b> Blood and 24-hour urine: to be sent to the central laboratory. Disease evaluation should continue to be performed until confirmed disease progression, death, start of a new anticancer treatment, withdrawal of consent for study participation, or study completion, whichever occurs first.). Subjects with disease progression who receive retreatment with JNJ-68284528 must continue with disease evaluation visits.																	
Quantitative Immunoglobulins							X				X	X	X	X	X		



**Table 6: Cohort E: Time and Events Schedule for Study Procedures/Assessments (JNJ-68284528 Infusion to Post-treatment)**

	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup>											Consolidation Treatment D+R (As early as day 21 and up to End of Cohort) <sup>a, b</sup>	
		Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>a</sup>	Lenalidomide
Serum M-protein quantitation by electrophoresis							X			X	X	X	X	
24-hour urine protein electrophoresis sample							X			X	X	X	X	
Serum calcium corrected for albumin							X			X	X	X	X	
Serum FLC and serum/urine immunofixation	Serum FLC and serum/urine immunofixation are to be performed prior to the start of conditioning regimen (Day -5 [ $\leq 7$ days]) and when CR is suspected or maintained; for subjects with measurable disease only by light chain criteria serum FLC will also be performed at every assessment when an SPEP is performed													
<b>Other Disease Evaluations</b>														
MRD (bone marrow aspirate) <sup>g</sup>	Sample should be collected: <ul style="list-style-type: none"> <li>For all dosed subjects at Day 56, and at 6 month, 12 month, 18 month (Day 520), 24 (Day 744) months (<math>\pm 16</math> days) regardless of the status of disease measured in blood and urine</li> <li>For subjects with suspected CR at the time of CR and then yearly for subjects that remain on study up to disease progression.</li> </ul>													
Bone marrow aspirate and core biopsy for disease evaluation	To confirm CR, sCR, and at disease progression (immunohistochemistry or immunofluorescence). Can be taken at the same time at MRD sample if needed.													
Skeletal Survey <sup>h</sup>	As clinically indicated to document disease progression or response.													
Assess extramedullary Plasmacytomas <sup>i</sup>	Measurable sites Day 28, Day 56, Day 78, Day 100 then every 4 weeks for physical examination (if applicable) and Day 78 and Day 156 then every 12 weeks for radiologic assessment (for subjects with a history of plasmacytomas or as clinically indicated for others).													
MRD assessment by imaging (optional)	Must be recorded in the CRF													
Biomarker evaluations	See Biomarker Time & Events Schedule (Table 7 and Table 8)													
<b>Patient Reported Outcomes (PRO):</b> To be completed before any clinical tests or procedures scheduled for the same day as the PRO assessments that would influence the subject's perceptions of their current health														
EORTC QLQ-C30			X				X			X	X	X		X; every 4 months ( $\pm 10$ days)
MySiM-Q (Unless translation not available)			X				X			X	X	X		X; every 4 months ( $\pm 10$ days)
PGIS			X				X			X	X	X		X; every 4 months ( $\pm 10$ days)
PGIC							X			X	X	X		
PRO CTCAE			X				X			X	X	X		
MRU	X				X		X			X		X		X; every 28 days until Day 180

**Table 6: Cohort E: Time and Events Schedule for Study Procedures/Assessments (JNJ-68284528 Infusion to Post-treatment)**

	JNJ-68284528 Infusion	Post Infusion (Day 1 to Day 100) (any subject who received an infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup>											Consolidation Treatment D+R (As early as day 21 and up to End of Cohort) <sup>a, b</sup>	
		Day 1 (Infusion)	Day 3	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days) <sup>a</sup>	Lenalidomide
<b>Ongoing Subject Review</b>														
Adverse Events	Continuous from the time of signing of ICF until 100 days after last administration of JNJ-68284528, or 30 days after last dose of D+R, whichever is later. Beyond the adverse event reporting period, adverse events that are considered related to study drug need to be reported until the end of the study. Collect new neurologic adverse events or exacerbation of existing neurologic adverse events up to 12 months after JNJ-68284528 infusion. Second primary malignancies should be followed from the time of signing of ICF to study completion <sup>1</sup> CRS should be evaluated according to the ASBMT consensus grading (Lee 2019) (Attachment 2) CAR-T cell-related neurotoxicity (eg, ICANS) should be graded according to the ASBMT consensus grading (Attachment 4) Events of HBV reactivations should be reported during the first year post-dosing of JNJ-68284528													
Concomitant medication	Continuous from the time of signing of ICF until at least 100 days after administration of JNJ-68284528, or 30 days after last dose of D+R, whichever is later. Thereafter, continue to report concomitant therapy given for any Adverse Events considered related to study drug until the end of the study.													
Survival Follow-up	After disease progression is documented, survival status will be obtained every 16 weeks until study completion													
Subsequent Anticancer Therapy	After disease progression is documented, subsequent anticancer therapy will be obtained every 16 weeks until study completion													

Abbreviations: aPTT=activated partial thromboplastin time; CR=complete response; sCR=stringent complete response; CRS= cytokine release syndrome; CT=computed tomography; C=Cycle; D=Day; D+R=daratumumab, lenalidomide; ECOG=Eastern Cooperative Oncology Group; ECG=electrocardiogram; EORTC-QLQ=European Organization for Research and Treatment of Cancer Quality of Life Questionnaire; FISH=fluorescence in situ hybridization; FLC=free light chain; ICF=informed consent form; INR=international normalized ratio; IPPI=investigational product preparation instructions; MRD=minimal residual disease; MRI=magnetic resonance imaging; MUGA=multiple-gated acquisition; MySIm-Q=Multiple Myeloma Symptom and Impact Questionnaire; PGIC=Patient Global Impression of Change; PGIS=Patient Global Impression of Severity; PRO=patient reported outcome; PT=prothrombin time; SC=subcutaneous; SIPPM=site investigational product; w=weeks.

- <sup>a</sup> Subjects who discontinue after Day 100 but before study completion should have urine and serum disease assessments performed prior to withdrawal if feasible at the time of discontinuation, unless performed within 14 days prior to discontinuation.
- <sup>b</sup> Post-treatment assessments to be obtained until progressive disease is documented or the start of subsequent anticancer therapy, with the exception of survival status and subsequent anticancer therapy, which are to be collected until study completion. PRO assessments are collected until study completion (continued after disease progression or subsequent anticancer therapy).
- <sup>c</sup> Pre-infusion ICE test and handwriting sample should be performed before pre-medication with diphenhydramine.
- <sup>d</sup> Immediately before the start of infusion, at the end of infusion, and 0.5, 1, 2 hours after end of infusion. Monitor until normalized after a CRS event.
- <sup>e</sup> Temperature will be checked at least twice a day up to Day 28. Subjects will be provided with a thermometer and instructed on the use of the thermometer and entering 2 temperatures including their maximum daily temperature in a diary. Diary will be reviewed at each visit, then collected on Day 28 and stored with subject source documents.
- <sup>f</sup> Hepatitis B: HBsAg, anti-HBc, anti-HBs, HBV DNA quantification (for subjects who are anti-HBs positive without a history of vaccination or for subjects who are anti-HBc positive with or without anti-HBs positive); Only in subjects who are positive for antibodies to hepatitis B core antigen (anti-HBc) with or without positive antibodies to hepatitis B surface antigen (anti-HBs): Monitor HBV-DNA, AST/ALT every 12 weeks (±7 days) for one year post-dose of JNJ-68284528. Hepatitis C: HCV antibody, HCV RNA (for subjects who are anti HCV positive); HIV serology. See Attachment 10

- <sup>g</sup> Bone marrow morphology from an aspirate and core biopsy to be assessed locally at all time points. Additional bone marrow aspirate samples will be collected for biomarkers (See [Table 8](#))
- <sup>h</sup> Results from skeletal survey performed as routine follow-up within 42 days before start of induction therapy may be used without these tests being repeated (Section 9.2.5). Additional imaging (X-ray, CT, or MRI) will be performed as clinically indicated (eg, to document response or progression).
- <sup>i</sup> Results from radiologic plasmacytoma assessments performed  $\leq 14$  days prior to the first dose of the conditioning as routine follow-up for subject's disease may be used. Extramedullary plasmacytomas should be assessed for all subjects with a history of plasmacytomas or if clinically indicated at screening, by clinical examination or radiologic imaging (Section 9.2.6).
- <sup>j</sup> The start of daratumumab should fall on Day 1 of a lenalidomide cycle to ensure cycle-day consistency. Subjects must complete 1 cycle of lenalidomide before starting daratumumab, even if hematologic recovery occurs sooner.
- <sup>k</sup> Hematology, chemistry and pill count are to be done every 28 days past 12 months; all other assessments will be collected every 56 days or as otherwise noted.
- <sup>l</sup> A tumor sample should be collected and DNA, RNA, or protein analysis may be performed to investigate the presence of lentiviral elements

**Table 7: Cohort E: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling (Screening to Conditioning Regimen)**

	Screening Phase	Induction Treatment <b>D-VRd</b> (Cycles 1-4, 21-Day Cycle) <i>Note: Cycle 4 will be administered during CAR-T cell manufacturing.</i> T = Induction treatment			Apheresis <i>Note: Occurs after Cycle 1 or 2, before Cycle 3</i> Minimum of 21 days after last dose of D-VRd	Cyclophosphamide and fludarabine conditioning regimen (Upon recovery from C4 Induction therapy)
	≤28 days Prior to Induction Treatment	D1T	D8T (±2 days)	D15T (±2 days)		Day -5,* -4, -3 Assessments may be conducted ≤72 hours pre-dose. <b>*Window of Day -7 to Day -5</b>
<b>Pharmacokinetics</b>						
PK CAR transgene levels blood sample						X (prior to first dose [≤7 days])
Soluble serum BCMA sample						X (prior to first dose [≤7 days])
PK CAR transgene levels bone marrow sample						X (prior to first dose [≤7 days])
Daratumumab PK sample (serum)		Pre-dose C1D1T; Pre-dose C3D1T			X (≤72 hours window)	
<b>Biomarker Sampling</b>						
Immuno-phenotyping (whole blood)	X	Pre-dose Day 1 of each cycle			X (≤72 hours window)	X (prior to first dose [≤7 days])
Flow cytometry, (aspirate) (bone marrow)	X (done during screening)	Pre-dose C1D1T				X (prior to first dose [≤7 days])
CyTOF (aspirate) (bone marrow)						X (prior to first dose [≤7 days])
CyTOF/ PBMC/ Plasma (whole blood)	X				X(≤72 hours window)	
PBMCs for functional in vitro assays					X (≤72 hours window)	
Cytogenetics (bone marrow)	X					
Whole Exome Sequencing (bone marrow)	X					
Replication Competent Lentivirus (RCL) (whole blood)						X (prior to first dose [≤7 days])
Serum protein analysis		Pre-dose C1D1T				X (prior to first dose [≤7 days])
Cytokine profiling (serum)						X (prior to first dose [≤7 days])

Abbreviations: BCMA=B-cell maturation antigen; CAR=chimeric antigen receptor; CR = complete response; CRS=cytokine release syndrome; CyTOF=cytometry by time of flight; PBMC=peripheral blood mononuclear cell; PD=progressive disease; PK=pharmacokinetic; sCR=stringent complete response

**Table 8: Cohort E: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling (JNJ-68284528 Infusion to Disease Progression or Study Completion)**

	JNJ-68284528 Infusion	Post Infusion (any subject who received infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup> and Post-treatment (Day 101 up to study completion) <sup>#</sup>													At PD	At Study Completion for subjects without PD
		Day 1 (Infusion)	Day 2 (± 2 hour)	Day 3 (± 4 hour)	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days)		
<b>Pharmacokinetics</b>																
PK CAR transgene levels blood sample <sup>b</sup>	Pre-dose (≤4 hr window); Post end of infusion (within 30 minutes)	24-hour post-end of infusion	X	X	X	X	X	X		X	X	X	X;	then every 8 weeks up to 1 year	X	X
Soluble serum BCMA sample	Pre-dose (≤4 hr window); Post end of infusion	24-hour post-end of infusion	X	X	X	X	X	X		X	X	X	X;	then every 8 weeks up to 1 year	X	X
PK CAR transgene levels bone marrow sample								X			X			X		
ADA sample (serum) <sup>b,c</sup>	X (Pre-dose)					X		X			X	X	X	X	X	X
Daratumumab PK sample (serum)	X (Pre-dose)				X			X			X					
<b>Biomarker Sampling</b>																
Immuno-phenotyping (whole blood) <sup>d,f</sup>	X Pre-dose (≤4 hour window)	24 hour post end of infusion		X	X	X	X	X	X	X	X	X	X;	then every 8 weeks up to 1 year	X	X <sup>d</sup>
Flow cytometry, (aspirate) (bone marrow) <sup>d</sup>								X			X			X	X	X

**Table 8: Cohort E: Time and Events Schedule for Pharmacokinetic and Biomarker Sampling (JNJ-68284528 Infusion to Disease Progression or Study Completion)**

	JNJ-68284528 Infusion	Post Infusion (any subject who received infusion of JNJ-68284528 should continue all subsequent assessments) <sup>a</sup> and Post-treatment (Day 101 up to study completion) <sup>#</sup>													At PD	At Study Completion for subjects without PD
	Day 1 (Infusion)	Day 2 (± 2 hour)	Day 3 (± 4 hour)	Day 7 (± 1 day)	Day 10 (± 1 day)	Day 14 (± 1 day)	Day 21 (± 1 day)	Day 28 (± 2 days)	Day 35 (± 2 days)	Day 42 (± 2 days)	Day 56 (± 2 days)	Day 78 (± 2 days)	Day 100 (± 2 days)	Day 184 (± 7 days)		
CyTOF (aspirate) (bone marrow) <sup>d,e</sup>								X		X	X			X	X	X
CyTOF/ PBMC/ Plasma (whole blood) <sup>d,e</sup>				X	X	X	X	X	X	X	X			X	X	X
PBMCs for functional in vitro assays					X			X		X	X		X; then every 8 weeks up to 1 year			
Cytogenetics (bone marrow)															X	
Whole Exome Sequencing (bone marrow)															X	
Replication Competent Lentivirus (RCL) (whole blood)	Pre-dose (≤4 hour window)	At approximately 3, 6, and 12 months; then yearly for 15 years post infusion														
Serum protein analysis	Pre-dose (≤4 hour window)	Additional serum protein sample will be taken at each SIFE/SPEP evaluation, as well as at each MRD sample collection														
Cytokine profiling <sup>f</sup> (serum)	Pre-dose, 2hrs Post (±10 minutes)	X	X	X	X	X	X	X		X	X	X	X			

Abbreviations: ADA=anti-drug antibody; BCMA=B-cell maturation antigen; CAR=chimeric antigen receptor; CR = complete response; CRS=cytokine release syndrome; CyTOF=cytometry by time of flight; ICANS= immune effector cell-associated neurotoxicity syndrome ; MRD=minimal residual disease; PBMC=peripheral blood mononuclear cell; PD=progressive disease; PK=pharmacokinetic; sCR=stringent complete response; SIFE=serum immunofixation electrophoresis; SPEP=serum protein electrophoresis

- <sup>a</sup> For subjects who discontinue the study before Day 100, the Day 100 assessments should be performed if feasible.
- <sup>b</sup> Collect additional samples when any of the following are observed or reported: 1) CRS or neurotoxicity event Grade  $\geq 2$  (at onset of the event, and 24 and 72 hours after) or as clinically indicated; and 2) as indicated based on emerging data
- <sup>c</sup> ADA sample should be collected if a subject withdraws from the study after JNJ-68284528 administration but prior to disease progression or study completion.
- <sup>d</sup> Sample should be collected at suspected CR
- <sup>e</sup> Sample should be collected at 12 months, relative to Day 1, for subjects that achieve CR/sCR and remain on study.
- <sup>f</sup> Collect additional samples when any of the following are suspected or reported: 1) CRS or CAR-T cell-related neurotoxicity (eg, ICANS) (any grade) (at onset of the event, and then every 24 hours until CRS or ICANS event has stabilized or is resolving at which time additional collections should occur at 24, 48, and 72 hours) or as clinically indicated; and 2) as indicated based on emerging data.
- <sup>g</sup> Pharmacokinetic and biomarker samples collection (serum samples only) may be performed at the subject's home by mobile study personnel (i.e. nurses and mobile phlebotomist) in the post-treatment period, after the Day 100.

**ABBREVIATIONS**

ADC	antibody-drug conjugate
ADL	Activities of Daily Living
AL	amyloid light-chain
ALC	absolute lymphocyte count
ALL	acute lymphocytic leukemia
ALT	alanine aminotransferase
ANC	absolute neutrophil count
APRIL	a proliferation inducing ligand
ASBMT	American Society for Blood and Bone Marrow Transplantation
ASCO	American Society of Clinical Oncology
ASCT	autologous stem cell transplantation
AST	aspartate aminotransferase
ASTCT	American Society for Transplantation and Cellular Therapy
AUC	area under the curve
BAFF	B-cell activating factor
BCMA	B-cell maturation antigen
β2M	beta 2 microglobulin
β-hCG	β human chorionic gonadotropin
BiPAP	Bilevel Positive Airway Pressure
BiTE	bispecific T-cell engager
BMI	body mass index
BNP	B-type natriuretic peptide
CABG	coronary artery bypass graft
CAR-T	chimeric antigen receptor T (cells)
CBC	complete blood count
CBR	clinical benefit rate
CI	confidence interval
CNS	central nervous system
COPD	chronic obstructive pulmonary disease
CPAP	Continuous Positive Airway Pressure
CR	complete response
CRES	CAR-T cell-related encephalopathy syndrome
CRF	case report form
CRS	cytokine release syndrome
CSR	clinical study report
CT	computed tomography
CTCAE	Common Terminology Criteria for Adverse Events
CytoF	cytometry by time of flight
DLT	dose limiting toxicity
DMSO	dimethyl sulfoxide
DOR	duration of response
D+R	daratumumab, lenalidomide
DTT	dithiothreitol
D-VRd	daratumumab, bortezomib, lenalidomide, dexamethasone
ECG	electrocardiogram
ECOG	Eastern Cooperative Oncology Group
eCRF	electronic case report form
eDC	electronic data capture
eGFR	estimated glomerular filtration rate
EORTC	European Organization for Research and Treatment of Cancer
FLC	free light chain
GCP	Good Clinical Practice
G-CSF	granulocyte colony-stimulating factor
GVHD	graft-versus-host disease
HBsAg	hepatitis B surface antigen



---

HBV	hepatitis B virus
HCV	hepatitis C virus
HLH/MAS	hemophagocytic lymphohistiocytosis/macrophage activation syndrome
HIV	human immunodeficiency virus
HRQoL	health-related quality of life
hr-NDMM	high risk, newly diagnosed multiple myeloma
IAT	Indirect Antiglobulin Test
ICANS	immune effector cell-associated neurotoxicity syndrome
ICE	immune effector cell-associated encephalopathy
ICF	informed consent form
ICH	International Council for Harmonisation of Technical Requirements for Pharmaceuticals for Human Use
ICP	intracranial pressure
IEC	independent ethics committee
Ig	immunoglobulins
IL	interleukin
IMiD	immunomodulatory drug
IMWG	International Myeloma Working Group
IPPI	investigational product preparation instructions
IRB	Institutional Review Board
IRR	infusion-related reaction
ISS	International Staging System
IV	intravenous(ly)
LDH	lactate dehydrogenase
LVEF	left ventricular ejection fraction
MedDRA	Medical Dictionary for Regulatory Activities
MR	minimal response
MRD	minimal residual disease
MRI	magnetic resonance imaging
MUGA	multiple-gated acquisition
MySim-Q	Multiple Myeloma Symptom and Impact Questionnaire
NGS	next generation sequencing
NYHA	New York Heart Association
ORR	overall response rate
OS	overall survival
PBMC	peripheral blood mononuclear cell
PD	progressive disease
PET	positron emission tomography
PFS	progression-free survival
PGIC	Patient Global Impression of Change
PGIS	Patient Global Impression of Severity
PI	proteasome inhibitor
POEMS	polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, and skin changes
PQC	product quality complaint
PR	partial response
PRO	patient reported outcomes
QIg	quantitative immunoglobulin
RBC	red blood cell
RCL	replication competent lentivirus
REMS	Risk Evaluation and Mitigation Strategy
rHuPH20	recombinant human hyaluronidase
RP2D	recommended Phase 2 dose
sBCMA	soluble BCMA
SC	subcutaneous
sCR	stringent complete response
SET	safety evaluation team
SIPPM	site investigational product procedures manual
SPEP	serum protein electrophoresis

---

SPM	second primary malignancy
SVR	sustained virologic response
TCR	T cell receptor
TEAE	treatment-emergent adverse event
TLS	tumor lysis syndrome
TNF-R	tumor necrosis factor receptor
TTR	time to response
ULN	upper limit of normal
UPEP	urine M-protein quantitation by electrophoresis
VGPR	very good partial response

## 1. INTRODUCTION

JNJ-68284528 (ciltacabtagene autoleucl) is an autologous chimeric antigen receptor T cell (CAR-T) therapy that targets B cell maturation antigen (BCMA), a molecule expressed on the surface of mature B lymphocytes and malignant plasma cells. Results from the Phase 1b portion of study 68284528MMY2001 indicate that JNJ-68284528 has a safety profile consistent with the known mechanism of action of the product and compelling efficacy in a population of heavily pre-treated subjects.

For the most comprehensive nonclinical and clinical information regarding JNJ-68284528, refer to the latest version of the Investigator's Brochure and Addenda for JNJ-68284528.

The term "sponsor" used throughout this document refers to the entities listed in the Contact Information page(s), which will be provided as a separate document.

### 1.1. Background

#### 1.1.1. Multiple Myeloma

Multiple myeloma is characterized by the production of monoclonal immunoglobulin (Ig) proteins or protein fragments (M proteins) that have lost their function (Kyle 2009; Palumbo 2011). The proliferation of multiple myeloma cells leads to subsequent displacement of normal bone marrow hematopoietic precursors and overproduction of M-proteins. Hallmarks of multiple myeloma include osteolytic lesions, anemia, increased susceptibility to infections, hypercalcemia, renal insufficiency or failure, and neurologic complications (Korde 2011; Palumbo 2011).

Treatment options for multiple myeloma have substantially improved over time and vary depending on the aggressiveness of the disease, underlying prognostic factors, physical condition of the patient, and existing co-morbidities. Therapeutic options include agents such as proteasome inhibitors (PIs), immunomodulatory drugs (IMiDs), monoclonal antibodies, and stem cell transplantation.

Despite these therapeutic achievements, the disease recurs and remains incurable. Thus, there is a need for novel therapeutic approaches.

#### 1.1.2. B cell Maturation Antigen

B cell maturation antigen (BCMA, also known as CD269 and TNFRSF17) is a 20 kilodalton, type III membrane protein that is part of the tumor necrosis receptor superfamily (Tai 2015). BCMA is predominantly expressed in B-lineage cells and plays a critical role in B cell maturation and subsequent differentiation into plasma cells (Tai 2015). BCMA binds 2 ligands that induce B cell proliferation: a proliferation inducing ligand (APRIL; CD256) and B-cell activating factor (BAFF; CD257) (Avery 2003; Darce 2007; Patel 2004). Upon binding of BCMA monomers to the APRIL trimer, activation and phosphorylation of p38MAPK, ELK, and NF- $\kappa$ B are triggered through intracellular tumor necrosis factor receptor (TNF-R)-associated factor (TRAF) molecules leading to pro-survival gene regulation (Bossen 2006; Hatzoglou 2000; Kimberley 2009).

In multiple myeloma cell lines and patient samples, BCMA is more stably expressed specifically on the B cell lineage than CD138, a key plasma cell marker which is also expressed on normal fibroblasts and epithelial cells (Palaiologou 2014). The expression characteristics of BCMA make it an ideal therapeutic target in the treatment of multiple myeloma (Frigyesi 2014; Tai 2015).

### 1.1.3. CAR-T Therapy


Chimeric antigen receptor T (CAR-T) cell therapy uses modified autologous T cells that are activated in a MHC-independent manner upon binding to their target. This results in lysis of the targeted cells. An ongoing Phase 1 clinical study with bb2121, a BCMA-directed CAR-T immunotherapy, demonstrated promising results for this strategy in relapsed/refractory multiple myeloma (Raje 2019). Of the first 33 consecutive subjects to receive an infusion of bb2121 76% experienced cytokine release syndrome (CRS), which was Grade 1 or 2 in 23 subjects (70%) and Grade 3 in 2 subjects (6%). Neurologic toxicity occurred in 42% of subjects. The objective response rate (partial response or better) was 85%, including 15 subjects (45%) with complete responses. A median progression-free survival (PFS) of 11.8 months was observed for the 30 subjects who received  $\geq 15 \times 10^7$  CAR-positive T-cells and 2.6 months for the 3 subjects who received  $< 15 \times 10^7$  CAR-positive T-cells. Sixteen subjects with a partial response or better and who could be evaluated for minimal residual disease (MRD), were MRD negative at the sensitivity threshold of  $10^{-4}$ .

### 1.1.4. JNJ-68284528

JNJ-68284528, consists of autologous CAR-T cells and the target antigen is BCMA. The novel design features dual targeting domains on BCMA, enabling tight binding of LCAR-B38M to the BCMA-expressing cells.

The LCAR-B38M coding sequence is comprised of a human CD8 alpha signal peptide (CD8 $\alpha$  SP), BCMA targeting domain consisting of 2 different VHH (single domain antibody, clone VHH1 and VHH2), human CD8 alpha hinge and transmembrane domain (CD8 $\alpha$  hinge+TM), human CD137 cytoplasmic domain, and a human CD3 zeta cytoplasmic domain (CD3 $\zeta$ ) (Figure 1). The expression of LCAR-B38M is driven and controlled by a human elongation factor 1 alpha promoter (hEF1 $\alpha$  promoter).

**Figure 1: LCAR-B38M Coding Region**



The JNJ-68284528 drug product used in this study and the LCAR-B38M CAR-T cell drug product used in the first-in-human Legend-2 study (See Section 1.1.5) express an identical CAR protein. The JNJ-68284528 drug product is produced using modified manufacturing and scale-up processes compared with the Legend-2 study material. The LCAR-B38M CAR-T cell designation will be used when referring to results from the Legend-2 study. JNJ-68284528 will be used to reference

the drug product in Study 68284528MMY2003 (CARTITUDE-2). A similar manufacturing process will be utilized to generate the commercial product used in Cohort A of Study 68284528MMY2003.

### **1.1.5. Clinical Studies**

#### **Legend-2**

Legend-2 is a first-in-human, single-arm, open-label, multicenter study to determine the safety and efficacy of LCAR-B38M CAR-T cells used to treat subjects with relapsed or refractory multiple myeloma. Study enrollment was completed in November 2017; 74 subjects have been treated on this study. The clinical cutoff for the analysis presented here is 06 February 2018 with updated survival, efficacy data, and CRS grading provided as of 31 December 2018. As of the 31 December 2018 update, the median follow-up was 17.41 months (range: 0.4–32.2 months).

All subjects had relapsed or refractory multiple myeloma with a median of 3 (range: 1-9) prior lines of therapy including PI therapy (73%), IMiD (87.8%), and both PI and IMiD (64.9%). Prior autologous stem cell transplant (ASCT) was reported for 24.3% of subjects. The median age at study entry was 54.5 years (range: 27-74 years).

Of the 74 subjects included in the Legend-2 study, 68 (91.9%) subjects had an adverse event of CRS (median time to onset 9 days [range: 1-19 days]). The majority of CRS events were of Grade 1 or Grade 2 severity. Six (8.1%) subjects experienced Grade 3 CRS and 1 (1.4%) subject experienced Grade 5 CRS. The fatal event of CRS occurred in a subject who experienced CRS and tumor lysis syndrome (TLS) and died on Day 13 after receiving the LCAR-B38M CAR-T cell infusion. For most subjects, symptoms of CRS were mild and reversible. Grade 1 neurotoxicity was reported for 1 subject.

In addition to the subject who died from CRS, 1 subject had a fatal adverse event of potential acute pulmonary embolism and potential acute coronary syndrome on Day 22 after infusion of LCAR-B38M CAR-T cells.

As of the 31 December 2018 update, the ORR (complete response [CR] + very good partial response [VGPR] + partial response [PR]) was 87.8%. Complete response was achieved by 54 (73.0%) subjects and 49 (66.2%) subjects were negative for MRD as assessed by bone marrow 8-color flow cytometry assay.

As of the 31 December 2018 update, the Kaplan-Meier estimate of median duration of response was 22.34 months (95% confidence interval [CI]: 11.79-29.14). The median progression-free survival (PFS) for the overall population was 16.2 months (95% CI: 10.61-28.16) and among the 49 subjects achieving MRD-negative CR, the median PFS was 28.16 months.

#### **Study 68284528MMY2001**

As of the 24 June 2019 data cutoff, 25 subjects had received an infusion of JNJ-68284528 in Study 68284528MMY2001. The safety profile is consistent with observations from the Legend-2 study.

- Twenty-four of the 25 subjects who received JNJ-68284528 had at least 1 treatment-emergent adverse event (TEAE).
  - The most commonly reported TEAEs (>20% of subjects) were CRS (88.0%), neutropenia (80.0%), anemia (76.0%), thrombocytopenia (72.0%), leukopenia (40.0%), lymphopenia (28.0%), fatigue (24.0%), headache (24.0%), cough (24.0%), and diarrhea (20.0%).
- Grade 3 or 4 TEAEs were reported for all 24 of the 25 subjects.
  - The most commonly reported Grade 3 or 4 TEAEs (>10% of subjects) were neutropenia (76.0%), anemia (48%), thrombocytopenia (60.0%), leukopenia (40.0%), lymphopenia (16%), and CRS (16.0%).
- Serious adverse events were reported for 6 subjects:
  - One subject had serious TEAEs of Grade 3 CAR-T cell-related encephalopathy syndrome (CRES) and Grade 3 CRS.
  - One subject had serious TEAEs of: CRS, with Grade 5 hemophagocytic lymphohistiocytosis (HLH) secondary to CRS, and Grade 4 acute kidney injury.
  - One subject had serious TEAEs of Grade 4 thrombocytopenia and Grade 3 pneumonia
  - One subject had a serious TEAE of Grade 2 influenza A
  - One subject had a serious TEAE of Grade 3 encephalitis
  - One subject had serious TEAEs of Grade 1 CRS and Grade 1 confusional state
- One subject died due to CRS, with Grade 5 hemophagocytic lymphohistiocytosis secondary to CRS.
- Twenty-two of the 25 subjects (88.0%) experienced CRS. Three subjects experienced Grade 3 CRS and 1 subject experienced Grade 5 CRS.
- CAR-T cell-related neurotoxicity was reported in 4 subjects. One subject had Grade 3 CRES, the second subject had Grade 1 neurologic adverse events (dysarthria, slow mentation, gait disturbance, and somnolence), the third subject had Grade 1 CRES, and the fourth subject had a Grade 1 immune effector cell-associated neurotoxicity syndrome (ICANS) event (difficulty in finding words). All events occurred in the setting of CRS.

The median duration of follow-up for the 21 subjects who received JNJ-68284528 and had at least 1 post-dose disease evaluation as of 24 June 2019 was 2.99 months (range: 1.3 to 9.9 months). The ORR (sCR + CR + VGPR + PR) for the 21 subjects with at least 1 post-dose disease evaluation as of the 24 June 2019 cutoff was 90.5% (including unconfirmed responses). Thirteen subjects (61.9%) having VGPR or better and 6 subjects (28.6%) having CR or better. A stringent complete response (sCR) was achieved by 4 subjects. Fifteen subjects had post-baseline bone marrow samples available for MRD assessment. All 10 subjects (100%) evaluable at the  $10^{-5}$  sensitivity level were negative for MRD by next generation flow cytometry and/or next generation sequencing (NGS; Adaptive v 2.0). In the remaining 5 subjects, two subjects were indeterminate at  $10^{-5}$  due to insufficient cell counts but were MRD negative at the sensitivity threshold of  $10^{-4}$  by NGS. No clone identification could be performed in 3 subjects by NGS. Follow-up for response continues, however these preliminary data suggest compelling efficacy in this population of heavily pre-treated subjects.

***Other Neurotoxicities***

As of 06 March 2020, neurotoxicity with symptoms suggestive of Parkinsonian features that occurred after a period of recovery from CRS and/or ICANS have been observed in 6 subjects from Study 68284528MMY2001 (n=97). Among 6 subjects, 3 subjects had confirmed Parkinsonian-like symptoms with some shared features in their clinical presentation, with initial presentation between Day 19 and Day 101 after JNJ-68284528 infusion. Symptoms include altered mental status, feeling or appearing agitated, personality changes (such as being more withdrawn, expressing little emotion), lack of facial expression, being less communicative (answering questions with 1-word), difficulty with short term memory, slowing of movements, difficulty performing simple tasks (such as buttoning a shirt), joint or muscle stiffness or inflexibility, shuffling gait (walking very slowly with small steps and heavy feet), impaired balance, bradykinesia, slight cogwheeling/rigidity in extremities, micrographia (small handwriting) and impaired ability to perform activities of daily living (ADLs). On exam, all 3 subjects presented with symptoms described as Parkinsonian. The brain MRI showed no acute abnormalities in these subjects. The remaining 3 subjects had a mixed presentation of neurotoxicities (including some Parkinsonian-like symptoms), which presented with complicating comorbidities. All 6 subjects referenced above had previously experienced CRS (4 Grade 2, 2 Grade 3), all of which resolved and 5 out of 6 subjects had ICANS (3 Grade 1, 1 Grade 2, 1 Grade 3). Three subjects referenced above have died. Although these subjects also experienced other adverse events which may have contributed to the cause of death, such as infection and sepsis, the role of neurotoxicity cannot be ruled out.

**Study 68284528MMY2002**

Study 68284528MMY2002 is a Phase 2, open-label, multicenter study in China (sponsored by Legend Biotech HK Limited and Janssen R&D; Investigational New Drug holder is Legend Biotech) to evaluate the efficacy and safety of LCAR-B38M CAR-T cells in adult subjects with relapsed or refractory multiple myeloma. Approximately 60 subjects will be enrolled into the study. The primary objective is to evaluate the efficacy of LCAR-B38M CAR-T cells.

The first subject was dosed on 22 March 2019. As of 9 December 2019, 25 subjects were enrolled into the study and 13 subjects received LCAR-B38M CAR-T cells. Of the 12 subjects analyzed in the second safety evaluation team meeting, 12 subjects were reported with CRS, 1 subject had Grade 1, 5 had Grade 2, 5 had Grade 3 and 1 had Grade 4 CRS. The median onset of CRS after CAR-T infusion was 7 days (range: 2-8) and median duration of CRS was 5 days (range: 4-8). One subject received an infusion of LCAR-B38M CAR-T cells on 06 May 2019 and died on 12 May 2019 due to hemorrhage secondary to thrombocytopenia. The investigator attributed the thrombocytopenia, CRS, and acute renal failure to LCAR-B38M CAR-T cell therapy. The CRS events experienced by other subjects all resolved. No neurotoxicity has been reported as of the cut-off date. Further details are provided in the most recent edition of the Investigator's Brochure.

## 1.2. Overall Rationale for the Study

BCMA is a cell surface antigen highly expressed on cells of the B cell lineage. Comparative studies show a lack of BCMA in most normal tissues and absence of expression on CD34-positive hematopoietic stem cells (Carpenter 2013; Hsi 2008). This selective expression and the biological importance for the proliferation and survival of myeloma cells makes BCMA a promising target for CAR-T based immunotherapy, JNJ-68284528. Results in 74 subjects from the Legend-2 study (Section 1.1.5) indicate an ORR of 87.8% with a CR rate of 64.9%. The observed response rates and the reversible adverse events for most subjects, support further investigation of this approach in the current study.

The safety and efficacy of JNJ-68284528 in heavily pretreated patients with multiple myeloma is being evaluated in the Phase 1b/2 Study 68284528MMY2001. As the degree of benefit of the CAR-T therapy is dependent on a patient's immune response, JNJ-68284528 could provide potentially meaningful efficacy and durability when administered to patients with multiple myeloma who have had less exposure to immunomodulatory and cytotoxic therapy. The objective of Study 68284528MMY2003 is to determine the safety and efficacy of JNJ-68284528 in various clinical settings. Multiple cohorts will run in parallel with unique patient populations of unmet medical need enrolled. Approximately 40 subjects will be enrolled in Cohort A, and approximately 20 subjects each will be enrolled in all other cohorts. The subject population for each of the cohorts is described below.

- Cohort A: Subjects with progressive disease after 1 to 3 prior lines of therapy for multiple myeloma including a proteasome inhibitor (PI) and immunomodulatory drug (IMiD) either individually or in combination. Subjects are required to be refractory to lenalidomide. Subjects who have received prior therapy that is targeted to BCMA are excluded from this cohort.
- Cohort B: Subjects who have received one line of previous therapy containing a PI and an IMiD and who have early relapse defined as disease progression  $\leq 12$  months after an ASCT or  $\leq 12$  months after the start of front-line therapy for subjects who have not had an ASCT. Subjects who have received prior therapy that is targeted to BCMA are excluded from this cohort.
- Cohort C: Relapsed or refractory disease in subjects previously treated with a proteasome inhibitor (PI), immunomodulatory drug (IMiD), anti-CD38 monoclonal antibody and BCMA-directed therapy (excluding cellular immunotherapy).
- Cohort D (JNJ-68284528 plus lenalidomide): Subjects with multiple myeloma who did not achieve a complete response after 4 to 8 total cycles of initial therapy, including induction, high-dose therapy and ASCT with or without consolidation induction.
- Cohort E (daratumumab, bortezomib, lenalidomide, and dexamethasone [D-VRd] induction, then JNJ-68284528, followed by consolidation treatment with daratumumab and lenalidomide [D+R]): Subjects with high risk, newly diagnosed and untreated multiple myeloma (hr-NDMM) for whom hematopoietic stem cell transplant is not planned as initial therapy.

Subjects who meet the eligibility criteria for Cohort A and Cohort B, must be enrolled in Cohort B.



### 1.3. Potential Safety Risks and Mitigation Strategies

#### JNJ-68284528 (All Cohorts)

The potential risks of JNJ-68284528 are identified from the following: 1) results of nonclinical studies; 2) mechanism of action; and 3) previous clinical experience with JNJ-68284528 and LCAR-B38M CAR-T cells. Treatment of additional subjects and prolonged follow-up may reveal additional risks.

By stimulating an inflammatory cascade, there is potential for toxicity in other tissues or organs by non-specific immune cell activation. Therefore, special attention will be given to both immunological and immunogenicity-related toxicities. Safety risks and mitigation strategies are outlined in [Table 9](#).

**Table 9: Risks Associated with JNJ-68284528 and Mitigation Strategies**

Risk	Mitigation Strategies
Cytokine release syndrome (CRS) <sup>a, b</sup>	<p>Monitor closely for CRS and follow guidance for management in Section 6.2.1. Body temperature should be monitored twice daily for 28 days post infusion. At the first sign of CRS (such as fever) subjects should be immediately hospitalized for evaluation. See <a href="#">Table 10</a> for other hospitalization requirements. Rarely, severe CRS can evolve into a presentation consistent with hemophagocytic lymphohistiocytosis/macrophage activation syndrome (HLH/MAS) that may require additional therapy. The use of myeloid growth factors, particularly granulocyte colony-stimulating factor (G-CSF), should be avoided during CRS. Tocilizumab intervention may be considered with presenting symptom of fever in the absence of clear infectious etiology. Early tocilizumab should be considered in subjects at high risk of severe CRS. Section 6.2.1 provides management guidelines for CRS. Notify the sponsor if subject is experiencing Grade 2 or higher CRS.</p>
Neurologic toxicities <sup>a, b</sup>	<p><b>CAR-T cell-related neurotoxicity (ie, immune effector cell-associated neurotoxicity syndrome [ICANS])<sup>a</sup></b></p> <p>Early recognition of neurologic adverse events is critical to management. Subjects should be advised to seek medical evaluation if they notice new onset of headache, convulsions, speech disorders, visual disorders, disturbances in consciousness, confusion and disorientation, and coordination, balance disorders, or mental status changes. Notify the sponsor if subject is experiencing any grade ICANS.</p> <p>At the first sign of neurotoxicity, neurology consultation and evaluation should be considered. Hospitalization is required for Grade <math>\geq 2</math> CAR-T cell-related neurotoxicity (eg, ICANS). Section 6.2.2 provides management guidelines for neurotoxicity.</p> <p><b>Other Neurotoxicities</b></p> <p>Monitor closely for other neurotoxicities with clinical presentation for 1-year post-infusion with JNJ-68284528. Symptoms include difficulty sleeping, feeling or appearing agitated, personality changes (such as being more withdrawn, expressing little emotion), lack of facial expression, being less communicative (answering questions with 1-word answers), difficulty with short term memory, slowing of movements, difficulty performing simple tasks (such as buttoning a shirt), joint or muscle stiffness or inflexibility, shuffling gait (walking very slowly with small steps and heavy feet), and micrographia (small handwriting), and impaired ability to perform ADLs. If those neurologic or psychiatric symptoms are noted, contact the medical monitor and refer the</p>

**Table 9: Risks Associated with JNJ-68284528 and Mitigation Strategies**

Risk	Mitigation Strategies
	<p>subject immediately to a neurologist for a full evaluation. Section 6.2.2.2 provides further details for management guidelines for neurotoxicity. If a subject is non-responsive to interventions for this neurotoxicity, consideration should be given for therapies directed at reduction or elimination of CAR-T cells.</p> <p>Subjects with Parkinsonian-like neurotoxicities were observed at a higher frequency in subjects with high burden of disease and in subjects experiencing higher grade CRS (Grade 2 and above) and any grade ICANS. This may be indicative that <math>\geq</math> Grade 2 CRS or any grade ICANS are an early indication of risk of other neurotoxicity that may develop after a period of recovery from CRS and/or ICANS. Therefore, <math>\geq</math> Grade 2 CRS or any grade ICANS may represent an opportunity for early intervention and more aggressive supportive care (including steroids) especially in patients treated with high tumor burden, may mitigate the risk of developing neurotoxicity later, after resolution of CRS. Infection and sepsis were concurrently seen in many of these patients.</p>
Tumor lysis syndrome (TLS) <sup>a</sup>	Monitor closely for TLS with frequent monitoring of chemistry parameters and follow guidance for management in Section 6.2.3.
Second primary malignancies (SPMs) <sup>a</sup>	<p>Second primary malignancies may occur in subjects receiving JNJ-68284528. SPMs should be managed per institutional standards. Second primary malignancies must be reported during the duration of the study, irrespective of when they occur, and subsequently will be collected in a long-term follow-up study yearly until 15 years post dosing of JNJ-68284528. A tumor sample should be collected and DNA, RNA, or protein analysis may be performed to investigate the presence of lentiviral elements if an SPM develops. Section 6.2.4 provides management guidelines for SPMs.</p>
Cytopenias	<p>Frequent monitoring of hematological parameters and provide supportive care (eg, radiated blood and thrombocyte concentrates, granulocyte-colony stimulating factor for neutropenia) as outlined by institutional guidelines. Pegylated myeloid growth factors (ie, pegfilgrastim) are prohibited. Prolonged neutropenia may increase the risk of infection. Severe thrombocytopenia may increase the risk of bleeding.</p> <p>Initiating lenalidomide after JNJ-68284528 might cause significant neutropenia and thrombocytopenia. Monitor complete blood counts (CBC) weekly for first 2 cycles, biweekly for cycle 3 and every 28 days thereafter. A dose interruption and reduction may be required (refer to local prescription information and guidance in Table 12, Table 13, and Table 14). Subjects should be monitored frequently for infection and bleeding. Supportive care should be provided per institutional standards. Section 6.2.5 provides management guidelines for cytopenia. Parvovirus B19 monitoring by PCR should be considered in subjects experiencing prolonged neutropenia or a decline in neutrophil counts following recovery.</p>
Hypogammaglobulinemia	Monitor immunoglobulin levels after treatment and treat according to local guidelines, including administration of immunoglobulin replacement and monitoring for infection. Additional assessments of immunoglobulin levels may be done as per local standards of care. Section 6.2.6 provides management guidelines for hypogammaglobulinemia. Subjects with IgG < 400 mg/dL and recurrent infections may receive prophylactic IVIG as per institutional guidelines.
Infections	Do not administer JNJ-68284528 to patients with active infection. Frequent monitoring for the presence of infections, with cultures or implementation of empiric antibiotic therapy as appropriate, based on clinical judgment and institutional standards. Extended use of anti-microbial therapies for at least 6 month (or longer as per institutional guidelines) or consistent with post ASCT consensus guidelines after JNJ-68284528 dosing is recommended (See Attachment 18).

**Table 9: Risks Associated with JNJ-68284528 and Mitigation Strategies**

Risk	Mitigation Strategies
	<p>Perform screening for hepatitis B virus (HBV), hepatitis C virus (HCV), and human immunodeficiency virus (HIV) and monitor as clinically indicated, and initiate treatment as appropriate. Consider CMV serology at baseline, monitor with PCR testing as clinically indicated per institutional guidance.</p> <p>HBV reactivation, in some cases resulting in fulminant hepatitis, hepatic failure and death, may occur in subjects treated with drugs directed against B cells such as JNJ-68284528. HBV reactivation has occurred in subjects who appear to have resolved hepatitis B infection. Prophylaxis for herpes zoster reactivation is recommended during study treatment as clinically indicated. Routinely monitor HBV DNA and AST/ALT for subjects with risk of HBV reactivation (<a href="#">Attachment 10</a>)</p>
Hypersensitivity reactions	<p>Allergic reactions may occur with the infusion of JNJ-68284528. Serious hypersensitivity reactions including anaphylaxis, may be due to dimethyl sulfoxide (DMSO), dextran 40, or residual ampicillin or kanamycin in JNJ-68284528. Subjects should be treated urgently per institutional standards, avoiding corticosteroid use if possible. Subjects should receive premedication prior to JNJ-68284528 dosing as noted in Section 6.1.3.3.</p>

a Adverse event of special interest (see Section 12.3.3)

b Identified Risk

## 2. OBJECTIVES, ENDPOINTS, AND HYPOTHESIS

### 2.1. Objectives and Endpoints

Objectives	Endpoints
<b>Primary</b>	
<ul style="list-style-type: none"> <li>To evaluate the overall minimal residual disease (MRD) negative rate of subjects who receive JNJ-68284528</li> </ul>	<ul style="list-style-type: none"> <li>MRD negative rate at a <math>10^{-5}</math> threshold as defined by the International Myeloma Working Group (IMWG) criteria using next generation sequencing (NGS)</li> </ul>
<b>Secondary</b>	
<ul style="list-style-type: none"> <li>To evaluate the efficacy of JNJ-68284528</li> </ul>	<ul style="list-style-type: none"> <li>Overall Response Rate (ORR) (partial response [PR] or better) as defined by the IMWG response criteria</li> <li>VGPR, CR, sCR rate as defined by the IMWG response criteria, clinical benefit rate (CBR; CBR = ORR [sCR + CR + VGPR + PR] + minimal response [MR])</li> <li>Duration of response (DOR) and time to response (TTR)</li> </ul>
<ul style="list-style-type: none"> <li>To further characterize MRD negativity</li> </ul>	<ul style="list-style-type: none"> <li>MRD negative rate at 12 months for subjects who achieved a complete response (CR MRD neg 12 month)</li> <li>Time to MRD negativity, duration of MRD negativity,</li> </ul>

Objectives	Endpoints
	<ul style="list-style-type: none"> <li>MRD negative rate across clinical response groups (CR, stringent complete response [sCR], very good partial response [VGPR])</li> </ul>
<ul style="list-style-type: none"> <li>To characterize the safety of JNJ-68284528</li> </ul>	<ul style="list-style-type: none"> <li>Incidence and severity of adverse events, laboratory results, and other safety parameters</li> </ul>
<ul style="list-style-type: none"> <li>To characterize the pharmacokinetics and pharmacodynamics of JNJ-68284528</li> </ul>	<ul style="list-style-type: none"> <li>Pharmacokinetic and pharmacodynamic markers including but not limited to depletion of soluble BCMA and BCMA expressing cells, systemic inflammatory cytokine concentrations and immune related proteins, and markers of CAR-T cell activation, expansion (proliferation), and persistence via monitoring CAR-T positive cell counts and CAR transgene level.</li> </ul>
<ul style="list-style-type: none"> <li>To assess the immunogenicity of JNJ-68284528</li> </ul>	<ul style="list-style-type: none"> <li>Presence of anti- JNJ-68284528 antibodies</li> </ul>
<b>Exploratory</b>	
<ul style="list-style-type: none"> <li>To further characterize the efficacy of JNJ-68284528</li> </ul>	<ul style="list-style-type: none"> <li>Progression-free survival (PFS), overall survival (OS)</li> <li>Imaging plus MRD negative rate (if positron emission tomography [PET] is locally available)</li> </ul>
<ul style="list-style-type: none"> <li>To explore changes in patient-reported outcomes (PRO) after treatment with JNJ-68284528</li> </ul>	<ul style="list-style-type: none"> <li>Time to worsening of symptoms using the MySim-Q total symptom score</li> <li>Change from baseline in HRQoL (symptoms, functioning, and well-being) using the European Organization for Research and Treatment of Cancer (EORTC) QLQ-C30, MySim-Q (Multiple Myeloma Symptom and Impact Questionnaire), Patient Global Impression of Change (PGIC), and Patient Global Impression of Severity (PGIS).</li> </ul>
<ul style="list-style-type: none"> <li>To assess symptomatic adverse events from the patient perspective via patient reported outcomes of the Common Terminology Criteria for Adverse Events (PRO-CTCAE).</li> </ul>	<ul style="list-style-type: none"> <li>PRO-CTCAE report using validated questions.</li> </ul>
<ul style="list-style-type: none"> <li>To characterize the impact of JNJ-68284528 CAR-T process on medical resource utilization</li> </ul>	<ul style="list-style-type: none"> <li>Number of subjects with type and length of inpatient stay and overall medical encounters</li> </ul>
<ul style="list-style-type: none"> <li>To characterize potential early clinical, translational, and imaging markers for neurotoxicity (predictive markers)</li> </ul>	<ul style="list-style-type: none"> <li>Qualitative changes in handwriting assessment</li> <li><math>T_{max}</math>, <math>C_{max}</math>, and phenotypic analysis of CAR-T cells</li> <li>Neuroimaging (CT/MRI/PET)</li> </ul>

## 2.2. Hypotheses

The primary hypothesis is that JNJ-68284528 will induce a deep-response, measured by MRD negative rate in the clinical settings investigated.

## 3. STUDY DESIGN AND RATIONALE

### 3.1. Overview of Study Design

This is a Phase 2, multicohort, open-label, multicenter study to determine whether treatment with JNJ-68284528 results in MRD negativity in adult subjects with multiple myeloma in various clinical settings. Multiple cohorts will run in parallel. Approximately 40 subjects will be enrolled in Cohort A, and approximately 20 subjects each will be enrolled in all other cohorts. The primary endpoint for all cohorts will be overall MRD negative rate.

Subjects will participate in one of the following cohorts based on eligibility criteria defined in Section 4.1 to Section 4.5, and summarized below:

- Cohort A will include subjects with progressive disease after 1 to 3 prior lines of therapy for multiple myeloma including a proteasome inhibitor (PI) and immunomodulatory therapy (IMiD) either individually or in combination. In addition, all study subjects will be lenalidomide refractory. Subjects who have received prior therapy that is targeted to BCMA are excluded from this cohort.
- Cohort B will include subjects who have had one line of previous therapy containing a PI and an IMiD and who have early relapse defined as having disease progression  $\leq 12$  months after an ASCT or  $\leq 12$  months after the start of front-line therapy for subjects who have not had an ASCT. Subjects who have received prior therapy that is targeted to BCMA are excluded from this cohort.
- Cohort C will include subjects with relapsed or refractory disease in subjects previously treated with a PI, IMiD, anti-CD38 monoclonal antibody and BCMA-directed therapy (excluding cellular immunotherapy).
- Cohort D (JNJ-68284528 plus lenalidomide) will include subjects with multiple myeloma who did not achieve a complete response after 4 to 8 total cycles of initial therapy, including induction, high-dose chemotherapy and ASCT with or without consolidation.
- Cohort E (D-VRd induction, JNJ-68284528, then D+R consolidation) will include subjects with high risk newly diagnosed and untreated multiple myeloma (hr-NDMM) for whom hematopoietic stem cell transplant is not planned as initial therapy.

Subjects who meet the eligibility criteria for Cohort A and Cohort B, must be enrolled in Cohort B.

Assessment of MRD negativity and all response assessments will be performed by central laboratory. Disease status will be evaluated according to the IMWG consensus recommendations for multiple myeloma treatment response criteria ([Attachment 1](#)). Response will be determined using a validated computer algorithm.

Safety evaluations will include a review of adverse events, laboratory test results, vital sign measurements, physical examination findings (including neurologic examination), assessment of cardiac function, Immune Effector Cell-associated Encephalopathy (ICE) score, handwriting assessment, and assessment of Eastern Cooperative Oncology Group (ECOG) performance status grade. Follow up of subjects for disease progression and survival will continue during the Post-treatment Phase. All study evaluations will be conducted according to the Time and Events Schedules (Table 1 to Table 8). At the investigator discretion and with sponsor approval, study visits (for all cohorts) in the post-treatment part of the study, as early as after Day 100 after JNJ-68284528 infusion, may be performed remotely via telemedicine technology that connects study subjects to their research coordinators and investigators, if not using the sponsor's telemedicine solution. Blood sample collection may also be performed at the subject's home by mobile study personnel (ie, nurses and mobile phlebotomist), if not using the sponsor's home health nursing solution in the post-treatment period, as early as after Day 100 after JNJ-68284528 infusion.

The sponsor will establish a data cutoff date for the clinical study report (CSR) analyses. The primary analysis for each cohort will be conducted approximately 1 year after the last subject in each cohort receives their initial dose of JNJ-68284528. Subjects will be followed for survival after the clinical cutoff for the primary CSR. For Cohort A, B, and C, the cohort completion will be defined as no later than 2 years after the last subject in each cohort has received their initial dose of JNJ-68284528. For Cohort D and Cohort E, cohort completion will be defined as 2 ½ years after the last subject receives their initial dose of JNJ-68284528.

### **Data Monitoring Committee**

A Data Monitoring Committee (DMC) will periodically review all safety and other relevant data collected in this study as well as data obtained in other JNJ-68284528 protocols. The DMC will provide recommendations to ensure subject safety and mitigate risk for all study subjects. Stopping rules based on safety signals will also be provided and maintained by the DMC. The roles and responsibilities of the DMC are to be specified in the DMC charter.

#### **3.1.1. Cohort A Study Design**

Subjects enrolled in Cohort A will undergo apheresis after screening to acquire peripheral blood mononuclear cells (PBMCs). For all subjects, JNJ-68284528 will be generated from the subject's T cells selected from the apheresis product. Subjects for whom apheresis or manufacturing fails will be allowed a second attempt at apheresis.


Bridging therapy (anti-plasma cell directed treatment between apheresis and lymphodepletion) will be allowed when clinically indicated (ie, to maintain a subject's clinical status while waiting for manufacturing of JNJ-68284528), with the permission of the sponsor. Bridging therapy should be a short-term treatment (approximately 1 to 2 cycles) to decrease tumor burden and may include therapies to which a subject has not been previously exposed. The sponsor will not permit subjects who are found to be in a confirmed CR after bridging therapy to receive JNJ-68284528. Subjects in Cohort A who do not receive an infusion of JNJ-68284528 will be replaced.

After JNJ-68284528 production and product release, all subjects will receive a conditioning regimen of IV cyclophosphamide 300 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> daily for 3 days. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an estimated glomerular filtration rate (eGFR) of 30 to 70 mL/min/1.73m<sup>2</sup>. JNJ-68284528 at the recommended phase 2 dose (RP2D) based on the MMY2001 study (0.75 x 10<sup>6</sup> CAR-positive viable T cells/kg; see Section 3.2) will be administered 5 to 7 days after the start of the conditioning regimen. The conditioning regimen will lead to lymphodepletion and help promote CAR-T cell expansion in the subject. Subjects who receive an infusion of JNJ-68284528 should continue all subsequent post-infusion assessments.

Of the approximately 40 subjects in Cohort A, approximately 20 will receive cilta-cel manufactured using the commercial process.

A diagram of the study design for Cohort A is provided in [Figure 2](#).

**Figure 2: Schematic Overview of the Study, Cohort A**






### 3.1.2. Cohort B Study Design

Subjects enrolled in Cohort B will undergo apheresis after screening to acquire peripheral blood mononuclear cells (PBMC). For all subjects JNJ-68284528 will be generated from the subject's T cells selected from the apheresis product. Subjects for whom apheresis or manufacturing fails will be allowed a second attempt at apheresis. Bridging therapy (anti-plasma cell directed treatment between apheresis and lymphodepletion) will be allowed when clinically indicated (ie, to maintain a subject's clinical status while waiting for manufacturing of JNJ-68284528), with the permission of the sponsor. Bridging therapy should be a short-term treatment (approximately 1 to 2 cycles) to decrease tumor burden and may include therapies to which a subject has not been previously exposed. The sponsor will not permit subjects who are found to be in a confirmed CR after bridging therapy to receive JNJ-68284528. Subjects in Cohort B who do not receive an infusion of JNJ-68284528 will be replaced.

After JNJ-68284528 production and product release, all subjects will receive a conditioning regimen of IV cyclophosphamide 300 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> daily for 3 days. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an eGFR of 30 to 70 mL/min/1.73m<sup>2</sup>. JNJ-68284528 at the RP2D based on the MMY2001 study (0.75 x 10<sup>6</sup> CAR-positive viable T cells/kg) will be administered 5 to 7 days after the start of the conditioning regimen. The conditioning regimen will lead to lymphodepletion and help promote CAR-T cell expansion in the subject. Subjects who receive an infusion of JNJ-68284528 should continue all subsequent post-infusion assessments. Subjects in Cohort B who do not receive an infusion of JNJ-68284528 will be replaced.

A diagram of the study design for Cohort B is provided in [Figure 3](#).

**Figure 3: Schematic Overview of the Study, Cohort B**

### 3.1.3. Cohort C Study Design


Subjects enrolled in Cohort C will undergo apheresis after screening to acquire peripheral blood mononuclear cells (PBMC). For all subjects, JNJ-68284528 will be generated from the subject's T cells selected from the apheresis product. Subjects for whom apheresis or manufacturing fails will be allowed a second attempt at apheresis.

Bridging therapy (anti-plasma cell directed treatment between apheresis and lymphodepletion) will be allowed when clinically indicated (ie, to maintain a subject's clinical status while waiting for manufacturing of JNJ-68284528), with the permission of the sponsor. Bridging therapy should be a short-term treatment (approximately 1 to 2 cycles) to decrease tumor burden and may include therapies to which a subject has not been previously exposed. The sponsor will not permit subjects who are found to be in CR after bridging therapy to receive JNJ-68284528. Subjects in Cohort C who do not receive an infusion of JNJ-68284528 will be replaced.

After JNJ-68284528 production and product release, all subjects will receive a conditioning regimen of IV cyclophosphamide 300 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> daily for 3 days. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an eGFR of 30 to 70 mL/min/1.73m<sup>2</sup>. JNJ-68284528 at the recommended phase 2 dose (RP2D) of 0.75 x 10<sup>6</sup> CAR-positive viable T cells/kg based on the 68284528MMY2001 study will be administered 5 to 7 days after the start of the conditioning regimen. The conditioning regimen will lead to lymphodepletion and help promote CAR-T cell expansion in the subject. Subjects who receive an infusion of JNJ-68284528 should continue all subsequent post-infusion assessments.

A diagram of the study design for Cohort C is provided in [Figure 4](#).

Figure 4: Schematic Overview of the Study, Cohort C



### 3.1.4. Cohort D Study Design

Subjects enrolled in Cohort D will undergo apheresis after screening to acquire peripheral blood mononuclear cells (PBMC). For all subjects, JNJ-68284528 will be generated from the subject's T cells selected from the apheresis product. Subjects for whom apheresis or manufacturing fails will be allowed a second attempt at apheresis. After apheresis and prior to administration of cyclophosphamide and fludarabine (conditioning regimen prior to JNJ-68284528 infusion) subjects will receive 1 or more cycles of lenalidomide at a dose of 10 mg per day upon adequate hematologic recovery from ASCT (absolute neutrophil count [ANC]  $\geq 1 \times 10^9/L$  and platelet count  $\geq 75 \times 10^9/L$ ) when minimum laboratory requirements are met. Alternative bridging therapy instead of, or in addition to, lenalidomide is permissible with sponsor approval. Additionally, if an alternative bridging therapy is used, assessments to address lenalidomide toxicity will not be required in [Table 3](#) (footnote v). The purpose of bridging therapy is to reduce the myeloma burden prior to lymphodepletion chemotherapy and JNJ-68284528 administration.

After JNJ-68284528 production and product release, all subjects will receive a conditioning regimen of IV cyclophosphamide 300 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> daily for 3 days. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an eGFR of 30 to 70 mL/min/1.73m<sup>2</sup>. JNJ-68284528 at the RP2D based on the MMY2001 study ( $0.75 \times 10^6$  CAR-positive viable T cells/kg) will be administered 5 to 7 days after the start of the conditioning regimen. The conditioning regimen will lead to lymphodepletion and help promote CAR-T cell expansion in the subject.

A strategy of staggered dosing with JNJ-68284528 will be applied to Cohort D. There must be an observation period of at least 4 weeks between administration of JNJ-68284528 to the first 5 subjects to allow for subject 28-day safety review prior to next subject dosing. In addition, the first 5 subjects will not receive lenalidomide after JNJ-68284528 therapy to allow time for the Data Monitoring Committee (described in [Section 3.1](#)) to review safety data. After the first 5 subjects are dosed with JNJ-68284528, the Data Monitoring Committee (DMC) will convene to review safety and any other relevant data.

Based on recommendation of the DMC, subsequent subjects in this cohort will be eligible to receive lenalidomide after JNJ-68284528. The same strategy of staggered dosing will also be applied to the first 5 subjects receiving lenalidomide after JNJ-68284528 (ie, the 6<sup>th</sup> through 10<sup>th</sup> subject enrolled in Cohort D). There must be an observation period of at least 4 weeks between administration of JNJ-68284528 (followed by lenalidomide) to these 5 subjects. The DMC will review safety and any other relevant data from the first 5 subjects who receive JNJ-68284528 followed by lenalidomide before a decision is made regarding the treatment plan for further subjects enrolled in Cohort D.

JNJ-68284528 plus lenalidomide treatment: Subjects will initiate lenalidomide maintenance therapy at a minimum of 21 days post JNJ-68284528 infusion and after resolution of any cytokine release syndrome (CRS) or neurologic toxicities per ASTCT criteria. Subjects will continue to receive lenalidomide until confirmed PD, unacceptable toxicity, or for 2 years post JNJ-68284528

infusion, whichever occurs first. The initial dose of lenalidomide post-JNJ-68284528 infusion will depend on the level of hematologic recovery.


**Lenalidomide after JNJ-68284528**

A minimum of 21 days after infusion of JNJ-68284528, after resolution of CRS or neurologic toxicities associated with JNJ-68284528, subjects in the cohort will receive lenalidomide treatment. The initial dose of lenalidomide will depend on the level of hematologic recovery as described in Section [6.1.4](#).

Subjects should continue all subsequent post-infusion assessments after infusion of JNJ-68284528. Subjects will receive lenalidomide until confirmed PD, unacceptable toxicity, or for 2 years post JNJ-68284528 infusion, whichever occurs first. Subjects in Cohort D who do not receive an infusion of JNJ-68284528 will be replaced.

A diagram of the study design for Cohort D is provided in [Figure 5](#).

**Figure 5: Schematic Overview of the Study, Cohort D**



a Additional cycles or alternative therapy permitted with sponsor approval

b See Section 3.1.4 for staggered dosing strategy

### 3.1.5. Cohort E Study Design

Subjects enrolled in Cohort E will receive 4 cycles of quadruplet induction regimen of daratumumab, bortezomib, lenalidomide and dexamethasone (D-VRd) as tolerated. All subjects in Cohort E will undergo apheresis after cycle 1 or 2 of the D-VRd induction regimen. At the investigator's discretion, subjects may undergo peripheral stem cell mobilization and harvesting after Cycle 3 or 4 of induction treatment (D-VRd) to allow for salvage ASCT after disease progression is confirmed.

For all subjects JNJ-68284528 will be generated from the subject's T cells selected from the apheresis product. Subjects for whom apheresis or manufacturing fails will be allowed a second attempt at apheresis.

After JNJ-68284528 production and product release, all subjects will receive a conditioning regimen of IV cyclophosphamide 300 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> daily for 3 days. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an eGFR of 30 to 70 mL/min/1.73m<sup>2</sup>. JNJ-68284528 at the recommended phase 2 dose (RP2D) of 0.75 x 10<sup>6</sup> CAR-positive viable T cells/kg based on the 68284528MMY2001 study will be administered 5 to 7 days after the start of the conditioning regimen. The conditioning regimen will lead to lymphodepletion and help promote CAR-T cell expansion in the subject. Subjects who receive an infusion of JNJ-68284528 should continue all subsequent post-infusion assessments.

After the infusion of JNJ-68284528, subjects will receive consolidation treatment with D+R until confirmed PD, unacceptable toxicity, or for 2 years post JNJ-68284528 infusion, whichever occurs first. Lenalidomide treatment will be initiated a minimum of 21 days after infusion of JNJ-68284528, and will only start after resolution of any grade CRS or neurologic toxicities associated with JNJ-68284528 per ASTCT criteria. The initial dose of lenalidomide will depend on the level of hematologic recovery as described in Section 6.1.4. Subjects will initiate daratumumab approximately 60-100 days after infusion of JNJ-68284528, upon adequate recovery from ongoing toxicities as described in Section 6.1.5.3. The start of daratumumab treatment will align with Day 1 of the next lenalidomide cycle. At least 1 cycle of lenalidomide must be completed prior to initiating daratumumab.


Subjects should continue all subsequent post-infusion assessments after infusion of JNJ-68284528. Subjects in Cohort E who do not receive an infusion of JNJ-68284528 will be replaced.

A strategy of staggered dosing will be applied to Cohort E. There will be a period of at least 2 weeks between administration of JNJ-68284528 to the first 5 subjects with periodic reviews by the DMC. After being treated with JNJ-68284528, the safety data of the first 5 subjects will be reviewed by the DMC to determine if there are any potential pharmacodynamic interactions between D-VRd induction treatment and JNJ-68284528. In addition, in order to evaluate the safety of adding D+R consolidation treatment to JNJ-68284528 the DMC will also continue to review the safety data of these first 5 subjects after the start of consolidation therapy in a periodic basis. The composition of the committee and procedures will be provided in the DMC charter. The DMC will continue to evaluate safety throughout the duration of the cohort per the DMC charter.



A diagram of the study design for Cohort E is provided in [Figure 6](#).

**Figure 6: Schematic Overview of the Study, Cohort E**



<sup>a</sup> Eligible subjects will receive 4 cycles of D-VRd induction therapy as tolerated

<sup>b</sup> Subjects in Cohort E may have peripheral stem cell mobilization and harvesting after Cycle 3 or 4 of induction therapy, at the investigator's discretion.

### 3.1.6. Retreatment with JNJ-68284528

Subjects in Cohorts A, B, and C may be considered for retreatment with JNJ-68284528. Subjects in Cohort D and Cohort E will not be allowed to receive retreatment.

Subjects must satisfy the following criteria to be eligible for retreatment, with approval from the sponsor:

- Progressive disease (PD) after best response of minimal response (MR) or better
- No ongoing Grade 2 non-hematologic toxicity (except for nausea, vomiting, hair loss). No ongoing Grade 3 or higher hematologic toxicity
- At least 6 months between first JNJ-68284528 infusion and detection of PD

Subjects must satisfy all inclusion and exclusion criteria (Section 4), except for exclusion criteria 1 (Cohorts A, B, C) and 2 (Cohorts A, B), to be eligible for retreatment. A maximum of 1 retreatment may occur per subject. Bridging therapy is allowed for subjects receiving retreatment with permission from the sponsor.

Assessments for subjects receiving retreatment should follow the Time and Events schedules (Table 1 and Table 2 for Cohorts A, B, and C) with the following exceptions:

- Subjects do not have to sign an additional informed consent form
- Subject height does not need to be collected again
- Assessments scheduled prior to apheresis will only occur if it is necessary to collect a second apheresis sample.
- PRO assessments will not be collected upon retreatment

Subjects who received retreatment with JNJ-68284528 and are in follow-up at the end of the study (2 years after the last subject in each cohort receives the initial dose of JNJ-68284528) will be monitored in the long-term follow-up study for 15 years from the time of last treatment (see Section 9.1.6.5). For subjects who receive retreatment, MRD negativity after retreatment with JNJ-68284528 will not be counted in the primary analysis for efficacy. The subject's best response before retreatment will be captured in the efficacy analysis.

### 3.2. Rationale of Dose and Administration Schedule Selection

The conditioning regimen of cyclophosphamide 300 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> daily for 3 doses will lead to lymphodepletion and help promote CAR-T cell expansion in the subject. Cyclophosphamide 300 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> before JNJ-68284528 infusion (Day 1) is consistent with the lymphodepletion regimen used in the marketed CAR-T products. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an eGFR of 30 to 70 mL/min/1.73m<sup>2</sup>.

JNJ-68284528 will be administered at a targeted infused dose of 0.75 x 10<sup>6</sup> CAR-positive viable T cells/kg.

As of the 24 June 2019 data cutoff, 25 subjects had received an infusion of JNJ-68284528 in Study 68284528MMY2001. Twenty-one subjects had at least 1 post-baseline disease evaluation. An overall response rate of 90.5% was observed, with 4 stringent complete response (sCR), 2 CRs, 7 very good partial responses (VGPR), and 6 partial responses (PR). The responses occurred rapidly, with most subjects exhibiting  $\geq 50\%$  myeloma protein reduction at the first disease assessment (scheduled at Day 28 after JNJ-68284528 dosing). The responses appear to be continuously deepening for all subjects who are in follow-up and no disease progression has been observed as of data cutoff.

The safety profile was consistent with observations from the Legend-2 study (Section 1.1.5). Data from these 25 subjects were reviewed by the Safety Evaluation Team (SET), which recommended maintaining the dose of  $0.75 \times 10^6$  CAR-positive viable T cells/kg for the Phase 2 portion of Study 68284528MMY2001.

Together, preliminary efficacy and safety data from Study 68284528MMY2001 showed that a dose of  $0.75 \times 10^6$  cells/kg of JNJ-68284528 CAR-T cell is highly efficacious in a patient population with no alternative treatment options.

### 3.3. Study Design Rationale

#### Rationale for Study Cohorts

The safety and efficacy of JNJ-68284528 in heavily pretreated patients with multiple myeloma is being evaluated in the Phase 1b/2 Study 68284528MMY2001. The degree of benefit of CAR-T therapy may be dependent on the fitness of the patient's immune system at the time of T-cell apheresis (Fraietta 2018). This study includes subjects in varying stages of multiple myeloma treatment in order to assess the effect of prior exposure to immunomodulatory and cytotoxic therapy on the efficacy of JNJ-68284528:

- Cohort A includes subjects with progressive disease after 1 to 3 prior lines of therapy for multiple myeloma including a proteasome inhibitor (PI) and immunomodulatory therapy (IMiD) either individually or in combination. Subjects will be refractory to lenalidomide. There are many approved triplet regimens for patients with multiple myeloma that have relapsed after 1 to 3 prior lines of therapy. However, these regimens have largely been tested in lenalidomide naïve or lenalidomide non-refractory patients. Pivotal Phase 3 studies (ASPIRE, ELOQUENT-2, Tourmaline-MM1, POLLUX, etc.) excluded lenalidomide refractory patients because these studies randomized against lenalidomide plus dexamethasone control arms. Given that lenalidomide is now frequently administered in front-line maintenance, and relapse settings in both the US and EU, there are few options for patients with lenalidomide-refractory disease, resulting in a high unmet need for these patients. In fact, patients who are lenalidomide refractory have a poor median PFS ranging from 8.6 and 9.5 months when treated with bortezomib, pomalidomide and dexamethasone (Richardson 2019) or carfilzomib and dexamethasone (Moreau 2017). Thus, novel therapies such as BCMA directed CAR-T cells are a promising treatment alternative.

- Cohort B includes subjects who have had one line of previous therapy containing an IMiD and a PI and who have early relapse defined as having disease progression <12 months after an ASCT or <12 months after the start of front-line therapy for subjects who have not had an ASCT. Patients that have an early relapse after standard front-line therapies have a particularly poor prognosis and represent an area of unmet need. This group represents approximately 15% of all patients that receive their first line of therapy. Three centers have reported that patients who relapse within 1 year from start of initial therapy (with PI or IMiD-based induction) have a median OS of 22 months and patients that relapse within 1 year of ASCT have a median OS of approximately 20 months, which is substantially shorter than expected outcomes in multiple myeloma today (median OS not reached [NR]) for both groups of patients, HR 13.7) ([Majithia 2016](#); [Ong 2016](#); [Jimenez-Zepeda 2015](#)). New treatments, such as BCMA directed CAR-T cells may offer an alternative to currently available treatments which have failed to address this unmet medical need.
- Cohort C includes subjects with relapsed or refractory disease who have been previously treated with a PI, IMiD, anti-CD38 monoclonal antibody and BCMA-directed therapy (excluding cellular immunotherapy). In spite of recent advancement in the treatment of multiple myeloma, most patients still relapse or become refractory to the last line of therapy and new approaches are needed. BCMA is considered a promising target for anti-myeloma therapy, and currently multiple BCMA-directed platforms (eg, antibody-drug conjugate [ADC], bispecific T-cell engager [BiTE], CAR-T) are in the investigational phase ([Cho 2018a](#); [Cho 2018b](#)). Recent data from belantamab mafodotin (GSK'916) ([Trudel 2019](#)) and AMG 420 ([Topp 2019](#)) have shown that patients still relapse after receiving these therapies and are left with limited options. Additionally, [Gore et al \(2018\)](#) reported that patients with ALL who relapsed after receiving blinatumomab (BiTE therapy), had an objective response to CAR-T therapy for the same target. The majority of other clinical trials for multiple myeloma patients receiving CAR-T have excluded patients who have previously received BCMA-directed therapy. There is an unmet need to address the sequence of treatments for BCMA-directed agents in multiple myeloma. Therefore, this cohort will determine whether BCMA CAR-T therapy remains effective after exposure to other BCMA-directed agents.
- Cohort D includes subjects with recently diagnosed multiple myeloma who did not achieve a CR after ASCT. This patient population does not achieve an optimal response to standard first line therapy including ASCT and is anticipated to have poorer clinical outcomes on standard lenalidomide maintenance treatment. Experience with JNJ-68284528 in Study 68284528MMY2001 and LCAR-B38M CAR-T cells in the Legend-2 study, demonstrates substantial response rates, MRD negative rates, and acceptable toxicity in a heavily pre-treated population of subjects. As the degree of benefit of CAR-T therapy is dependent on a patient's immune response, JNJ-68284528 could have potential meaningful efficacy and durability when administered to patients with multiple myeloma who have had less exposure to immunomodulatory and cytotoxic therapy. Therefore, addition of JNJ-68284528 to lenalidomide treatment could provide long-term benefit in this patient population compared to standard of care lenalidomide maintenance. In addition, lenalidomide has been shown to promote immunologic memory in T-cell populations. In a pre-clinical study, infusion of CS1 CAR-T followed by daily low dose of lenalidomide in myeloma tumor bearing mice preferentially expanded CD8<sup>+</sup>CAR-T cells. Lenalidomide treatment also exerted a costimulatory effect on T-cell response by increasing production of IL-2 and IFN- $\gamma$  and inhibiting production of anti-inflammatory cytokines; resulting in enhanced anti-myeloma

activity and persistence of CAR-T cells in vivo (Wang 2018). Similar effects have been observed in CD19 CAR-T and EGFRvIII CAR-T in-vitro and in-vivo studies (Kuramitsu 2015; Otahal 2016). The combination of anti-BCMA CAR-T cells and lenalidomide treatment is currently under study in patients with multiple myeloma (NCT03070327).

- Cohort E includes subjects with hr-NDMM for whom hematopoietic stem cell transplant is not planned as initial therapy. The treatment of high risk newly diagnosed patients with multiple myeloma is evolving. Induction therapy followed by transplant is not feasible in all patients due to age or patient frailty. Despite advances in treatment, including quadruplet therapies, patients with hr-NDMM still respond more poorly than patients with standard risk myeloma (Kurki 2016; Voorhees 2020). Therefore, novel strategies are needed.

Due to the effectiveness of triplet and quadruplet therapies, some patients are deferring initial transplant. The most active combination to date is the VRd regimen. The superiority of the VRd regimen has been established by the results of the Phase 3 SWOG S0777 study (Durie 2017). This study demonstrated both increased PFS and OS in the VRd arm compared with Rd alone. Further, the Phase 3 study described by Attal and colleagues (2017) showed that a deep level of response, CR and MRD negative status can be achieved with the VRd regimen without transplant.

The sponsor has observed compelling clinical data with daratumumab in combination with either bortezomib or lenalidomide in the relapsed/refractory setting, and bortezomib in the frontline setting. The addition of daratumumab with VRd is anticipated to improve the response rates and the depth of response and may lead to improved long-term outcomes in newly diagnosed patients with multiple myeloma. The Phase 2 Study 54767414MMY2004 (NCT02874742), compared daratumumab with VRd (D-VRd) against VRd in patients with newly diagnosed multiple myeloma that were transplant eligible. The study found 42.4% of patients treated with D-VRd achieved a sCR, compared to 32.0% of patients who received VRd alone by the end of consolidation (Voorhees 2019). Overall, the safety profile of D-VRd was consistent with the safety profile for each therapy separately.

The Phase 3 study of D-VRd (54767414MMY3014) in the same patient population described above utilizes the subcutaneous (SC) formulation of daratumumab instead of the IV formulation utilized in the Phase 2 VRd study 54767414MMY2004. This is expected to provide similar exposure and limit additional toxicity for patients treated with this quadruplet regimen, from infusion-related reactions due to daratumumab. The initial PK of SC daratumumab administration shows similar exposure to IV and improved safety with a lower IRR rate compared with the IV formulation (Usmani 2019). Additionally, another Phase 3 study of D-VRd is currently being conducted in newly diagnosed multiple myeloma patients in whom transplant is not intended, 54767414MMY3019 (NCT03652064).

Despite prolonging the PFS and OS in patients with newly diagnosed multiple myeloma who are transplant ineligible, the disease remains incurable, especially for patients with high risk disease. Experience with JNJ-68284528 in Study 68284528MMY2001 and LCAR-B38M CAR-T cells in the Legend-2 study, demonstrates substantial response rates, MRD negative rates, and acceptable toxicity in a heavily pre-treated population of subjects. As the degree of benefit of CAR-T therapy is dependent on a patient's immune response, JNJ-68284528 could have potential meaningful efficacy and durability when administered in the frontline setting

to patients with multiple myeloma who have had less exposure to immunomodulatory and cytotoxic therapy. In order to maximize disease control, daratumumab and lenalidomide will be given as consolidation therapy. Daratumumab, lenalidomide, and dexamethasone has been proven to be efficacious in transplant ineligible patients in the Phase 3 study 54767414MMY3008 (NCT02252172). The use of lenalidomide to potentiate the efficacy of JNJ-68284528 is addressed in the design rationale of Cohort D. The immunomodulatory effects of daratumumab may reduce the immunosuppressive effects of the microenvironment, and thereby improve the efficacy of JNJ-68284528 CAR T-cells. Multiple myeloma induces bone marrow microenvironment changes which may lead to decreased efficacy of T-cell mediated cellular therapy (de Haart 2013). Regulatory T cells (T<sub>regs</sub>), regulatory B cells (B<sub>regs</sub>), and myeloid-derived suppressor cells (MDSCs) may impair the efficacy of the infused CAR T-cells, similar to their suppression of non-engineered T-cells. Studies have shown that the CD38-targeting antibody daratumumab has strong immunomodulatory effects including elimination of CD38-positive Tregs, CD38-positive B-regs, and CD38-positive MDSCs (Krejci 2016; van de Donk 2018; Adams 2019; van de Donk 2017; Adams 2018). This results in expansion of T-cells and a better cytotoxic capacity of daratumumab-exposed T cells (higher granzyme B content).


### **Rationale for MRD as Primary Endpoint**

Achievement of MRD negativity by flow or next generation sequencing (NGS) at a cutoff of both  $10^{-4}$  and  $10^{-5}$  is associated with improvement of both PFS and OS (Lahuerta 2008; Landgren 2016; Munshi 2016). Meta- and pooled analyses of MRD data have demonstrated that MRD negative status is the strongest independent prognostic biomarker for PFS and OS in newly diagnosed multiple myeloma. Based upon the strength of these early analyses, the acceptance of MRD as a validated, clinical endpoint may be achieved in the near future.

Furthermore, data from 2 studies utilizing daratumumab in combination with either Rd (54767414MMY3003 [NCT02076009]) or Vd (54767414MMY3004 [NCT02136134]) in subjects with relapsed/refractory multiple myeloma have shown over a 3-fold increase in the number of subjects who achieved MRD negative status (Figure 7) (Avet 2016). Subjects achieving MRD negative status demonstrated improved PFS (Figure 7). These studies utilized the clonoSEQ MRD assay (Adaptive Biotechnology) that is an analytically validated NGS assay. The use of the clonoSEQ assay allows for centralized analysis and the consistent, accurate evaluation of MRD status that will support the primary endpoint.



**Figure 7: Progression-free Survival According to MRD Status  $10^{-5}$  in Studies 54767414MMY3003 and 54767414MMY3004**



In this study, minimal residual disease will be evaluated at suspected CR when a subject has achieved a deep clinical response. Additional landmarked bone marrow aspirate samples will be obtained to allow for the statistical analysis of the association of MRD with PFS/OS and to evaluate the durability of MRD negativity in these subjects.

The primary endpoint will evaluate the overall MRD negative rate. Secondary endpoints will assess the durability of MRD negativity, thus assessing whether subjects who obtain MRD negativity are able to maintain that depth of response. The sponsor will assess the impact of this durability of MRD negativity on the long-term outcome of PFS.

**4. SUBJECT POPULATION**

The inclusion and exclusion criteria for enrolling subjects in this study are described in the following subsections. If there is a question about the inclusion or exclusion criteria below, the investigator must consult with the appropriate sponsor representative and resolve any issues before enrolling a subject in the study. Waivers are not allowed.

Eligibility criteria are grouped by cohort. Inclusion and exclusion criteria for Cohort A are described in Section 4.1, Cohort B in Section 4.2, Cohort C in Section 4.3, Cohort D in Section 4.4, and Cohort E in Section 4.5.

Subjects who meet the eligibility criteria for both Cohort A and Cohort B, must be enrolled into Cohort B.

## 4.1. Cohort A Eligibility Criteria

### 4.1.1. Cohort A Inclusion Criteria

Each potential subject must satisfy all of the following criteria to be enrolled in Cohort A of the study:

1a. Criterion modified per Amendment 1

1a.1. Criterion modified per Amendment 2

1a.2. Have received a minimum of 1 to a maximum of 3 prior lines of therapy including a proteasome inhibitor (PI) and immunomodulatory therapy (IMiD) either individually or in combination.

- Undergone at least 1 complete cycle of treatment for each line of therapy, unless PD was the best response to that line of therapy.
- Lenalidomide refractory; confirmed PD, per IMWG consensus guidelines ([Rajkumar 2011](#)), during treatment or  $\leq 60$  days after cessation of treatment. Progression on lenalidomide maintenance or non-responsive disease while on lenalidomide for at least 1 cycle will meet this criterion. For subjects with more than 1 prior line of therapy, there is no requirement to be lenalidomide refractory to the most recent line of prior therapy.

2a. Criterion modified per Amendment 1

2a.1. Documented evidence of progressive disease based on investigator's determination of response by the IMWG criteria on or within 6 months of their last regimen ([Attachment 1](#)). Confirmation may be from either central or local testing.

3a.  $\geq 18$  years of age.

4a. Documented diagnosis of multiple myeloma according to IMWG diagnostic criteria ([Attachment 5](#)).

5a. Criterion modified per Amendment 3

5a.1. Measurable disease at Screening as defined by any of the following:

- Serum monoclonal paraprotein (M-protein) level  $\geq 1.0$  g/dL or urine M-protein level  $\geq 200$  mg/24 hours; or
- Light chain multiple myeloma in whom only measurable disease is by serum FLC levels in the serum: Serum immunoglobulin free light chain  $\geq 10$  mg/dL and abnormal serum immunoglobulin kappa lambda free light chain ratio.



Note: Local laboratory assessments may be used to establish measurable disease at Screening, with local laboratory result  $\geq 125\%$  of requirements (eg, M-protein  $\geq 1.25$  g/dL if using local labs).

- For subjects with neither serum nor urine measurable disease, baseline positron emission tomography/ computed tomography (PET/CT) or whole body magnetic resonance imaging (MRI) may be used to satisfy the measurable disease criteria. See [Attachment 22](#) for disease characterization of imaging and [Attachment 23](#) for reporting requirements.

6a. ECOG Performance Status grade of 0 or 1 ([Attachment 7](#)).

7a. Criterion modified per Amendment 1

7a.1. Clinical laboratory values meeting the following criteria during the Screening Phase:

<b>Hematology</b>	
<b>Hemoglobin</b>	$\geq 8.0$ g/dL ( $\geq 5$ mmol/L) (without prior red blood cell [RBC] transfusion within 7 days before the laboratory test; recombinant human erythropoietin use is permitted)*
<b>Platelets</b>	$\geq 50 \times 10^9$ /L (must be without transfusion support in the 7 days prior to the laboratory test)
<b>Absolute Lymphocyte Count (ALC)</b>	$\geq 0.3 \times 10^9$ /L
<b>Absolute Neutrophil Count (ANC)</b>	$\geq 0.75 \times 10^9$ /L (prior growth factor support is permitted but must be without support in the 7 days prior to the laboratory test)
<b>Chemistry</b>	
<b>AST and ALT</b>	$\leq 3.0 \times$ upper limit of normal (ULN)
<b>Estimated Glomerular Filtration Rate</b>	$\geq 40$ mL/min/1.73 m <sup>2</sup> based upon Modified Diet in Renal Disease formula calculation ( <a href="#">Attachment 8</a> ) or a 24-hour urine collection.
<b>Total bilirubin</b>	$\leq 2.0 \times$ ULN; except in subjects with congenital bilirubinemia, such as Gilbert syndrome (in which case direct bilirubin $\leq 1.5 \times$ ULN is required)
<b>Corrected serum calcium</b>	$\leq 12.5$ mg/dL ( $\leq 3.1$ mmol/L) or free ionized calcium $\leq 6.5$ mg/dL ( $\leq 1.6$ mmol/L)

\* For subjects who meet the inclusion criteria at screening, transfusion of RBCs is permitted after screening as needed to maintain a hemoglobin level  $\geq 8.0$  g/dL.

8a. Criterion modified per Amendment 1

8a.1. A woman of childbearing potential must have a negative highly sensitive serum pregnancy test ( $\beta$ -human chorionic gonadotropin [ $\beta$ -hCG]) at screening.

9a. Criterion modified per Amendment 1

9a.1. When a woman is of childbearing potential (See [Attachment 16](#)) the following are required:

- Subject must agree to practice a highly effective method of contraception (failure rate of <1% per year when used consistently and correctly) and agree to remain on a highly effective method of contraception from the time of signing the informed consent form (ICF) until 1 year after receiving a JNJ-68284528 infusion. Examples of highly effective contraceptives include:
  - user-independent methods: 1) implantable progestogen-only hormone contraception associated with inhibition of ovulation; 2) intrauterine device; intrauterine hormone-releasing system; 3) vasectomized partner;
  - user-dependent methods: 1) combined (estrogen- and progestogen-containing) hormonal contraception associated with inhibition of ovulation: oral or intravaginal or transdermal; 2) progestogen-only hormone contraception associated with inhibition of ovulation (oral or injectable)

In addition to the highly effective method of contraception a man:

- Who is sexually active with a woman of childbearing potential must agree to use a barrier method of contraception (eg, condom with spermicidal foam/gel/film/cream/suppository) from the time of signing the ICF until 1 year after receiving a JNJ-68284528 infusion
- Who is sexually active with a woman who is pregnant must use a condom

Women and men must agree not to donate eggs (ova, oocytes) or sperm, respectively, during the study and for 1 year after receiving a JNJ-68284528 infusion.

Note: Hormonal contraception may be susceptible to interaction with the study treatment, which may reduce the efficacy of the contraceptive method.

10a. Subject must sign an ICF indicating that he or she understands the purpose of and procedures required for the study and is willing to participate in the study. Consent is to be obtained prior to the initiation of any study-related tests or procedures that are not part of standard-of-care for the subject's disease.

11a. Willing and able to adhere to the prohibitions and restrictions specified in this protocol.

#### 4.1.2. Cohort A Exclusion Criteria

Any potential subject who meets any of the following criteria will be excluded from participating in Cohort A of the study:

1a. Prior treatment with CAR-T therapy directed at any target.

2a. Any therapy that is targeted to BCMA.

3a. Criterion modified per Amendment 1

3a.1. Criterion modified per Amendment 2

3a.2. Active malignancies (ie, progressing or requiring treatment change in the last 24 months) other than the disease being treated under study. The only allowed exceptions are:

- non-muscle invasive bladder cancer (NMIBC) treated within the last 24 months that is considered completely cured.
- skin cancer (non-melanoma or melanoma) treated within the last 24 months that is considered completely cured.
- non-invasive cervical cancer treated within the last 24 months that is considered completely cured.
- localized prostate cancer (N0M0):
  - with a Gleason score of  $\leq 6$ , treated within the last 24 months or untreated and under surveillance,
  - with a Gleason score of 3+4 that has been treated more than 6 months prior to full study screening and considered to have a very low risk of recurrence, or
  - history of localized prostate cancer and receiving androgen deprivation therapy and considered to have a very low risk of recurrence.
- breast cancer: adequately treated lobular carcinoma in situ or ductal carcinoma in situ, or history of localized breast cancer and receiving antihormonal agents and considered to have a very low risk of recurrence.
- malignancy that is considered cured with minimal risk of recurrence.

4a. Prior antitumor therapy as follows, prior to apheresis:

- Targeted therapy, epigenetic therapy, or treatment with an investigational drug or used an invasive investigational medical device within 14 days or at least 5 half-lives, whichever is less.
- Monoclonal antibody treatment for multiple myeloma within 21 days.
- Cytotoxic therapy within 14 days.
- Proteasome inhibitor therapy within 14 days.
- Immunomodulatory agent therapy within 7 days.
- Radiotherapy within 14 days. However, if the radiation portal covered  $\leq 5\%$  of the bone marrow reserve, the subject is eligible irrespective of the end date of radiotherapy.

- 5a. Criterion modified per Amendment 1
- 5a.1. Ongoing toxicity from previous anticancer therapy must resolve to baseline levels or to Grade 1 or less except for alopecia or peripheral neuropathy.
- 6a. The following cardiac conditions:
- New York Heart Association (NYHA) stage III or IV congestive heart failure
  - Myocardial infarction or coronary artery bypass graft (CABG)  $\leq 6$  months prior to enrollment
  - History of clinically significant ventricular arrhythmia or unexplained syncope, not believed to be vasovagal in nature or due to dehydration
  - History of severe non-ischemic cardiomyopathy
  - Impaired cardiac function (left ventricular ejection fraction [LVEF]  $< 45\%$ ) as assessed by echocardiogram or multiple-gated acquisition (MUGA) scan (performed  $\leq 8$  weeks of apheresis)
- 7a. Received a cumulative dose of corticosteroids equivalent to  $\geq 70$  mg of prednisone within the 7 days prior to apheresis
- 8a. Known active, or prior history of central nervous system (CNS) involvement or exhibits clinical signs of meningeal involvement of multiple myeloma.
- 9a. Stroke or seizure within 6 months of signing ICF.
- 10a. Plasma cell leukemia at the time of screening ( $> 2.0 \times 10^9/L$  plasma cells by standard differential), Waldenström's macroglobulinemia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, and skin changes), or primary amyloid light-chain (AL) amyloidosis.
- 11a. Seropositive for human immunodeficiency virus (HIV).
- 12a. Vaccinated with live, attenuated vaccine within 4 weeks prior to apheresis.
- 13a. Hepatitis B infection as defined according to [Attachment 10](#). In the event the infection status is unclear, quantitative levels are necessary to determine the infection status. ([Hwang 2015](#))
- 14a. Hepatitis C infection defined as (anti-hepatitis C virus [HCV] antibody positive or HCV-RNA positive) or known to have a history of hepatitis C. For subjects with known history of HCV infection, confirmation of sustained virologic response [SVR] is required for study eligibility, defined as  $\geq 24$  weeks after completion of antiviral therapy.

15a. Criterion modified per Amendment 1

15a.1. Subject must not require continuous supplemental oxygen.

16a. Criterion modified per Amendment 3

16a.1. Contraindications, known life threatening allergies, hypersensitivity, or intolerance to cyclophosphamide, fludarabine, or JNJ-68284528 excipients, including DMSO (refer to Investigator's Brochure).

17a. Criterion modified per Amendment 1

17a.1. Criterion Modified per Amendment 2

17a.2. Serious underlying medical condition, such as:

- Evidence of active viral or bacterial infection requiring systemic antimicrobial therapy, or uncontrolled systemic fungal infection
- Active autoimmune disease or a history of autoimmune disease within 3 years
- Overt clinical evidence of dementia or altered mental status
- Any history of Parkinson's disease or other neurodegenerative disorder

18a. Any issue that would impair the ability of the subject to receive or tolerate the planned treatment at the investigational site, to understand informed consent or any condition for which, in the opinion of the investigator, participation would not be in the best interest of the subject (eg, compromise the well-being) or that could prevent, limit, or confound the protocol-specified assessments.

19a. Criterion modified per Amendment 1

19a.1. Pregnant or breast-feeding, or planning to become pregnant while enrolled in this study or within 1 year after receiving JNJ-68284528 infusion.

20a. Criterion modified per Amendment 1

20a.1. Plans to father a child while enrolled in this study or within 1 year after receiving a JNJ-68284528 infusion.

21a. Major surgery within 2 weeks prior to apheresis, or has surgery planned during the study or within 2 weeks after study treatment administration. (Note: subjects with planned surgical procedures to be conducted under local anesthesia may participate.)

22a. Received either of the following:

- An allogenic stem cell transplant within 6 months before apheresis. Subjects who received an allogeneic transplant must be off all immunosuppressive medications for 6 weeks without signs of graft-versus-host disease (GVHD).
- An autologous stem cell transplant  $\leq 12$  weeks before apheresis

23a. Received an investigational intervention (including investigational vaccines) or used an invasive investigational medical device within 15 days prior to apheresis or is currently enrolled in an investigational study

## 4.2. Cohort B Eligibility Criteria

### 4.2.1. Cohort B Inclusion Criteria

Each potential subject must satisfy all of the following criteria to be enrolled in Cohort B of the study:

1b. Criterion modified per Amendment 1

1b.1. Have undergone one line of prior therapy including a PI and an IMiD, per local standard of care.

2b. Criterion modified per Amendment 3

2b.1. Disease progression per IMWG criteria  $\leq 12$  months after ASCT or disease progression  $\leq 12$  months from the start of anti-myeloma therapy for subjects who have not had an ASCT. Confirmation may be from either central or local testing.

3b.  $\geq 18$  years of age.

4b. Documented diagnosis of multiple myeloma according to IMWG diagnostic criteria ([Attachment 5](#)).

5b. Criterion modified per Amendment 3

5b.1. Measurable disease at Screening as defined by any of the following:

- Serum monoclonal paraprotein (M-protein) level  $\geq 1.0$  g/dL or urine M-protein level  $\geq 200$  mg/24 hours; or
- Light chain multiple myeloma in whom only measurable disease is by serum FLC levels in the serum: Serum immunoglobulin free light chain  $\geq 10$  mg/dL and abnormal serum immunoglobulin kappa lambda free light chain ratio.

Note: Local laboratory assessments may be used to establish measurable disease at Screening, with local laboratory result  $\geq 125\%$  of requirements (eg, M-protein  $\geq 1.25$  g/dL if using local labs).

- For subjects with neither serum nor urine measurable disease, baseline PET/CT or whole body MRI may be used to satisfy the measurable disease criteria. See [Attachment 22](#) for disease characterization of imaging and [Attachment 23](#) for reporting requirements.

6b. ECOG Performance Status grade of 0 or 1 ([Attachment 7](#)).

7b. Criterion modified per Amendment 1

7b.1. Clinical laboratory values meeting the following criteria during the Screening Phase:

<b>Hematology</b>	
<b>Hemoglobin</b>	$\geq 8.0$ g/dL ( $\geq 5$ mmol/L) (without prior red blood cell [RBC] transfusion within 7 days before the laboratory test; recombinant human erythropoietin use is permitted)*
<b>Platelets</b>	$\geq 50 \times 10^9/L$ (must be without transfusion support in the 7 days prior to the laboratory test)
<b>Absolute Lymphocyte Count (ALC)</b>	$\geq 0.3 \times 10^9/L$
<b>Absolute Neutrophil Count (ANC)</b>	$\geq 0.75 \times 10^9/L$ (prior growth factor support is permitted but must be without support in the 7 days prior to the laboratory test)
<b>Chemistry</b>	
<b>AST and ALT</b>	$\leq 3.0 \times$ upper limit of normal (ULN)
<b>Estimated Glomerular Filtration Rate</b>	$\geq 40$ mL/min/1.73 m <sup>2</sup> based upon Modified Diet in Renal Disease formula calculation ( <a href="#">Attachment 8</a> ) or a 24-hour urine collection.
<b>Total bilirubin</b>	$\leq 2.0 \times$ ULN; except in subjects with congenital bilirubinemia, such as Gilbert syndrome (in which case direct bilirubin $\leq 1.5 \times$ ULN is required)
<b>Corrected serum calcium</b>	$\leq 12.5$ mg/dL ( $\leq 3.1$ mmol/L) or free ionized calcium $\leq 6.5$ mg/dL ( $\leq 1.6$ mmol/L)

\* For subjects who meet the inclusion criteria at screening, transfusion of RBCs is permitted after screening as needed to maintain a hemoglobin level  $\geq 8.0$  g/dL.

8b. Criterion modified per Amendment 1

8b.1. A woman of childbearing potential must have a negative highly sensitive serum pregnancy test ( $\beta$ -human chorionic gonadotropin [ $\beta$ -hCG]) at screening.

9b. Criterion modified per Amendment 1

9b.1. When a woman is of childbearing potential (See [Attachment 16](#)) the following are required:

- Subject must agree to practice a highly effective method of contraception (failure rate of <1% per year when used consistently and correctly) and agree to remain on a highly effective method of contraception from the time of signing the informed consent form (ICF) until 1 year after receiving a JNJ-68284528 infusion. Examples of highly effective contraceptives include:
  - user-independent methods: 1) implantable progestogen-only hormone contraception associated with inhibition of ovulation; 2) intrauterine device; intrauterine hormone-releasing system; 3) vasectomized partner;
  - user-dependent methods: 1) combined (estrogen- and progestogen-containing) hormonal contraception associated with inhibition of ovulation: oral or intravaginal or transdermal; 2) progestogen-only hormone contraception associated with inhibition of ovulation (oral or injectable)

In addition to the highly effective method of contraception a man:

- Who is sexually active with a woman of childbearing potential must agree to use a barrier method of contraception (eg, condom with spermicidal foam/gel/film/cream/suppository) from the time of signing the ICF until 1 year after receiving JNJ-68284528 infusion
- Who is sexually active with a woman who is pregnant must use a condom

Women and men must agree not to donate eggs (ova, oocytes) or sperm, respectively, during the study and for 1 year after JNJ-68284528 infusion.

Note: Hormonal contraception may be susceptible to interaction with the study treatment, which may reduce the efficacy of the contraceptive method.

10b. Subject must sign an ICF indicating that he or she understands the purpose of and procedures required for the study and is willing to participate in the study. Consent is to be obtained prior to the initiation of any study-related tests or procedures that are not part of standard-of-care for the subject's disease.

11b. Willing and able to adhere to the prohibitions and restrictions specified in this protocol.



#### 4.2.2. Cohort B Exclusion Criteria

Any potential subject who meets any of the following criteria will be excluded from participating in Cohort B of the study:

- 1b. Prior treatment with CAR-T therapy directed at any target.
- 2b. Any therapy that is targeted to BCMA.
- 3b. Criterion modified per Amendment 1
  - 3b.1. Criterion modified per Amendment 2
  - 3b.2. Active malignancies (ie, progressing or requiring treatment change in the last 24 months) other than the disease being treated under study. The only allowed exceptions are:
    - non-muscle invasive bladder cancer (NMIBC) treated within the last 24 months that is considered completely cured.
    - skin cancer (non-melanoma or melanoma) treated within the last 24 months that is considered completely cured.
    - non-invasive cervical cancer treated within the last 24 months that is considered completely cured.
    - localized prostate cancer (N0M0):
      - with a Gleason score of  $\leq 6$ , treated within the last 24 months or untreated and under surveillance,
      - with a Gleason score of 3+4 that has been treated more than 6 months prior to full study screening and considered to have a very low risk of recurrence, or
      - history of localized prostate cancer and receiving androgen deprivation therapy and considered to have a very low risk of recurrence.
    - breast cancer: adequately treated lobular carcinoma in situ or ductal carcinoma in situ, or history of localized breast cancer and receiving antihormonal agents and considered to have a very low risk of recurrence.
    - malignancy that is considered cured with minimal risk of recurrence.
- 4b. Prior antitumor therapy as follows, prior to apheresis:
  - Targeted therapy, epigenetic therapy, or treatment with an investigational drug or used an invasive investigational medical device within 14 days or at least 5 half-lives, whichever is less.
  - Monoclonal antibody treatment for multiple myeloma within 21 days.
  - Cytotoxic therapy within 14 days.
  - Proteasome inhibitor therapy within 14 days.

- Immunomodulatory agent therapy within 7 days.
  - Radiotherapy within 14 days. However, if the radiation portal covered  $\leq 5\%$  of the bone marrow reserve, the subject is eligible irrespective of the end date of radiotherapy.
- 5b. Criterion modified per Amendment 1
- 5b.1. Ongoing toxicity from previous anticancer therapy must resolve to baseline levels or to Grade 1 or less except for alopecia or peripheral neuropathy.
- 6b. The following cardiac conditions:
- NYHA stage III or IV congestive heart failure
  - Myocardial infarction or CABG  $\leq 6$  months prior to enrollment
  - History of clinically significant ventricular arrhythmia or unexplained syncope, not believed to be vasovagal in nature or due to dehydration
  - History of severe non-ischemic cardiomyopathy
  - Impaired cardiac function (LVEF  $< 45\%$ ) as assessed by echocardiogram or MUGA scan (performed  $\leq 8$  weeks of apheresis)
- 7b. Received a cumulative dose of corticosteroids equivalent to  $\geq 70$  mg of prednisone within the 7 days prior to apheresis
- 8b. Known active, or prior history of CNS involvement or exhibits clinical signs of meningeal involvement of multiple myeloma.
- 9b. Stroke or seizure within 6 months of signing ICF.
- 10b. Plasma cell leukemia at the time of screening ( $> 2.0 \times 10^9/L$  plasma cells by standard differential), Waldenström's macroglobulinemia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, and skin changes), or primary AL amyloidosis.
- 11b. Seropositive for HIV.
- 12b. Vaccinated with live, attenuated vaccine within 4 weeks prior to apheresis.
- 13b. Hepatitis B infection as defined according to [Attachment 10](#). In the event the infection status is unclear, quantitative levels are necessary to determine the infection status. ([Hwang 2015](#))
- 14b. Hepatitis C infection defined as (anti-hepatitis C virus [HCV] antibody positive or HCV-RNA positive) or known to have a history of hepatitis C. For subjects with known history of HCV infection, confirmation of sustained virologic response [SVR] is required for study eligibility, defined as  $\geq 24$  weeks after completion of antiviral therapy.

- 15b. Criterion modified per Amendment 1
- 15b.1. Subject must not require continuous supplemental oxygen.
- 16b. Criterion modified per Amendment 3
- 16b.1. Contraindications, known life threatening allergies, hypersensitivity, or intolerance to cyclophosphamide, fludarabine, or JNJ-68284528 excipients, including DMSO (refer to Investigator's Brochure).
- 17b. Criterion modified per Amendment 1
- 17b.1. Criterion modified per Amendment 2
- 17b.2. Serious underlying medical condition, such as:
- Evidence of active viral or bacterial infection requiring systemic antimicrobial therapy, or uncontrolled systemic fungal infection
  - Active autoimmune disease or a history of autoimmune disease within 3 years
  - Overt clinical evidence of dementia or altered mental status
  - Any history of Parkinson's disease or other neurodegenerative disorder
- 18b. Any issue that would impair the ability of the subject to receive or tolerate the planned treatment at the investigational site, to understand informed consent or any condition for which, in the opinion of the investigator, participation would not be in the best interest of the subject (eg, compromise the well-being) or that could prevent, limit, or confound the protocol-specified assessments.
- 19b. Criterion modified per Amendment 1
- 19b.1. Pregnant or breast-feeding, or planning to become pregnant while enrolled in this study or within 1 year after receiving JNJ-68284528 infusion.
- 20b. Criterion modified per Amendment 1
- 20b.1. Plans to father a child while enrolled in this study or within 1 year after receiving JNJ-68284528 infusion.
- 21b. Major surgery within 2 weeks prior to apheresis, or has surgery planned during the study or within 2 weeks after study treatment administration. (Note: subjects with planned surgical procedures to be conducted under local anesthesia may participate.)
- 22b. Received either of the following:
- An allogenic stem cell transplant within 6 months before apheresis. Subjects who received an allogeneic transplant must be off all immunosuppressive medications for 6 weeks without signs of graft-versus-host disease (GVHD).

- An autologous stem cell transplant  $\leq 12$  weeks before apheresis
- 23b. Received an investigational intervention (including investigational vaccines) or used an invasive investigational medical device within 15 days prior to apheresis or is currently enrolled in an investigational study

### 4.3. Cohort C Eligibility Criteria

#### 4.3.1. Cohort C Inclusion Criteria

Each potential subject must satisfy all of the following criteria to be enrolled in Cohort C of the study:

- 1c. Previously treated with a PI, an IMiD, an anti-CD38 monoclonal antibody and BCMA-directed therapy (prior exposure can be from different monotherapy or combination lines of therapy). Subjects may be enrolled in this cohort regardless of dose level/schedule or response obtained to prior BCMA directed therapy.
- Subject must have received prior therapy with at least one BCMA-directed agent (eg, ADC or BiTE), excluding cellular immunotherapy.
- 2c. Documented evidence of progressive disease based on investigator's determination of response by the IMWG criteria ([Attachment 1](#)), either:
- on or within 12 months of their last line of therapy, or,
  - on or within 6 months of prior therapy, and refractory or non-responsive to their most recent line of therapy.

Confirmation may be from either central or local testing.

- 3c.  $\geq 18$  years of age.
- 4c. Documented diagnosis of multiple myeloma according to IMWG diagnostic criteria ([Attachment 5](#)).
- 5c. Criterion modified per Amendment 3
- 5c.1. Measurable disease at Screening as defined by any of the following:
- Serum monoclonal paraprotein (M-protein) level  $\geq 1.0$  g/dL or urine M-protein level  $\geq 200$  mg/24 hours; or
  - Light chain multiple myeloma in whom only measurable disease is by serum FLC levels in the serum: Serum immunoglobulin free light chain  $\geq 10$  mg/dL and abnormal serum immunoglobulin kappa lambda free light chain ratio.

Note: Local laboratory assessments may be used to establish measurable disease at Screening, with local laboratory result  $\geq 125\%$  of requirements (eg, M-protein  $\geq 1.25$  g/dL if using local labs).

- For subjects with neither serum nor urine measurable disease, baseline PET/CT or whole body MRI may be used to satisfy the measurable disease criteria. See [Attachment 22](#) for disease characterization of imaging response and [Attachment 23](#) for reporting requirements.
- 6c. ECOG Performance Status grade of 0 or 1 ([Attachment 7](#)).
- 7c. Criterion modified per Amendment 3
- 7c.1. Clinical laboratory values meeting the following criteria during the Screening Phase:

<b>Hematology</b>	
<b>Hemoglobin</b>	≥8.0 g/dL (≥5 mmol/L) (without prior red blood cell [RBC] transfusion within 3 days before the laboratory test; recombinant human erythropoietin use is permitted)*
<b>Platelets</b>	≥50 x 10 <sup>9</sup> /L (must be without transfusion support in the 3 days prior to the laboratory test)
<b>Absolute Lymphocyte Count (ALC)</b>	≥0.3 x 10 <sup>9</sup> /L
<b>Absolute Neutrophil Count (ANC)</b>	≥0.75×10 <sup>9</sup> /L (prior growth factor support is permitted but must be without support in the 7 days prior to the laboratory test)
<b>Chemistry</b>	
<b>AST and ALT</b>	≤3.0 × upper limit of normal (ULN)
<b>Estimated Glomerular Filtration Rate</b>	≥40 mL/min/1.73 m <sup>2</sup> based upon Modified Diet in Renal Disease formula calculation ( <a href="#">Attachment 8</a> ) or a 24-hour urine collection.
<b>Total bilirubin</b>	≤2.0 × ULN; except in subjects with congenital bilirubinemia, such as Gilbert syndrome (in which case direct bilirubin ≤1.5 × ULN is required)
<b>Corrected serum calcium</b>	≤12.5 mg/dL (≤3.1 mmol/L) or free ionized calcium ≤6.5 mg/dL (≤1.6 mmol/L)

\* For subjects who meet the inclusion criteria at screening, transfusion of RBCs is permitted after screening as needed to maintain a hemoglobin level ≥8.0 g/dL.

- 8c. Women of childbearing potential must have a negative pregnancy test at screening using a highly sensitive serum pregnancy test (β human chorionic gonadotropin [β-hCG]).
- 9c. When a woman is of childbearing potential (See [Attachment 16](#)) the following are required:
- Subject must agree to practice a highly effective method of contraception (failure rate of <1% per year when used consistently and correctly) and agree to remain on a highly effective method of contraception from the time of signing the informed consent form (ICF) until 1 year after receiving a JNJ-68284528 infusion. Examples of highly effective contraceptives include:
    - user-independent methods: 1) implantable progestogen-only hormone contraception associated with inhibition of ovulation; 2) intrauterine device; intrauterine hormone-releasing system; 3) vasectomized partner;

- user-dependent methods: 1) combined (estrogen- and progestogen-containing) hormonal contraception associated with inhibition of ovulation: oral or intravaginal or transdermal; 2) progestogen-only hormone contraception associated with inhibition of ovulation (oral or injectable)

In addition to the highly effective method of contraception a man:

- Who is sexually active with a woman of childbearing potential must agree to use a barrier method of contraception (eg, condom with spermicidal foam/gel/film/cream/suppository) from the time of signing the ICF until 1 year after receiving a JNJ-68284528 infusion.
- Who is sexually active with a woman who is pregnant must use a condom

Women and men must agree not to donate eggs (ova, oocytes) or sperm, respectively, during the study and for 1 year after JNJ-68284528 infusion.

Note: Hormonal contraception may be susceptible to interaction with the study treatment, which may reduce the efficacy of the contraceptive method.

10c. Subject must sign an ICF indicating that he or she understands the purpose of, and procedures required for the study and is willing to participate in the study. Consent is to be obtained prior to the initiation of any study-related tests or procedures that are not part of standard-of-care for the subject's disease.

11c. Willing and able to adhere to the prohibitions and restrictions specified in this protocol.

#### **4.3.2. Cohort C Exclusion Criteria**

Any potential subject who meets any of the following criteria will be excluded from participating in Cohort C of the study:

1c. Prior treatment with cellular immunotherapy (eg, CAR-T) directed at any target.

2c. Criterion modified per Amendment 2

2c.1. Active malignancies (ie, progressing or requiring treatment change in the last 24 months) other than the disease being treated under study. The only allowed exceptions are:

- non-muscle invasive bladder cancer (NMIBC) treated within the last 24 months that is considered completely cured.
- skin cancer (non-melanoma or melanoma) treated within the last 24 months that is considered completely cured.
- non-invasive cervical cancer treated within the last 24 months that is considered completely cured.

- localized prostate cancer (N0M0):
  - with a Gleason score of  $\leq 6$ , treated within the last 24 months or untreated and under surveillance,
  - with a Gleason score of 3+4 that has been treated more than 6 months prior to full study screening and considered to have a very low risk of recurrence, or
  - history of localized prostate cancer and receiving androgen deprivation therapy and considered to have a very low risk of recurrence.
- breast cancer: adequately treated lobular carcinoma in situ or ductal carcinoma in situ, or history of localized breast cancer and receiving antihormonal agents and considered to have a very low risk of recurrence.
- malignancy that is considered cured with minimal risk of recurrence.

3c. Criterion modified per Amendment 2

3c.1. Prior antitumor therapy as follows, prior to apheresis:

- Targeted therapy, epigenetic therapy, or treatment with an investigational drug or used an invasive investigational medical device within 14 days.
- Monoclonal antibody treatment for multiple myeloma within 21 days.
- BCMA-directed Antibody-drug Conjugate or Bispecific T-cell Engager Antibody therapy for multiple myeloma within 5 half-lives of the drug. However, if treated with belantamab mafodotin (GSK2857916) then within 21 days of the drug.
- Cytotoxic therapy within 14 days.
- Proteasome inhibitor therapy within 14 days.
- Immunomodulatory agent therapy within 7 days.
- Radiotherapy within 14 days. However, if the radiation portal covered  $\leq 5\%$  of the bone marrow reserve, the subject is eligible irrespective of the end date of radiotherapy.

4c. Ongoing toxicity from previous anticancer therapy must resolve to baseline levels or to Grade 1 or less except for alopecia or peripheral neuropathy.

5c. The following cardiac conditions:

- New York Heart Association (NYHA) stage III or IV congestive heart failure
- Myocardial infarction or coronary artery bypass graft (CABG)  $\leq 6$  months prior to enrollment
- History of clinically significant ventricular arrhythmia or unexplained syncope, not believed to be vasovagal in nature or due to dehydration
- History of severe non-ischemic cardiomyopathy

- Impaired cardiac function (left ventricular ejection fraction [LVEF] <45%) as assessed by echocardiogram or multiple-gated acquisition (MUGA) scan (performed  $\leq$ 8 weeks of apheresis)
- 6c. Received a cumulative dose of corticosteroids equivalent to  $\geq$ 70 mg of prednisone within the 7 days prior to apheresis
- 7c. Known active, or prior history of central nervous system (CNS) involvement or exhibits clinical signs of meningeal involvement of multiple myeloma.
- 8c. Stroke or seizure within 6 months of signing ICF.
- 9c. Plasma cell leukemia at the time of screening ( $>2.0 \times 10^9/L$  plasma cells by standard differential), Waldenström's macroglobulinemia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, and skin changes), or primary AL amyloidosis.
- 10c. Seropositive for human immunodeficiency virus (HIV).
- 11c. Vaccinated with live, attenuated vaccine within 4 weeks prior to apheresis.
- 12c. Hepatitis B infection as defined according to [Attachment 10](#). In the event the infection status is unclear, quantitative levels are necessary to determine the infection status. ([Hwang 2015](#))
- 13c. Hepatitis C infection defined as (anti-hepatitis C virus [HCV] antibody positive or HCV-RNA positive) or known to have a history of hepatitis C. For subjects with known history of HCV infection, confirmation of sustained virologic response [SVR] is required for study eligibility, defined as  $\geq$ 24 weeks after completion of antiviral therapy.
- 14c. Subject must not require continuous supplemental oxygen.
- 15c. Criterion modified per Amendment 3
- 15c.1. Contraindications, known life threatening allergies, hypersensitivity, or intolerance to cyclophosphamide, fludarabine, or JNJ-68284528 excipients, including DMSO (refer to Investigator's Brochure).
- 16c. Criterion modified per Amendment 2
- 16c.1. Serious underlying medical condition, such as:
- Evidence of active viral or bacterial infection requiring systemic antimicrobial therapy, or uncontrolled systemic fungal infection
  - Active autoimmune disease or a history of autoimmune disease within 3 years
  - Overt clinical evidence of dementia or altered mental status



- Any history of Parkinson's disease or other neurodegenerative disorder
- 17c. Any issue that would impair the ability of the subject to receive or tolerate the planned treatment at the investigational site, to understand informed consent or any condition for which, in the opinion of the investigator, participation would not be in the best interest of the subject (eg, compromise the well-being) or that could prevent, limit, or confound the protocol-specified assessments.
- 18c. Pregnant or breast-feeding, or planning to become pregnant while enrolled in this study or within 1 year after receiving JNJ-68284528 infusion.
- 19c. Plans to father a child while enrolled in this study or within 1 year after receiving JNJ-68284528 infusion.
- 20c. Major surgery within 2 weeks prior to apheresis, or has surgery planned during the study or within 2 weeks after study treatment administration. (Note: subjects with planned surgical procedures to be conducted under local anesthesia may participate.)
- 21c. Received either of the following:
- An allogenic stem cell transplant within 6 months before apheresis. Subjects who received an allogeneic transplant must be off all immunosuppressive medications for 6 weeks without signs of graft-versus-host disease (GVHD).
  - An autologous stem cell transplant  $\leq 12$  weeks before apheresis
- 22c. Received an investigational intervention (including investigational vaccines) or used an invasive investigational medical device within 15 days prior to apheresis or is currently enrolled in an investigational study

#### 4.4. Cohort D Eligibility Criteria

##### 4.4.1. Cohort D Inclusion Criteria

Each potential subject must satisfy all of the following criteria to be enrolled in Cohort D of the study:

- 1d. Have newly diagnosed multiple myeloma per IMWG criteria ([Rajkumar 2014](#)) with a history of 4 to 8 total cycles of initial therapy, including induction, high-dose therapy, and ASCT with or without consolidation (Subjects previously treated for smoldering myeloma are not eligible).

Subject treated with consolidation must have received  $\leq 2$  cycles.

- Received an IMiD or PI or both in combination with a steroid as a part of the induction or consolidation regimen

- Treatment with alkylating therapy (for example cyclophosphamide) and/or monoclonal antibodies (for example, daratumumab) during induction/consolidation is permitted
  - Subjects who have not received consolidation therapy should be approximately 100 days post-ASCT during screening
  - Subjects treated with consolidation therapy should be approximately 160 days post-ASCT during screening
- 2d. Have overall best response <CR and  $\geq$ stable disease, and have not yet evolved to Progressive Disease as assessed per IMWG 2016 criteria
- 3d.  $\geq$ 18 years of age.
- 4d. ECOG Performance Status score of 0 or 1
- 5d. Clinical laboratory values meeting the following criteria during the Screening Phase:

<b>Hematology</b>	
<b>Hemoglobin</b>	$\geq$ 8.0 g/dL ( $\geq$ 5 mmol/L) (without prior red blood cell [RBC] transfusion within 7 days before the laboratory test; recombinant human erythropoietin use is permitted)*
<b>Platelets</b>	$\geq$ 75 x 10 <sup>9</sup> /L (must be without transfusion support in the 7 days prior to the laboratory test)
<b>Absolute Lymphocyte Count (ALC)</b>	$\geq$ 0.3 x 10 <sup>9</sup> /L
<b>Absolute Neutrophil Count (ANC)</b>	$\geq$ 1x10 <sup>9</sup> /L (prior growth factor support is permitted but must be without support in the 7 days prior to the laboratory test)
<b>Chemistry</b>	
<b>AST and ALT</b>	$\leq$ 3.0 x upper limit of normal (ULN)
<b>Estimated Glomerular Filtration Rate</b>	$\geq$ 40 mL/min/1.73 m <sup>2</sup> based upon Modified Diet in Renal Disease formula calculation ( <a href="#">Attachment 8</a> ) or a 24-hour urine collection.
<b>Total bilirubin</b>	$\leq$ 2.0 x ULN; except in subjects with congenital bilirubinemia, such as Gilbert syndrome (in which case direct bilirubin $\leq$ 1.5 x ULN is required)
<b>Corrected serum calcium</b>	$\leq$ 12.5 mg/dL ( $\leq$ 3.1 mmol/L) or free ionized calcium $\leq$ 6.5 mg/dL ( $\leq$ 1.6 mmol/L)

\* For subjects who meet the inclusion criteria at screening, transfusion of RBCs is permitted after screening as needed to maintain a hemoglobin level  $\geq$ 8.0 g/dL

- 6d. Women of childbearing potential must have a negative highly sensitive serum pregnancy test ( $\beta$ -human chorionic gonadotropin [ $\beta$ -hCG]) at screening.

7d. Criterion modified per Amendment 2

7d.1. Criterion modified per Amendment 3

7d.2. When a woman is of childbearing potential (See [Attachment 16](#)) the following are required:

- Subject must agree to practice 2 methods of reliable birth control simultaneously from 4 weeks prior to initiating treatment with lenalidomide until 1 year after receiving a JNJ-68284528 infusion or for 4 weeks following discontinuation of lenalidomide (whichever is later). One of the birth control methods should be a highly effective method of contraception (failure rate of <1% per year when used consistently and correctly; see examples below) and one other effective method (ie, male latex or synthetic condom, diaphragm, or cervical cap) and subject must agree to remain on both methods. Examples of highly effective contraceptives include:
  - user-independent methods: 1) implantable progestogen-only hormone contraception associated with inhibition of ovulation; 2) intrauterine device; intrauterine hormone-releasing system; 3) vasectomized partner;
  - user-dependent method: progestogen-only hormone contraception associated with inhibition of ovulation (oral or injectable). Estrogen-containing hormonal contraception is contraindicated due to increase risk of thromboembolic events with lenalidomide.
  - women of childbearing potential must follow the contraception criteria outlined in the global REVLIMID<sup>®</sup> pregnancy prevention program or equivalent local Risk Evaluation and Mitigation Strategy (REMS), whichever is more stringent, as applicable in their region.

In addition to the highly effective method of contraception a man:

- Must always use a condom during any sexual contact with a woman of childbearing potential, even if they have undergone a successful vasectomy, from the time of signing the ICF until 1 year after receiving a JNJ-68284528 infusion or for 4 weeks after discontinuing lenalidomide (whichever is later).
- Who is sexually active with a woman who is pregnant must use a condom.
- Should agree to practice contraception according to and for the time frame specified in the global REVLIMID<sup>®</sup> pregnancy prevention program or equivalent local REMS, whichever is more stringent, as applicable in their region.

Women and men must agree not to donate eggs (ova, oocytes) or sperm, respectively, during the study and for 1 year after receiving a JNJ-68284528 infusion or for 4 weeks after discontinuing lenalidomide (whichever is later).

Note: Hormonal contraception may be susceptible to interaction with the study treatment, which may reduce the efficacy of the contraceptive method.

- 8d. Subject must sign an ICF indicating that he or she understands the purpose of and procedures required for the study and is willing to participate in the study. Consent is to be obtained prior to the initiation of any study-related tests or procedures that are not part of standard-of-care for the subject's disease.
- 9d. Willing and able to adhere to the prohibitions and restrictions specified in this protocol.

#### 4.4.2. Cohort D Exclusion Criteria

Any potential subject who meets any of the following criteria will be excluded from participating in Cohort D of the study:

- 1d. Received prior treatment with CAR-T therapy directed at any target.
- 2d. Received any therapy that is targeted to BCMA.
- 3d. Criterion modified per Amendment 2
- 3d.1. Active malignancies (ie, progressing or requiring treatment change in the last 24 months) other than the disease being treated under study. The only allowed exceptions are:
- non-muscle invasive bladder cancer (NMIBC) treated within the last 24 months that is considered completely cured.
  - skin cancer (non-melanoma or melanoma) treated within the last 24 months that is considered completely cured.
  - non-invasive cervical cancer treated within the last 24 months that is considered completely cured.
  - localized prostate cancer (N0M0):
    - with a Gleason score of  $\leq 6$ , treated within the last 24 months or untreated and under surveillance,
    - with a Gleason score of 3+4 that has been treated more than 6 months prior to full study screening and considered to have a very low risk of recurrence, or
    - history of localized prostate cancer and receiving androgen deprivation therapy and considered to have a very low risk of recurrence.
  - breast cancer: adequately treated lobular carcinoma in situ or ductal carcinoma in situ, or history of localized breast cancer and receiving antihormonal agents and considered to have a very low risk of recurrence.
  - malignancy that is considered cured with minimal risk of recurrence.

- 4d. Prior anti-tumor therapy, as follows, prior to apheresis:
- Targeted therapy, epigenetic therapy, or treatment with an investigational drug or used as invasive investigational medical device within 14 days or at least 5 half-lives, whichever is less.
  - Monoclonal antibody treatment for multiple myeloma within 21 days.
  - Cytotoxic therapy within 14 days.
  - Proteasome inhibitor therapy within 14 days.
  - Immunomodulatory agent therapy within 7 days.
  - Radiotherapy within 14 days. However, if the radiation portal covered  $\leq 5\%$  of the bone marrow reserve, the subject is eligible irrespective of the end date of radiotherapy.
- 5d. Ongoing toxicity from previous anticancer therapy must resolve to baseline levels or to Grade 1 or less, except for alopecia or peripheral neuropathy.
- 6d. Received a cumulative dose of corticosteroids equivalent to  $\geq 70$  mg of prednisone within the 14 days prior to apheresis.
- 7d. The following cardiac conditions:
- New York Heart Association (NYHA) stage III or IV congestive heart failure
  - Myocardial infarction or coronary artery bypass graft  $\leq 6$  months prior to enrollment
  - History of clinically significant ventricular arrhythmia or unexplained syncope, not believed to be vasovagal in nature or due to dehydration
  - History of severe non-ischemic cardiomyopathy
  - Impaired cardiac function (left ventricular ejection fraction  $< 45\%$ ) as assessed by echocardiogram or multiple-gated acquisition (MUGA) scan (performed  $\leq 8$  weeks of apheresis)
- 8d. Known active, or prior history of central nervous system involvement of myeloma or exhibits clinical signs of meningeal involvement of multiple myeloma.
- 9d. Stroke or seizure within 6 months of signing ICF.
- 10d. Plasma cell leukemia at the time of screening ( $> 2.0 \times 10^9/L$  plasma cells by standard differential), Waldenström's macroglobulinemia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, and skin changes), or primary (AL) amyloidosis.
- 11d. Seropositive for human immunodeficiency virus (HIV).

- 12d. Vaccinated with live, attenuated vaccine within 4 weeks prior to apheresis.
- 13d. Subject must not require continuous supplemental oxygen.
- 14d. Criterion modified per Amendment 3
- 14d.1. Contraindications, known life threatening allergies, hypersensitivity, or intolerance to cyclophosphamide, fludarabine, lenalidomide, or JNJ-68284528 excipients, including DMSO (refer to Investigator's Brochure).
- 15d. Hepatitis B infection as defined according to [Attachment 10](#). In the event the infection status is unclear, quantitative levels are necessary to determine the infection status. ([Hwang 2015](#))
- 16d. Hepatitis C infection defined as (anti-hepatitis C virus [HCV] antibody positive or HCV-RNA positive) or known to have a history of hepatitis C. For subjects with known history of HCV infection, confirmation of sustained virologic response [SVR] is required for study eligibility, defined as  $\geq 24$  weeks after completion of antiviral therapy.
- 17d. Criterion modified per Amendment 2
- 17d.1. Serious underlying medical condition, such as:
- Evidence of active viral or bacterial infection requiring systemic antimicrobial therapy, or uncontrolled systemic fungal infection
  - Active autoimmune disease or a history of autoimmune disease within 3 years
  - Overt clinical evidence of dementia or altered mental status
  - Any history of Parkinson's disease or other neurodegenerative disorder
- 18d. Any issue that would impair the ability of the subject to receive or tolerate the planned treatment at the investigational site, to understand informed consent or any condition for which, in the opinion of the investigator, participation would not be in the best interest of the subject (eg, compromise the well-being) or that could prevent, limit, or confound the protocol-specified assessments.
- 19d. Pregnant or breast-feeding, or planning to become pregnant while enrolled in this study and until 1 year after receiving a JNJ-68284528 infusion or for 4 weeks following discontinuation of lenalidomide (whichever is later).
- 20d. Plans to father a child while enrolled in this study until 1 year after receiving a JNJ-68284528 infusion or for 4 weeks following discontinuation of lenalidomide (whichever is later).

- 21d. Major surgery within 2 weeks prior to apheresis, or has surgery planned during the study or within 2 weeks after JNJ-68284528. (Note: subjects with planned surgical procedures to be conducted under local anesthesia may participate)

#### 4.5. Cohort E Eligibility Criteria

##### 4.5.1. Cohort E Inclusion Criteria

Each potential subject must satisfy all of the following criteria to be enrolled in Cohort E of the study:

- 1e. Have newly diagnosed multiple myeloma without prior therapy and classified as high risk per International Staging System (ISS) stage III criteria, defined as: beta 2 microglobulin ( $\beta 2M$ )  $\geq 5.5$  mg/L ([Attachment 19](#), [Greipp 2005](#)).
- 2e. Not considered candidate for high-dose chemotherapy with stem cell transplantation due to:
  - Being age  $\geq 65$  years; or
  - Age 18-65 years with presence of comorbid condition(s) likely to have a negative impact on tolerability of high-dose chemotherapy with SCT; or
  - Refusal of high-dose chemotherapy with SCT as initial treatment.
- 3e. Documented diagnosis of multiple myeloma according to IMWG diagnostic criteria ([Attachment 5](#)).
- 4e. Measurable disease at Screening as defined by any of the following:
  - Serum monoclonal paraprotein (M-protein) level  $\geq 1.0$  g/dL or urine M-protein level  $\geq 200$  mg/24 hours; or
  - Light chain multiple myeloma in whom only measurable disease is by serum FLC levels in the serum: Serum immunoglobulin free light chain  $\geq 10$  mg/dL and abnormal serum immunoglobulin kappa lambda free light chain ratio.

Note: Local laboratory assessments may be used to establish measurable disease at Screening, with local laboratory result  $\geq 125\%$  of requirements (eg, M-protein  $\geq 1.25$  g/dL if using local labs).
- 5e. ECOG Performance Status grade of 0 or 1 ([Attachment 7](#)).

6e. Clinical laboratory values meeting the following criteria during the Screening Phase:

<b>Hematology</b>	
<b>Hemoglobin</b>	≥8.0 g/dL (≥5 mmol/L) (without prior red blood cell [RBC] transfusion within 7 days before the laboratory test; recombinant human erythropoietin use is permitted)*
<b>Platelets</b>	≥75 x 10 <sup>9</sup> /L (must be without transfusion support in the 7 days prior to the laboratory test)
<b>Absolute Lymphocyte Count (ALC)</b>	≥0.3 x 10 <sup>9</sup> /L
<b>Absolute Neutrophil Count (ANC)</b>	≥1 × 10 <sup>9</sup> /L (prior growth factor support is permitted but must be without support in the 7 days prior to the laboratory test)
<b>Chemistry</b>	
<b>AST and ALT</b>	≤3.0 × upper limit of normal (ULN)
<b>Estimated Glomerular Filtration Rate</b>	≥40 mL/min/1.73 m <sup>2</sup> based upon Modified Diet in Renal Disease formula calculation ( <a href="#">Attachment 8</a> ) or a 24-hour urine collection.
<b>Total bilirubin</b>	≤2.0 × ULN; except in subjects with congenital bilirubinemia, such as Gilbert syndrome (in which case direct bilirubin ≤1.5 × ULN is required)
<b>Corrected serum calcium</b>	≤12.5 mg/dL (≤3.1 mmol/L) or free ionized calcium ≤6.5 mg/dL (≤1.6 mmol/L)

\* For subjects who meet the inclusion criteria at screening, transfusion of RBCs is permitted after screening as needed to maintain a hemoglobin level ≥8.0 g/dL.

7e. A woman of childbearing potential must have a negative highly sensitive serum pregnancy test (β-human chorionic gonadotropin [β-hCG]) at screening.

8e. When a woman is of childbearing potential (See [Attachment 16](#)) the following are required:

- Subject must agree to practice 2 methods of reliable birth control simultaneously from 4 weeks prior to initiating treatment with lenalidomide until 1 year after receiving a JNJ-68284528 infusion or for 4 weeks following discontinuation of lenalidomide or for 3 months after discontinuation of daratumumab (whichever is later). One of the birth control methods should be a highly effective method of contraception (failure rate of <1% per year when used consistently and correctly; see examples below) and one other effective method (ie, male latex or synthetic condom, diaphragm, or cervical cap) and subject must agree to remain on both methods. Examples of highly effective contraceptives include:
  - user-independent methods: 1) implantable progestogen-only hormone contraception associated with inhibition of ovulation; 2) intrauterine device; intrauterine hormone-releasing system; 3) vasectomized partner;



- user-dependent method: progestogen-only hormone contraception associated with inhibition of ovulation (oral or injectable). Estrogen-containing hormonal contraception is contraindicated due to increase risk of thromboembolic events with lenalidomide.
- women of childbearing potential must follow the contraception criteria outlined in the global REVLIMID<sup>®</sup> pregnancy prevention program or equivalent local REMS, whichever is more stringent, as applicable in their region.

In addition to the highly effective method of contraception, a man:

- Must always use a condom during any sexual contact with a woman of childbearing potential, even if they have undergone a successful vasectomy, from the time of signing the ICF until 1 year after receiving a JNJ-68284528 infusion or for 4 weeks after discontinuing lenalidomide (whichever is later).
- Who is sexually active with a woman who is pregnant must use a condom.
- Should agree to practice contraception according to and for the time frame specified in the global REVLIMID<sup>®</sup> pregnancy prevention program or equivalent local REVLIMID<sup>®</sup> pregnancy prevention program, whichever is more stringent.

Women and men must agree not to donate eggs (ova, oocytes) or sperm, respectively, during the study and for 1 year after receiving a JNJ-68284528 infusion or for 4 weeks after discontinuing lenalidomide (whichever is later).

Note: Hormonal contraception may be susceptible to interaction with the study treatment, which may reduce the efficacy of the contraceptive method.

- 9e. Subject must sign an ICF indicating that he or she understands the purpose of and procedures required for the study and is willing to participate in the study. Consent is to be obtained prior to the initiation of any study-related tests or procedures that are not part of standard-of-care for the subject's disease.
- 10e. Willing and able to adhere to the prohibitions and restrictions specified in this protocol.

#### **4.5.2. Cohort E Exclusion Criteria**

Any potential subject who meets any of the following criteria will be excluded from participating in Cohort E of the study:

- 1e. Active malignancies (ie, progressing or requiring treatment change in the last 24 months) other than the disease being treated under study. The only allowed exceptions are:
- non-muscle invasive bladder cancer (NMIBC) treated within the last 24 months that is considered completely cured.
  - skin cancer (non-melanoma or melanoma) treated within the last 24 months that is considered completely cured.

- non-invasive cervical cancer treated within the last 24 months that is considered completely cured.
  - localized prostate cancer (N0M0):
    - with a Gleason score of  $\leq 6$ , treated within the last 24 months or untreated and under surveillance,
    - with a Gleason score of 3+4 that has been treated more than 6 months prior to full study screening and considered to have a very low risk of recurrence, or
    - history of localized prostate cancer and receiving androgen deprivation therapy and considered to have a very low risk of recurrence.
  - breast cancer: adequately treated lobular carcinoma in situ or ductal carcinoma in situ, or history of localized breast cancer and receiving antihormonal agents and considered to have a very low risk of recurrence.
  - malignancy that is considered cured with minimal risk of recurrence.
- 2e. Prior therapy for plasma cell disorder-multiple myeloma other than a short course of corticosteroids (not to exceed 40 mg of dexamethasone, or equivalent per day for a maximum of 4 days, total of 160 mg dexamethasone or equivalent).
- 3e. Frailty index of  $\geq 2$  according to Myeloma Geriatric Assessment score ([Palumbo 2015](#); [Attachment 21](#))
- 4e. Peripheral neuropathy or neuropathic pain Grade 2 or higher, as defined by the National Cancer Institute-Common Terminology Criteria for Adverse Events (NCI-CTCAE) Version 5.
- 5e. The following cardiac conditions:
- New York Heart Association (NYHA) stage III or IV congestive heart failure
  - Myocardial infarction or coronary artery bypass graft (CABG)  $\leq 6$  months prior to enrollment
  - History of clinically significant ventricular arrhythmia or unexplained syncope, not believed to be vasovagal in nature or due to dehydration
  - History of severe non-ischemic cardiomyopathy
  - Impaired cardiac function (left ventricular ejection fraction [LVEF]  $< 45\%$ ) as assessed by echocardiogram or multiple-gated acquisition (MUGA) scan (performed  $\leq 8$  weeks of apheresis)
  - Screening 12-lead ECG showing a baseline QTcF interval  $> 470$  msec
- 6e. Received a strong CYP3A4 inducer within 5 half-lives prior to induction therapy

- 
- 7e. Known active, or prior history of central nervous system (CNS) involvement or exhibits clinical signs of meningeal involvement of multiple myeloma.
- 8e. Stroke or seizure within 6 months of signing ICF.
- 9e. Plasma cell leukemia at the time of screening ( $>2.0 \times 10^9/L$  plasma cells by standard differential), Waldenström's macroglobulinemia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, and skin changes), or primary amyloid light-chain (AL) amyloidosis.
- 10e. Seropositive for human immunodeficiency virus (HIV).
- 11e. Vaccinated with live, attenuated vaccine within 4 weeks prior to apheresis.
- 12e. Hepatitis B infection as defined according to [Attachment 10](#). In the event the infection status is unclear, quantitative levels are necessary to determine the infection status. ([Hwang 2015](#))
- 13e. Hepatitis C infection defined as (anti-hepatitis C virus [HCV] antibody positive or HCV-RNA positive) or known to have a history of hepatitis C. For subjects with known history of HCV infection, confirmation of sustained virologic response [SVR] is required for study eligibility, defined as  $\geq 24$  weeks after completion of antiviral therapy.
- 14e. Subject must not require continuous supplemental oxygen.
- 15e. The following pulmonary conditions
- Chronic obstructive pulmonary disease (COPD) with a Forced Expiratory Volume in 1 second (FEV1)  $<50\%$  of predicted normal (for subjects  $\geq 65$  years old FEV1  $<50\%$  or DLCO  $<50\%$ ).
  - Moderate or severe persistent asthma within the past 2 years (see [Attachment 24](#)), or currently has uncontrolled asthma of any classification. (Note that subjects who currently have controlled intermittent asthma or controlled mild persistent asthma are allowed in the study).
- 16e. Contraindications, known life threatening allergies, hypersensitivity, or intolerance to boron or mannitol, hyaluronidase, sorbitol, corticosteroids, monoclonal antibodies or human proteins, cyclophosphamide, fludarabine, lenalidomide, daratumumab, bortezomib, dexamethasone, or JNJ-68284528 excipients, including DMSO (refer to Investigator's Brochure).
- 17e. Serious underlying medical condition, such as:
- Evidence of active viral or bacterial infection requiring systemic antimicrobial therapy, or uncontrolled systemic fungal infection

- Active autoimmune disease or a history of autoimmune disease within 3 years
  - Overt clinical evidence of dementia or altered mental status
- 18e. Any issue that would impair the ability of the subject to receive or tolerate the planned treatment at the investigational site, to understand informed consent or any condition for which, in the opinion of the investigator, participation would not be in the best interest of the subject (eg, compromise the well-being) or that could prevent, limit, or confound the protocol-specified assessments.
- 19e. Pregnant or breast-feeding, or planning to become pregnant while enrolled in this study and until: 1 year after receiving a JNJ-68284528 infusion, or for 4 weeks following discontinuation of lenalidomide, or until 3 months after daratumumab (whichever is later).
- 20e. Plans to father a child while enrolled in this study until: 1 year after receiving a JNJ-68284528 infusion, or for 4 weeks following discontinuation of lenalidomide, or until 3 months after daratumumab (whichever is later).
- 21e. Major surgery within 2 weeks prior to study start, or has surgery planned during the study or within 2 weeks after study treatment administration. (Note: subjects with planned surgical procedures to be conducted under local anesthesia may participate.)
- 22e. Received an investigational intervention (including investigational vaccines) or used an invasive investigational medical device within 15 days prior to apheresis or is currently enrolled in an investigational study.

#### **4.6. Prohibitions and Restrictions**

Potential subjects must be willing and able to adhere to the following prohibitions and restrictions during the course of the study to be eligible for participation:

1. Refer to Section 8 (Prestudy and Concomitant Therapy) for details regarding prohibited and restricted therapy during the study.
2. Agree to follow all requirements that must be met during the study as noted in the Inclusion and Exclusion Criteria (eg, contraceptive requirements).

### **5. TREATMENT ALLOCATION AND BLINDING**

Randomization will not be used in this study. Subjects will be enrolled into a cohort based on inclusion/exclusion criteria defined for each cohort.

Subjects eligible for both Cohorts A and B, must be enrolled in Cohort B.

As these are single arm study cohorts, blinding procedures are not applicable.

## 6. DOSAGE AND ADMINISTRATION

For this study, study treatment refers to the cyclophosphamide/fludarabine conditioning regimen and JNJ-68284528 for all cohorts. In addition, for Cohort D lenalidomide given post JNJ-68284528 is also considered a study treatment. For Cohort E daratumumab, bortezomib, lenalidomide, and dexamethasone are also considered study treatments. All dosing information must be recorded in the Dosage Administration page of the electronic case report form (eCRF).

### 6.1. Study Treatment Administration

#### 6.1.1. Criteria for Apheresis (all cohorts unless otherwise specified)

The investigator should contact the sponsor if evidence of rapid disease progression or suspected CNS involvement is observed between screening and apheresis. Subjects must meet the following criteria to proceed with apheresis:

- hemoglobin  $\geq 8$  g/dL (PRBC transfusion is permitted)
- platelet count  $\geq 50 \times 10^9/L$  (platelet transfusion is permitted)
- Negative pregnancy test for women of childbearing potential up to 72 hours prior to apheresis
- No supplemental oxygen use to maintain adequate oxygenation
- ECOG performance status grade of 0 or 1
- No investigational agents and anti-myeloma therapy within the timeframe as detailed in the exclusion criteria. For Cohort E, apheresis should occur after 1 or 2 cycles of D-VRd induction treatment and at least 21 days after the last dose of D-VRd in a cycle.
- No focal radiotherapy as specified in the exclusion criteria, except palliative radiotherapy for symptomatic management of bone disease
- No evidence of serious active viral, bacterial, or uncontrolled systemic fungal infection. Subjects on anti-infective agents within 7 days prior to apheresis must receive approval to proceed from sponsor.
- No major surgery as specified in the exclusion criteria
- No new arrhythmia or other cardiac adverse events unless controlled with medical management and approved by the medical monitor

For subjects who require a repeat apheresis, the following assessments should be collected before the second apheresis: weight, hematology laboratory assessments, chemistry laboratory assessments, and echocardiogram or MUGA (if clinically indicated).

If the second apheresis falls outside of the 28-day screening window, all screening assessments (except bone marrow collection) must be repeated (not applicable to Cohort E). Subjects in Cohort E may have a second apheresis after the screening period, as subjects in this cohort will begin induction treatment prior to apheresis. If a second apheresis results in a prolonged treatment-free interval between the 4<sup>th</sup> cycle of induction therapy and initiation of the conditioning regimen, a 5<sup>th</sup> cycle may be given with sponsor approval.

## **6.1.2. Administration of Conditioning Regimen (Cyclophosphamide and Fludarabine) (All Cohorts)**

The site will be notified in writing by the Janssen team that manufacture of JNJ-68284528 has been completed. Each subject will receive a conditioning regimen of intravenous (IV) cyclophosphamide 300 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> daily for 3 doses; sponsor approval must be obtained to modify the conditioning regimen schedule or dose. The dose of fludarabine should be reduced to 24 mg/m<sup>2</sup> for subjects with an eGFR of 30 to 70 mL/min/1.73m<sup>2</sup>. JNJ-68284528 will be administered as a single infusion 5 to 7 days after the start of the conditioning regimen (the first day of conditioning is Day -7 to Day -5, and the day of JNJ-68284528 infusion is Day 1). Cyclophosphamide and fludarabine should be administered using administration procedures and supportive care according to the site's standard of care. JNJ-68284528 should be administered as described in the site investigational product procedures manual (SIPPM) and investigational product preparation instructions (IPPI).

### **6.1.2.1. Criteria for Conditioning Regimen (Cyclophosphamide and Fludarabine Dosing) (All Cohorts)**

The investigator should contact the sponsor if evidence of rapid disease progression or significant change in the subject's clinical status is observed before the start of the conditioning regimen. In addition, subjects must meet the following criteria to proceed with cyclophosphamide and fludarabine dosing:

- Transfusion support is permitted to maintain a hemoglobin of >8.0 g/dL as needed and platelets of  $\geq 50 \times 10^9/L$  until 3 days before the hematology laboratory test, preceding the start of the conditioning regimen.
- Myeloid growth factors are permitted at the investigator's discretion up to 1 day prior to the start of the conditioning regimen. Pegylated myeloid growth factors (ie, pegfilgrastim) are prohibited.
- eGFR  $\geq 30$  mL/min per 1.73 m<sup>2</sup>. The dose of fludarabine should be reduced according to the guidance in Section 6.1.2.
- The investigator must contact the sponsor if the subject has any sign of a reduction in kidney function, which may be manifested by a clinically significant increase in serum creatinine, clinically significant decrease in eGFR, and/or a clinically significant decrease in urine output compared to baseline
- ECOG performance status grade of 0 or 1
- Aspartate aminotransferase (AST)  $\leq 3 \times$  upper limit of normal (ULN)
- Alanine aminotransferase (ALT)  $\leq 3 \times$  ULN.
- Total bilirubin  $\leq 2.0 \times$  ULN; except in subjects with congenital bilirubinemia, such as Gilbert syndrome (in which case direct bilirubin  $\leq 1.5 \times$  ULN is required)
- Negative pregnancy test for women of childbearing potential up to 72 hours prior to the first dose of the conditioning regimen

- Subjects must not have received any antitumor therapy prior to conditioning in the time period outlined in the Exclusion Criteria.
- No active non-hematologic Grade 3 toxicity secondary to bridging therapy
- No signs of infection. For subjects requiring systemic antimicrobial treatment or with temperature  $>38.0^{\circ}\text{C}/100.4^{\circ}\text{F}$  within 7 days prior to the first dose of conditioning regimen, the investigator must receive approval from the sponsor.
- No cumulative dose of corticosteroids equivalent to  $\geq 70$  mg prednisone within the 7 days prior to conditioning regimen dosing. The sponsor should be called for approval if a subject receives corticosteroids at a dose equivalent to  $>10$  mg prednisone per day in the week prior to the start of the conditioning regimen.
- No live, attenuated vaccines within 4 weeks prior to conditioning regimen dosing
- No supplemental oxygen use to maintain adequate oxygenation
- No new arrhythmia or other cardiac adverse events unless controlled with medical management and approved by the medical monitor
- Echocardiogram or MUGA scan for subjects who receive bridging therapy that includes agents with known cardiac toxicity, including but not limited to anthracyclines and carfilzomib (per prescribing information), verification of non-impaired cardiac function (LVEF  $\geq 45\%$ ) should be performed after completion of bridging therapy and prior to the first dose of the conditioning regimen.

### 6.1.3. JNJ-68284528 Administration (All Cohorts)

JNJ-68284528 will be administered as summarized in [Table 10](#).

**Table 10: JNJ-68284528 Administration**

<b>Dose</b>	JNJ-68284528 will be administered in one infusion. The target dose will be the RP2D, of $0.75 \times 10^6$ CAR-positive viable T cells/kg (range: $0.5\text{-}1.0 \times 10^6$ CAR-positive viable T cells/kg) as described in Section 3.3. The maximum total dose of cells to be administered to any subject is $1.0 \times 10^8$ CAR-positive viable T cells (ie, the maximum weight adjusted dose calculated for a 100-kg subject).  If after apheresis and CAR-T cell preparation the quantity of JNJ-68284528 manufactured is not sufficient for dosing at the lower end of the dosing range, dosing for that subject may proceed, provided that a measurable quantity of JNJ-68284528 CAR-positive viable T cells that pass quality testing are generated.
<b>Route/Regimen</b>	JNJ-68284528 IV infusion is to be administered under the supervision of site staff. Refer to the IPPI for JNJ-68284528 infusion instructions.
<b>Dosing Instructions</b>	The actual dose for study treatment administration will be based on the subject's weight (kg) at apheresis.
<b>Schedule of Administration</b>	One intravenous infusion
<b>Hospitalization Requirements</b>	Dependent on the subject's status, medical history, concurrent comorbidities, adequate social support (full-time company of a competent adult) or potential risk factors for developing CAR-T toxicities, including CRS and neurotoxicity, it will be at the Investigator's discretion, patient's willingness, and Sponsor approval whether the subject: <ul style="list-style-type: none"> <li>• will be admitted for inpatient monitoring from the day of infusion (Day 1) through Day 14 post JNJ-68284528 infusion (with potential discharge on Day 10 if there are no CRS, neurotoxicity or other clinically significant events),</li> </ul> OR

**Table 10: JNJ-68284528 Administration**

	<ul style="list-style-type: none"> <li>will receive JNJ-68284528 infusion as an outpatient in close proximity (within 30 min) to the hospital, be monitored for outpatient follow-up and then be admitted for the required inpatient monitoring from Day 5 to Day 14 after JNJ-68284528 infusion (with potential discharge on Day 10 if there are no CRS, neurotoxicity or other clinically significant events) (<a href="#">Attachment 14</a>)</li> </ul> <p>This evaluation should occur at the time of apheresis, prior to administration of the conditioning regimen and again prior to JNJ-68284528 infusion, and in consultation with approval from the sponsor to determine whether outpatient administration and follow-up after JNJ-68284528 infusion is suitable for a given subject and site. The patient must be clinically evaluated after JNJ-68284528 infusion for at least 6 hours before being discharged from the outpatient facility</p> <p>Subject to institutional guidance, local regulations, investigator discretion and sponsor approval, if outpatient JNJ-68284528 administration is being considered, please refer to <a href="#">Attachment 14</a> (JNJ-68284528 Outpatient Administration Guidelines) and Time and Events Schedule <a href="#">Table 1</a>, <a href="#">Table 3</a>, <a href="#">Table 5</a>, and <a href="#">Table 6</a>.</p> <p>For countries or specific hospitals which require hospitalization for all patients treated with cellular therapy, the more stringent requirements for hospitalization will prevail</p> <p>Subjects will be asked to remain within a 1-hour travel time of the hospital and in the company of a competent adult at all times for 1 additional week after hospital discharge, or until study Day 21, whichever is sooner.</p> <p>At the first sign of CRS (such as fever), subjects should be immediately hospitalized for evaluation. Further details regarding management of CRS are described in <a href="#">Table 23</a>.</p> <p>Hospitalization is required for Grade 2, 3, or 4 CAR-T cell-related neurotoxicity (eg, ICANS) temporally associated to CRS. Further details regarding management of CAR-T cell related neurotoxicity are described in <a href="#">Table 24</a>.</p> <p>Hospitalization for neurotoxicity that is not temporally associated with CRS, or any other neurologic adverse events, is at the discretion of the investigator</p>
<b>Vital Sign and Clinical Safety Monitoring</b>	Monitor vital signs as indicated in the Time and Events Schedule ( <a href="#">Table 1</a> , <a href="#">Table 3</a> , <a href="#">Table 5</a> , and <a href="#">Table 6</a> ).

**6.1.3.1. Exceptional Release Criteria**

In the event a JNJ-68284528 product that did not meet pre-specified release criteria is produced during the manufacturing procedures, the sponsor will evaluate the risk/benefit for administration of the affected product and determine if the supply of the product to the treating physician could be considered. If required, approval from the relevant health authorities for use of the product will be obtained in compliance with local regulations regarding notification and approval. In the event the supply of the affected product is deemed appropriate, the investigator will inform the study subject that the product did not meet release specifications prior to administration. The investigator will discuss with the subject the potential risks and benefits of receiving the product, treatment alternatives, including the potential for a second apheresis and discuss alternatives with the patient, including repeat apheresis and alternative therapies.



### 6.1.3.2. Evaluation Prior to Administration of JNJ-68284528 (All Cohorts)

#### JNJ-68284528 Dosing Delays:

Subjects will be evaluated for safety on the day of JNJ-68284528 infusion. If a significant health status change (eg, clinical deterioration, rapidly progressing disease, etc.) occurs following the start of the conditioning regimen (see Section 6.1.2), the investigator must contact the sponsor prior to dosing.

Infusion of JNJ-68284528 must be delayed if any of the following events occur:

- Signs of active infection. Do not administer JNJ-68284528 to patients with active infection. For subjects requiring systemic anti-microbial treatment, or with temperature  $\geq 38.0^{\circ}\text{C}/100.4^{\circ}\text{F}$  Celsius within 48 hours before JNJ-68284528 infusion, investigator must consult with the sponsor prior to dosing.
- Grade  $\geq 3$  non-hematologic toxicities of cyclophosphamide and fludarabine conditioning (except for Grade 3 nausea, vomiting, diarrhea, or constipation). Investigator must consult with the sponsor prior to dosing.

If resolution of these events to Grade  $\leq 1$  takes more than 14 days, the conditioning regimen should be re-administered (cyclophosphamide 300 mg/m<sup>2</sup> and fludarabine 30 mg/m<sup>2</sup> daily for 3 days) after a minimum of 21 days following the first dose of the first conditioning regimen (cyclophosphamide and fludarabine).

### 6.1.3.3. Pre-JNJ-68284528 Infusion Supportive Therapy (All Cohorts)

Prior to JNJ-68284528 infusion, subjects should receive premedication as noted below (Table 11). Corticosteroids should not be used during pre-infusion.

**Table 11: Pre-infusion Medications**

Medication	Dose	Administration
Antihistamine	diphenhydramine (25 -50 mg IV or PO) or equivalent	Oral – administer 1 hour ( $\pm 15$ minutes) prior to JNJ-68284528 infusion
		Or  IV– start infusion 30 minutes ( $\pm 15$ minutes) prior to JNJ-68284528 infusion
Antipyretic	acetaminophen (650 mg to 1,000 mg) or equivalent	Oral or IV - administer 30 minutes ( $\pm 15$ minutes) prior to JNJ-68284528 infusion

### 6.1.4. Lenalidomide (Cohort D)

Starting with the 6<sup>th</sup> subject enrolled in Cohort D (based on DMC recommendation), lenalidomide will be self-administered at a starting dose based on hematologic parameters (Table 12) orally each day on Days 1 through 28 (continuously) of each 28-day cycle until confirmed PD, unacceptable toxicity, or 2 years after initiating lenalidomide treatment (post JNJ-68284528 infusion), whichever occurs first. The first 5 subjects receiving lenalidomide post-JNJ-68284528 will be

staggered similarly as the first 5 subjects without lenalidomide, ie, an observation period of at least 4 weeks will be applied between administration of JNJ-68284528 to the 6<sup>th</sup> through 10<sup>th</sup> subject to allow for subject 28-day safety review prior to next subject dosing. After these 5 subjects receive at least 1 cycle of lenalidomide post-JNJ-68284528, the DMC will reconvene to review safety and any other relevant data. Based on recommendation of the DMC, subsequent subjects in Cohort D may be permitted to enroll and receive JNJ-68284528 followed by lenalidomide without a staggered approach. Women of childbearing potential are required to practice 2 methods of reliable birth control simultaneously beginning 4 weeks prior to receiving lenalidomide.

- After apheresis and prior to administration of cyclophosphamide and fludarabine (conditioning regimen prior to JNJ-68284528 infusion) subjects will receive 1 cycle of lenalidomide at a dose of 10 mg per day upon adequate hematologic recovery from ASCT (absolute neutrophil count [ANC]  $\geq 1 \times 10^9/L$  and platelet count  $\geq 75 \times 10^9/L$ ). Additional cycles of lenalidomide may be given, with sponsor approval, if timing to infusion of JNJ-68284528 is delayed. Alternative bridging therapy in addition, or instead of lenalidomide, is also permitted per sponsor approval. Additionally, if an alternative bridging therapy is used, assessments to address lenalidomide toxicity will not be required in [Table 3](#) (footnote v). The purpose of the bridging therapy is to reduce the myeloma disease burden prior to lymphodepletion chemotherapy and JNJ-58284528 administration.
- After infusion of JNJ-68284528: Applicable subjects (after the first 5 subjects who will receive JNJ-58284528 alone) will initiate lenalidomide a minimum of 21 days post JNJ-68284528 after resolution of cytokine release syndrome (CRS) or neurologic toxicities per ASTCT criteria. The initial dose of lenalidomide will depend on the level of hematologic recovery.
  - Criteria for lenalidomide administration after JNJ-68284528 infusion are summarized in [Table 12](#).

**Table 12: Criteria for Lenalidomide Administration after JNJ-68284528 Infusion – Cohort D**

Hematologic Parameter		Starting Dose of Lenalidomide
ANC	Platelet Count	
$\geq 1.0 \times 10^9/L$	AND $\geq 75 \times 10^9/L$	10 mg daily
0.75 $\times 10^9/L$ to $< 1.0 \times 10^9/L$	AND $\geq 50 \times 10^9/L$	Start 5 mg daily, increase to 10mg per day when ANC is $\geq 1.0 \times 10^9/L$ and the platelet count is $\geq 75 \times 10^9/L$
$\geq 0.75 \times 10^9/L$	AND 50 $\times 10^9/L$ to $< 75 \times 10^9/L$	
$< 0.75 \times 10^9/L$	OR $< 50 \times 10^9/L$	Lenalidomide <u>must</u> be held if <u>either one</u> of these criteria are present

Starting dose of lenalidomide for subjects with renal impairment is provided in [Table 15](#).

Initiation and continuation of lenalidomide treatment is dependent on no additional safety concerns by investigator or sponsor.

- If well tolerated after 3 cycles of lenalidomide treatment at 10 mg, the dose of lenalidomide may be increased to 15 mg per day at the discretion of the investigator.

- For subjects with an eGFR <60 mL/min/1.73m<sup>2</sup>, the lenalidomide dose should be reduced to 5 mg per day. Other dose adjustments should be performed based on local prescribing information and in alignment with the guidance provided in [Table 15](#).

Lenalidomide should be taken orally at about the same time each day, either with or without food. Lenalidomide capsules should be swallowed whole with water. The capsules should not be opened, broken, or chewed. Lenalidomide pill counts should be performed every cycle (ie, every 28 days).

Management guidelines for potential risks of lenalidomide, including dose adjustment, are provided in [Table 13](#) (dose modification guidelines), [Table 14](#) (dose reduction steps) and [Table 15](#) (dosing for subjects with renal impairment).

**Table 13: Dose Modification Guidelines for Lenalidomide**

Toxicity	Lenalidomide Dose Modification
Neutropenia <sup>a</sup> (any of the following): ANC <0.5 x 10 <sup>9</sup> /L	Interrupt lenalidomide treatment.
Return to ANC ≥0.5 x 10 <sup>9</sup> /L when neutropenia is the only observed toxicity	Resume lenalidomide at 1 dose level lower once daily.
For each subsequent drop below <0.5 x 10 <sup>9</sup> /L	Interrupt lenalidomide treatment.
Return to ANC ≥0.5 x 10 <sup>9</sup> /L	Resume lenalidomide at next lower dose level once daily
Thrombocytopenia Platelet count <30 x 10 <sup>9</sup> /L	Interrupt lenalidomide treatment.
Platelet count return to ≥30 x 10 <sup>9</sup> /L	Resume lenalidomide at 1 dose level lower once daily.
For each subsequent drop below 30 x 10 <sup>9</sup> /L	Interrupt lenalidomide treatment.
Platelet count return to ≥30 x 10 <sup>9</sup> /L	Resume lenalidomide at next lower dose level once daily
Grade 2 or 3 skin rash	Lenalidomide interruption or discontinuation should be considered at the investigator's discretion
Angioedema, Grade 4 rash, exfoliative or bullous rash, or if Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) or Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) is suspected or confirmed Progressive Multifocal Leukoencephalopathy (PML)	Permanently discontinue lenalidomide.
Other Grade 3 or 4 toxicities judged to be related to lenalidomide <sup>b</sup>	Interrupt lenalidomide treatment and restart at next lower dose level when toxicity has resolved to ≤ Grade 2 at the investigator's discretion  OR  Consider modification of dosing schedule to 3 weeks of lenalidomide therapy followed by one week rest for each 28 day cycle.

<sup>a</sup> If neutropenia is the only toxicity at any dose level, add G-CSF and maintain the dose level of lenalidomide, at the investigator's discretion.

<sup>b</sup> Adjustments based on other grade 1-2 toxicities (non-hematologic) may be made at investigator's discretion based on standard of care.

**Table 14: Lenalidomide Dose Reduction Steps**

	<b>Starting dose (5mg)</b>	<b>Starting dose (10 mg)</b>	<b>If dose increased (15 mg)</b>
Dose level -1	5mg (Days 1-21 every 28 day cycle)	5 mg (Days 1-28)	10 mg (Days 1-28)
Dose level -2	Not applicable	5 mg (Days 1-21 every 28-day cycle)	5 mg (Days 1-28)
Dose level -3	Not applicable	Not applicable	5 mg (Days 1-21 every 28-day cycle)

**Table 15: Lenalidomide Starting Dose After JNJ-68284528 Infusion for Subjects with Renal Impairment**

<b>Renal function (eGFR) / MDRD</b>	<b>Starting Dose<sup>a</sup></b>
> 60 mL/min/1.73m <sup>2</sup>	10 mg once daily
30 ≤ eGFR ≤ 60 mL/min/1.73m <sup>2</sup>	5 mg once daily
eGFR < 30 mL/min/1.73m <sup>2</sup> , not requiring dialysis	2.5 mg once daily
eGFR < 30 mL/min/1.73m <sup>2</sup> , <i>requiring dialysis</i>	2.5 mg once daily. On dialysis days, the dose should be administered following dialysis

<sup>a</sup> Subsequent dose increase or decrease based on individual patient tolerance.

Management guidelines for potential risks of lenalidomide, including dose adjustment, are provided in the product labeling (REVLIMID<sup>®</sup> USPI and SmPC).

### 6.1.5. Cohort E: Induction and Post JNJ-68284528 Therapy

Subjects in Cohort E will receive D-VRd induction therapy prior to the conditioning therapy and JNJ-68284528 infusion. Subjects will receive D+R consolidation treatment for 2 years post-JNJ-68284528 infusion or until confirmed PD or unacceptable toxicity, whichever occurs first.

Subjects who prematurely discontinue the D-VRd induction treatment (for toxicity or any other reason) may still proceed to receive lymphodepletion and CAR-T therapy after agreement with the sponsor. Subjects receiving D+R consolidation treatment may discontinue one or more agents of this combination if required due to development of toxicities related to these agents.

#### 6.1.5.1. Cohort E: D-VRd Induction Treatment

For subjects participating in Cohort E, details pertaining to D-VRd induction therapy are included in [Table 16](#). Sponsor approval is required for any modification to dosing of the D-VRd induction regimen.

**Table 16: D-VRd Induction Therapy**

	<b>D-VRd Induction therapy (4 Cycles, 21 days per cycle)</b>
Dexamethasone <sup>a</sup> 20 mg PO or IV	On Days 1, 2, 4, 5, 8, 9, 11,12
Bortezomib 1.3 mg/m <sup>2</sup> SC	On Days 1, 4, 8, and 11 of each cycle
Lenalidomide <sup>b</sup> 25 mg PO	On Days 1-14 of 21-day cycle
Daratumumab <sup>c</sup> 1800 mg SC	On Days 1, 8, 15 for Cycles 1-2 [weekly] then Day 1 of Cycles 3-4

IV=intravenous; PO=oral; SC=subcutaneous

- Dispense on Induction Day 1 for self-administration (on daratumumab dosing days, dexamethasone pre-medication for daratumumab injection will replace the daily dose of dexamethasone).
- Dispense on Induction Day 1 of each cycle for self-administration
- Refer to SIPP or approved prescribing information for drug preparation and administration recommendations

### 6.1.5.1.1. Daratumumab

Daratumumab-IV infusion requires a large volume (500 mL to 1000 mL) of infusate, resulting in a median infusion time for the first infusion of 7 hours. Subsequent infusions are approximately 3 to 4 hours. To shorten the infusion time and decrease the risk of infusion-related reactions (IRRs), a technology based on a recombinant human hyaluronidase PH20 (rHuPH20) was developed to facilitate subcutaneous (SC) administration of protein therapeutics. rHuPH20 is the active ingredient of the commercial product Hylenex<sup>®</sup> recombinant (hyaluronidase human injection), which was approved for use in the US in December 2005. Daratumumab has been administered in combination with rHuPH20 in a Phase 1 study (54767414MMY1004; NCT02519452). In some regions, rHuPH20 is also approved in combination with protein therapeutics for SC administration, such as HyQvia (Immune globulin infusion 10% [human] with recombinant human hyaluronidase), as well as anticancer medications such as Herceptin<sup>®</sup> SC (trastuzumab) and MabThera<sup>®</sup> SC (rituximab). Only daratumumab SC will be used in this cohort. Daratumumab SC (trade name, DARZALEX FAST-PRO) was approved by the US FDA in May 2020 and the European Commission in June 2020.

#### 6.1.5.1.1.1. Daratumumab Subcutaneous Preparation

Dara-SC will be provided as a fixed-dosed (1800 mg), combination drug product containing rHuPH20 drug substance (2000 U/mL) and daratumumab drug substance (120 mg/mL) in a single vial. Manuals with detailed descriptions for preparation and administration of daratumumab will be supplied to each pharmacy and site.

#### 6.1.5.1.1.2. Daratumumab Administration

Daratumumab (1800 mg) will be administered by SC injection by manual push over 3-5 minutes in the abdominal subcutaneous tissues in left/right locations, alternating between individual doses. The volume of the SC solution will be 15 mL for the 1800 mg dose. Refer to the locally approved label (USPI, SMPC, or equivalent) or IPPI for additional guidance on SC administration of Dara-SC, as applicable. All subjects will be observed for at least 6 hours after the end of the SC injection during Cycle 1 Day 1 and, if deemed necessary by the investigator, after subsequent injections. Reasons for continued observation on subsequent daratumumab injections may include but are not limited to the following: subjects with a higher risk of respiratory complications (eg, subjects with mild asthma or subjects with COPD who have an FEV1 <80% at screening or developed FEV1

<80% during the study without any medical history), subjects with IRR with first injection of study drug, and subject with a decreased condition on day of dosing compared to prior dosing day. The dose of daratumumab will remain constant throughout the study.

Daratumumab will be administered weekly in Cycles 1-2, every 3 weeks in Cycles 3-4. Every effort should be made to keep subjects on the planned dosing schedule see Time and Events Schedules: Cohort E (Table 5 and Table 6) for acceptable window for treatment. Cycles 1-4 are 21 days.

All daratumumab administrations will be in an outpatient setting. Subjects will receive pre-injection medications and post-injection medications as outlined in Sections 6.1.5.3.

As noted in the Time and Events Schedule (Table 5), vital signs should be monitored extensively on Cycle 1 Day 1 before, and after the first administration of daratumumab. For all other administrations, vital signs should be measured before the start of injection and at the end of the injection. If the subject experiences any significant medical event, then the investigator should assess whether the subject should stay overnight for observation. If the subject has not experienced a significant medical event but is hospitalized overnight only for observation, then the hospitalization should not be reported as a serious adverse event (Section 12.3.2).

On dosing days where the VRd combination is given with daratumumab they should be administered in the following order: lenalidomide, dexamethasone, daratumumab, and bortezomib. Daratumumab will be manufactured and provided under the responsibility of the sponsor. Refer to the daratumumab locally approved label (USPI, SmPC, or equivalent) or IB (as applicable) for a list of excipients.

Daratumumab must be held if any of the following criteria are met at any time, to allow for recovery from toxicity, regardless of relationship to study drug. Daratumumab administration should only be re-initiated after resolution to Grade 2 or better:

- Grade 4 hematologic toxicity, except for Grade 4 lymphopenia
- Grade 3 or higher thrombocytopenia
- Febrile neutropenia
- Neutropenia with infection, of any grade
- Grade 3 or higher non-hematologic toxicities with the following exceptions:
  - Grade 3 nausea that responds to antiemetic treatment within 7 days
  - Grade 3 vomiting that responds to antiemetic treatment within 7 days
  - Grade 3 diarrhea that responds to antidiarrheal treatment within 7 days
  - Grade 3 fatigue that was present at baseline or that lasts for <7 days after the last administration of daratumumab
  - Grade 3 asthenia that was present at baseline or that lasts for <7 days after the last administration of daratumumab

Any adverse event deemed to be related to daratumumab that requires a dose hold of more than 28 days will result in permanent discontinuation of daratumumab unless, upon consultation with the sponsor and the review of safety and efficacy, continuation is agreed upon.

A daratumumab dose held for more than 3 days from the per-protocol administration date for any reason other than toxicities suspected to be related to daratumumab should be brought to the attention of the sponsor at the earliest possible time. Patients missing  $\geq 3$  consecutive planned doses of study drug for reasons other than toxicity should be withdrawn from treatment, unless, upon consultation with the sponsor and the review of safety and efficacy, continuation is agreed upon.

#### **6.1.5.1.2. Bortezomib**

The amount (in mg) of bortezomib to be administered will be determined by body surface area, calculated per a standard nomogram, there is no maximum dose ([Mosteller 1987](#)). The calculated dose of bortezomib may be rounded to the nearest tenth of a mg (or as per institutional practice).

Subjects will receive 1.3 mg/m<sup>2</sup> bortezomib as a subcutaneous injection twice weekly on Days 1, 4, 8, and 11 of each 21-day cycle for Cycles 1-4. For subjects who experience injection-site reactions at the subcutaneous administration site, bortezomib may be administered by IV injection (see local prescribing information). Subjects will not receive bortezomib after the first 4 cycles of treatment. On treatment days when both bortezomib and daratumumab are administered, bortezomib must be administered after the daratumumab administration. If a subject's weight changes by more than 10% from baseline, the dose of bortezomib will be re-calculated. Bortezomib dosing may be delayed up to 48 hours, however subsequent doses must be adjusted to account for the delay. Note that there should be approximately 3 days between doses of bortezomib. Skipped doses of bortezomib will not be made up later in the cycle. For subjects with unacceptable toxicity at the local injection-site despite dose modifications or change in injection concentration, bortezomib can be administered intravenously as a 3 to 5 sec bolus injection. Refer to local prescribing information for further details on either SC or IV administration.

#### **6.1.5.1.3. Lenalidomide**

In Cycles 1 through 4, lenalidomide will be self-administered at a dose of 25 mg orally each day on Days 1 through 14 of each 21-day cycle for subjects with CrCl  $\geq 60$  mL/min. Lenalidomide will be continued until disease progression or unacceptable toxicity whichever occurs first. See [Table 17](#), Renal Impairment for lenalidomide dose adjustments in subjects with renal impairment. On daratumumab administration days, it is recommended that lenalidomide should be administered either prior to or at the same time (preferred) as pre-injection medication. Lenalidomide should be taken as a single dose at the same time daily. If a daily lenalidomide dose is missed, it may be taken if <12 hours have elapsed since the time that it should have been taken. Otherwise, the missed lenalidomide dose should be skipped and not be made up for. Lenalidomide can be taken with or without food. Breaking or dividing the lenalidomide capsule is strongly discouraged.

**6.1.5.1.4. Dexamethasone**

Dexamethasone (or steroid equivalent in accordance with local standards; see [Attachment 12](#) for conversion table) will be self-administered orally, 20 mg on Days 1, 2, 4, 5, 8, 9, 11, 12 of each 21-day cycle for Cycles 1-4. For subjects older than 75 years or underweight (BMI <18.5), the dexamethasone dose may be administered at a dose of 20 mg on days 1, 4, 8, and 11. After dosing with JNJ-68284528, D+R consolidation will include dexamethasone 20 mg oral or IV dose administered as a preinjection medication only on daratumumab injection days during the two years of consolidation therapy of D+R.

**6.1.5.1.5. Dose Modification Guidelines for Bortezomib, Lenalidomide, and Dexamethasone**

Dose modification guidelines for bortezomib, lenalidomide, and dexamethasone are provided in [Table 17](#).

Note that the dose modifications in [Table 17](#) are suggested, but physician discretion and clinical judgment should prevail.

Sponsor approval is required for any modification to dosing of the D-VRd induction regimen.



**Table 17: Dose modification Guidelines for Bortezomib, Lenalidomide, and Dexamethasone**

Body System	NCI-CTC Adverse Event and or Symptom and Category	Bortezomib <sup>g</sup>	Lenalidomide <sup>h</sup>	Dexamethasone <sup>i</sup>
<b>Allergic reactions</b>	Allergic reaction or hypersensitivity Grade 2 OR 3	Hold all therapy. If the toxicity resolves to $\leq$ Grade 1, restart VRd. Reduce by 1 dose-level the suspected medication(s) AND implement appropriate anti-allergic prophylaxis therapy. If the reaction was anaphylactic in nature, do not resume VRd. NOTE: If the reaction was cutaneous in nature, refer to the cutaneous category below.		
	Allergic reaction or hypersensitivity Grade 4	Discontinue VRd.		
<b>Cardiovascular</b>	Fluid Retention (ie, edema) >Grade 3 (limiting function and unresponsive to therapy or anasarca)			Administer diuretics as needed and decrease dexamethasone dose by 1 dose-level; if edema persists despite above measures, decrease dose another dose-level. Permanently discontinue dexamethasone if symptoms persist despite second dose reduction.
<b>Constitutional</b>	Fatigue <sup>a</sup> $\geq$ Grade 3 (ie, severe fatigue interfering with activities of daily living)	Hold the dose until resolved to Grade $\leq$ 2. Consider reduction of lenalidomide or bortezomib by 1 dose-level or change to bortezomib dosing once per week.		
<b>Cutaneous</b>	Non-blistering rash Grade 2	Hold bortezomib therapy. Begin treatment with antihistamines and/or low-dose steroids as per institutional practice. If the toxicity resolves to $\leq$ Grade 1, reduce dose by 1 level and restart bortezomib. Restart with lower concentration formulation. If recurrent consider IV bortezomib.	Consider holding lenalidomide.	

**Table 17: Dose modification Guidelines for Bortezomib, Lenalidomide, and Dexamethasone**

Body System	NCI-CTC Adverse Event and or Symptom and Category	Bortezomib <sup>g</sup>	Lenalidomide <sup>h</sup>	Dexamethasone <sup>i</sup>
	Non-blistering rash ≥ Grade 3 or 4	Hold bortezomib therapies. Begin treatment with antihistamines and/or low-dose steroids as per institutional practice. If the toxicity resolves to ≤ Grade 1, reduce dose by 1 level and restart bortezomib and lenalidomide and continue antihistamines and/or low-dose steroids as per institutional practice. Restart with lower concentration formulation. If recurrent consider IV bortezomib. For Grade 4 toxicity permanently discontinue bortezomib and lenalidomide permanently.		
	Desquamating (blistering) rash-any grade or erythema multiform ≥ Grade 3	Discontinue bortezomib and lenalidomide permanently. Hold other therapies. Begin treatment with antihistamines and/or low-dose steroids as per institutional practice. If the toxicity resolves to ≤ Grade 1, restart other medications.		
<b>Gastrointestinal</b>	Constipation <sup>b</sup> ≥ Grade 3	Hold bortezomib therapy. Upon recovery to ≤ Grade 1, restart bortezomib at 1 dose-reduced level.		
	Diarrhea <sup>c</sup> ≥ Grade 3	Hold bortezomib and consider loperamide therapy. Upon recovery to ≤ Grade 1, restart bortezomib at 1 dose-reduced level.		
	Dyspepsia, gastric or duodenal ulcer, gastritis Grade 1-2 (requiring medical management)			Treat with histamine-2 blockers, sucralfate, or proton pump inhibitor. If symptoms persist despite above measures, decrease dexamethasone dose by 1 dose-level.

**Table 17: Dose modification Guidelines for Bortezomib, Lenalidomide, and Dexamethasone**

<b>Body System</b>	<b>NCI-CTC Adverse Event and or Symptom and Category</b>	<b>Bortezomib<sup>g</sup></b>	<b>Lenalidomide<sup>h</sup></b>	<b>Dexamethasone<sup>i</sup></b>
	Dyspepsia, gastric or duodenal ulcer, gastritis ≥ Grade 3 (requiring hospitalization or surgery)			Hold dexamethasone and consider treatment with histamine-2 blockers, sucralfate, or proton pump inhibitor. Restart and reduce dexamethasone by 1 dose level if symptoms are adequately controlled. If symptoms persist despite above measures, permanently discontinue dexamethasone.
	Acute Pancreatitis			Permanently discontinue dexamethasone.
<b>Hematological</b>	Neutropenia Grade 3 (without complications)	No dose reduction required of bortezomib. Consider treatment with G-CSF.	Hold therapy with lenalidomide until recovery to baseline OR ≤ Grade 2. Consider G-CSF support. Upon recovery if isolated neutropenia, maintain lenalidomide at current dose level. If other hematologic toxicities present reduce lenalidomide by 1 dose level. Maintain bortezomib at current dose. If recurrent episode, reduce lenalidomide by 1 dose-level.	

**Table 17: Dose modification Guidelines for Bortezomib, Lenalidomide, and Dexamethasone**

Body System	NCI-CTC Adverse Event and or Symptom and Category	Bortezomib <sup>g</sup>	Lenalidomide <sup>h</sup>	Dexamethasone <sup>i</sup>
	Neutropenia associated with fever ( $\geq 38.5^{\circ}\text{C}$ ): Grade 3 or neutropenia Grade 4	Hold therapy with all drugs until recovery to baseline OR $\leq$ Grade 2. Consider G-CSF support. Upon recovery if isolated neutropenia, maintain lenalidomide at current dose level. If other hematologic toxicities present reduce lenalidomide by 1 dose level. Maintain bortezomib at current dose. If recurrent episode, reduce lenalidomide and bortezomib by 1 dose-level.		
	Thrombocytopenia Grade 3 (without complications)	No dose reduction required for bortezomib.	Reduce lenalidomide by 1 dose-level for the remainder of the cycle.	
	Platelet count $\leq 30 \times 10^9/\text{L}$ or ANC $\leq 0.75 \times 10^9/\text{L}$ on a bortezomib dosing day	Hold bortezomib dose.		
	Platelet count $< 25,000/\mu\text{L}$ (ie, Grade 4) or Grade 3 thrombocytopenia with bleeding	Hold therapy with all drugs until recovery to baseline OR $\leq$ Grade 2. Upon recovery, reduce bortezomib 1 dose level, hold lenalidomide for remainder of the cycle and decrease by 1 dose level at start of next cycle.		
<b>Infection</b>	Herpes Zoster <sup>d</sup> activation or reactivation ANY grade	Hold ALL therapies until lesions are dry. If not already underway, begin antiviral treatment Once the infection is resolved all medications can be restarted without a dose reduction; however, continued antiviral prophylaxis is required.		
<b>Musculoskeletal</b>	Muscle weakness $>$ Grade 2 (symptomatic and interfering with function +/- interfering with activities of daily living)			Decrease dexamethasone dose by 1 dose-level. If weakness persists despite above measures, decrease dose by 1 <i>further</i> dose-level. If symptoms <i>still</i> persist, permanently discontinue dexamethasone.
<b>Metabolic</b>	Hyperglycemia $\geq$ Grade 3			Treatment with insulin or oral hypoglycemics. If uncontrolled despite above measures, decrease dose by 1 dose-level until levels are satisfactory.

**Table 17: Dose modification Guidelines for Bortezomib, Lenalidomide, and Dexamethasone**

Body System	NCI-CTC Adverse Event and or Symptom and Category	Bortezomib <sup>g</sup>	Lenalidomide <sup>h</sup>	Dexamethasone <sup>i</sup>
<b>Neurological<sup>c</sup></b>	Peripheral Neuropathy (Sensory or Motor) and/or Neuropathic Pain	Grade 1 (paresthesias and/or loss of reflexes) without pain or loss of function	No action required.	
		Grade 1 with pain or Grade 2 (interfering with function but not with activities of daily living)	Change schedule to once per week; if recurrence Reduce bortezomib by 1 dose-level	
		Grade 2 with pain or Grade 3 (interfering with activities of daily living)	Hold bortezomib until toxicity resolves to <Grade 2. When toxicity resolves, reinstate with a reduction by 1 dose-level and change bortezomib treatment schedule to once per week	
		Grade 4 (permanent sensory loss that interferes with function) and/or severe autonomic neuropathy	Discontinue bortezomib permanently.	
<b>Neuro-psychological</b>	Confusion or mood alteration >Grade 2 (interfering with function ± interfering with activities of daily living)			Hold dexamethasone until symptoms resolve. Restart with 1 dose-level reduction. If symptoms persist despite above measures, permanently discontinue dexamethasone.
<b>Thromboembolic</b>	Venous and /or pulmonary thromboembolism ≥ Grade 3 [Deep vein thrombosis or cardiac thrombosis intervention indicate; eg: anticoagulation, lysis, filter, invasive procedure.]		Stop until toxicity resolves and, if not already given, start anticoagulation therapy. Restart lenalidomide and dexamethasone at full dose after adequate anticoagulation.	

**Table 17: Dose modification Guidelines for Bortezomib, Lenalidomide, and Dexamethasone**

Body System	NCI-CTC Adverse Event and or Symptom and Category	Bortezomib <sup>g</sup>	Lenalidomide <sup>h</sup>	Dexamethasone <sup>i</sup>
Renal Impairment	Moderate renal impairment-CrCl <sup>f</sup> 30-59 mL/min		Lenalidomide should be given at a dose of 10 mg daily	
	Severe renal impairment-CrCl <sup>f</sup> <30 mL/min (not requiring dialysis)		Lenalidomide should be given at a dose of 15 mg every 48 hrs	
	End-stage renal disease-CrCl <sup>f</sup> <30 mL/min (requiring dialysis)		Lenalidomide should be given at a dose of 5 mg daily. Administer dose after dialysis.	
Other toxicities	Any reported ≥ Grade 3	Determine drug attribution of the toxicity and hold the therapy(ies) as appropriate. If toxicity resolves to ≤ Grade 1, resume therapy with 1 level of dose reduction for suspect drug.		
Abbreviations: ANC=absolute neutrophil count; CrCl=creatinine clearance; IV=intravenous; NCI-CTCAE=National Cancer Institute Common Terminology Criteria for Adverse Events; G-CSF=Granulocyte colony stimulating factor; VRd=VELCADE, lenalidomide, and dexamethasone				
<p><sup>a</sup> Determine if fatigue is possibly not medication-related but due to an underlying cause (eg, infection, progression of disease, diarrhea, anemia, depression) and treat these symptoms/causes as appropriate.</p> <p><sup>b</sup> Prior to dose reduction of medications, consider/eliminate other possible causes of constipation.</p> <p><sup>c</sup> Prior to dose reduction of medications, consider/eliminate other possible causes (ie, bacterial or viral infections) of diarrhea.</p> <p><sup>d</sup> In the event that a subject is already receiving antiviral treatment at the time of the Herpes Zoster activation, consider switching to or adding another antiviral agent.</p> <p><sup>e</sup> The neurotoxicity-directed questionnaire is a useful tool for determining the presence and intensity of neuropathic pain and/or peripheral neuropathy from the subject's perspective. Neuropathic symptoms are more prominent than abnormalities on the clinical examination. After the subject completes the neurotoxicity-directed questionnaire, the questionnaire should be reviewed to assist with the evaluation of the onset and intensity of peripheral neuropathy and other neurotoxicities that may require intervention or dose modification.</p> <p><sup>f</sup> CrCl = creatinine clearance. Estimated by creatinine clearance as calculated by the Cockcroft-Gault formula and adjusted for body weight in subjects with a body mass index &gt;30 kg/m<sup>2</sup>. The eGFR (MDRD) or CKD-epi formulas can also be utilized to assess renal function.</p> <p><sup>g</sup> See Table 18 for dose levels</p> <p><sup>h</sup> See Table 19 for dose levels</p> <p><sup>i</sup> See Table 20 for dose levels</p>				

**Table 18: Dose Modifications for Bortezomib**

Starting Dose	First Dose Reduction	Second Dose Reduction	Third Dose Reduction
Bortezomib 1.3 mg/m <sup>2</sup>	Bortezomib 1.0 mg/m <sup>2</sup>	Bortezomib 0.7 mg/m <sup>2</sup>	Discontinue bortezomib

**Table 19: Dose Modifications for Lenalidomide**

Starting Dose	First Dose Reduction	Second Dose Reduction	Third Dose Reduction	Fourth Dose Reduction	Fifth Dose reduction
Lenalidomide 25 mg	Lenalidomide 20 mg	Lenalidomide 15 mg	Lenalidomide 10 mg	Lenalidomide 5 mg	Discontinue Lenalidomide

**Table 20: Dose Modifications for Dexamethasone**

First Dose Reduction	Second Dose Reduction	Third Dose Reduction
Reduce dexamethasone by 50% from starting dose	Skip dexamethasone on days when daratumumab is not given	Discontinue dexamethasone

### 6.1.5.2. Cohort E: D+R Consolidation Treatment

After infusion of JNJ-68284528, all subjects in Cohort E will initiate lenalidomide a minimum of 21 days post JNJ-68284528 after resolution of CRS or neurologic toxicities associated with JNJ-68284528. The initial dose of lenalidomide will depend on the level of hematologic recovery.

- Criteria for lenalidomide administration after JNJ-68284528 infusion are summarized in [Table 21](#).

**Table 21: Criteria for Lenalidomide Administration after JNJ-68284528 Infusion – Cohort E**

Hematologic Parameter			Starting Dose of Lenalidomide
ANC		Platelet Count	
$\geq 1.0 \times 10^9/L$	AND	$\geq 75 \times 10^9/L$	10 mg daily
$0.75 \times 10^9/L$ to $< 1.0 \times 10^9/L$	AND	$\geq 50 \times 10^9/L$	Start 5mg daily, increase to 10mg per day when ANC is $\geq 1.0 \times 10^9/L$ <u>and</u> the platelet count is $\geq 75 \times 10^9/L$
$\geq 0.75 \times 10^9/L$	AND	$50 \times 10^9/L$ to $< 75 \times 10^9/L$	
$< 0.75 \times 10^9/L$	OR	$< 50 \times 10^9/L$	Lenalidomide <u>must</u> be held if <u>either one</u> of these criteria are present

ANC=absolute neutrophil count

- For subjects with an eGFR  $< 60 \text{ mL/min/1.73m}^2$ , dose reduction of lenalidomide should be performed based on local prescribing information and in alignment with the guidance provided in [Table 15](#).
- Initiation and continuation of lenalidomide treatment is dependent on no additional safety concerns by investigator or sponsor.
  - If well tolerated after 3 cycles of lenalidomide treatment, the dose of lenalidomide may be increased to 15 mg per day at the discretion of the investigator;

- further titration at investigator discretion, by 5mg increments may proceed after 6 cycles until a maximum dose of 25 mg daily (eg, lenalidomide Cycle 1: 10 mg, Cycle 2: 10 mg, Cycle 3: 10 mg, Cycle 4: 15 mg, Cycle 5: 15 mg, Cycle 6: 15 mg, Cycle 7: 20 mg, Cycle 8: 25 mg).

Lenalidomide will be self-administered orally each day on Days 1 through 21 (21 days on, 7 days off) of each 28-day cycle until confirmed PD, unacceptable toxicity, or 2 years after JNJ-68284528 infusion, whichever occurs first.

The initial dose of lenalidomide will depend on the level of hematologic recovery. The criteria for lenalidomide administration after JNJ-68284528 infusion and the management guidelines for potential risks of lenalidomide, including dose adjustment, are described in [Table 13](#) (dose modification guidelines), [Table 19](#) (dose reduction steps) and [Table 15](#) (dosing for subjects with renal impairment).

Daratumumab will be initiated approximately 60-100 days after JNJ-68284528 and upon adequate recovery from ongoing toxicities. Initiate daratumumab once platelet and absolute neutrophil count (ANC) are  $\geq 50 \times 10^9/L$  and  $\geq 1.0 \times 10^9/L$ , respectively. The 1<sup>st</sup> dose of daratumumab should be on Day 1 of a subsequent cycle of lenalidomide in order to align cycle days, see [Table 22](#). Subjects must complete 1 cycle of lenalidomide before starting daratumumab, even if hematologic recovery occurs sooner.

**Table 22: D+R Consolidation Therapy**

	<b>D+R Consolidation therapy (up to 2 years, 28 days per cycle)</b>
Daratumumab <sup>a</sup> 1800 mg SC <sup>b</sup>	Day 1
Lenalidomide <sup>c,d</sup> 10 mg PO	On Days 1-21 of 28-day cycle

IV=intravenous; PO=oral; SC=subcutaneous

- Refer to SIPP or approved prescribing information for drug preparation and administration recommendations
- Medications to be administered prior to daratumumab infusion are described in [Section 6.1.5.3](#)
- Dispense on Consolidation Day 1 of each cycle for self-administration
- The initial dose of lenalidomide after JNJ-68284528 infusion will depend on the level of hematologic recovery and may be titrated at investigator discretion to a maximum dose of 25 mg as described in [Section 6.1.5.2](#) and [Table 21](#).

Daratumumab 1800 mg will be administered by SC injection, every 4 weeks onwards until confirmed PD, unacceptable toxicity, or 2 years after the first cycle of lenalidomide, whichever occurs first.

Daratumumab must be held if any of the criteria specified in [Section 6.1.5.1.1.2](#) are met at any time, to allow for recovery from toxicity, regardless of relationship to study drug or discontinued as specified. Daratumumab administration should only be re-initiated after resolution to Grade 2 or better.



### 6.1.5.3. Cohort E: Medications Related to Daratumumab Dosing

To decrease the risk of infusion-related reactions, all subjects will receive the following medications 1 to 3 hours prior to daratumumab administration:

- Paracetamol (acetaminophen) 650-1000 mg IV or orally (PO).
- An antihistamine: diphenhydramine 25-50 mg IV or PO, or equivalent (see [Attachment 25](#) for a list of antihistamines that may be used). Avoid IV promethazine.
- Dexamethasone 20 mg IV or oral
  - The dexamethasone 20 mg oral or IV dose administered as a preinjection medication on daratumumab injection days, replaces the oral dexamethasone dose for that day (if applicable).
  - If adverse events due to steroid during pre-medication for consolidation therapy with daratumumab. A reduction to 10 mg IV or PO dexamethasone will be permitted for preinjection medication. If dexamethasone is contraindicated due to adverse events, the sponsor should be consulted regarding before continuing daratumumab dosing.
- Substitutions for dexamethasone are allowed (refer to [Attachment 12](#)).
- Predose administration of a leukotriene inhibitor (montelukast 10 mg PO) is recommended on Cycle 1 Day 1 of induction therapy. The leukotriene inhibitor may be administered up to 24 hours before the daratumumab dose.

If necessary, due to timing constraints, all PO pre-administration medications may be administered outside of the clinic on the day of the injection, provided they are taken 1-3 hours before the daratumumab dose.

Prophylaxis for herpes zoster reactivation is recommended (see Section [8.1](#)).

In addition, for any patients with a history of chronic obstructive pulmonary disease, consider prescribing post-infusion medications such as short and long-acting bronchodilators, and inhaled corticosteroids.

## 6.2. Management Guidelines for Potential Risks (All Cohorts)

### 6.2.1. Management of Cytokine Release Syndrome

In the Legend-2 study, CRS was reported in approximately 92% of subjects who received LCAR-B38M CAR-T cells. Most CRS events were Grade 1 or Grade 2. All events of CRS started with fever after the infusion of CAR-T therapy (See Section [1.1.5](#)). Of the subjects who developed CRS, approximately 84% experienced transiently increased aspartate aminotransferase (AST). AST increase was Grade 3 or Grade 4 in 31% and 6% of subjects with CRS, respectively. If CRS is suspected, subjects should be monitored for increased AST, and consumptive coagulopathy, indicated by an increase in D-dimers and a decrease in fibrinogen if CRS is suspected.

Symptoms indicative of CRS may include, but are not limited to, fever (with or without rigors), arthralgia, nausea, vomiting, tachypnea, hypoxia, tachycardia, hypotension, headache, confusion, tremor, delirium, dyspnea, pulmonary edema, and capillary leak ([Lee 2014](#)). Potentially life-

threatening complications of CRS may include cardiac dysfunction, adult respiratory distress syndrome, neurologic toxicity, renal failure, hepatic failure, and disseminated intravascular coagulation.

Laboratory testing to monitor for disseminated intravascular coagulation, a manifestation of CRS, should be carried out in addition to daily monitoring of chemistry and hematology assessments (including ferritin and CRP) when fever or other signs of potential CRS are present (see [Table 1](#), [Table 3](#), [Table 6](#), and [Table 7](#)). In addition, pulmonary, renal and hepatic function will be monitored closely (see [Table 1](#) and [Table 3](#)). Cytokine release syndrome will be captured as an adverse event of special interest (see Section [12.3.3](#)).

Rarely, severe CRS can evolve into a presentation consistent with hemophagocytic lymphohistiocytosis/macrophage activation syndrome (HLH/MAS) that may require additional therapy. In these cases, laboratory testing may reveal high serum levels of ferritin, lactate dehydrogenase, soluble CD25, and cytokines (such as IFN $\gamma$  and IL-6), and low serum levels of fibrinogen ([Neelapu 2018](#)).

Trained clinical personnel should be prepared to intervene in the event of CRS. Resources necessary for resuscitation (ie, agents such as epinephrine and aerosolized bronchodilator; medical equipment such as oxygen, tracheostomy equipment, and a defibrillator) should be readily available. Tocilizumab must be available prior to administration of JNJ-68284528. Vital signs and laboratory parameters must be monitored at regular intervals until normal. Additional specimens for pharmacokinetic and pharmacodynamic testing should be collected as per the schedule outlined in the Time and Events schedules ([Table 2](#), [Table 4](#), [Table 7](#), and [Table 8](#)).

Infection and CRS may have a similar presentation. Therefore, investigators are strongly encouraged to evaluate for an infection at the first signs or symptoms of CRS. Cultures and imaging should be obtained: the clinical signs and symptoms should determine which tests are appropriate.

Recommendations for the clinical management of CRS are provided in [Table 23](#). At the first sign of CRS (such as fever), subjects should be immediately hospitalized for evaluation. The use of myeloid growth factors, particularly granulocyte colony-stimulating factor (G-CSF), should be avoided during CRS. Tocilizumab intervention may be considered with presenting symptom of fever per investigator discretion in the absence of clear infectious etiology and early tocilizumab should be considered in subjects at high risk of severe CRS (for example, high baseline tumor burden, early fever onset, or persistent fever after 24 hours of symptomatic treatment). Other monoclonal antibodies targeting cytokines (for example, anti-IL1 and/or anti-TNF $\alpha$ ) may be used based on institutional practice, especially for cases of CRS which does not respond to tocilizumab. Therapy directed at reduction or elimination of CAR-T cells, including chemotherapy, may be considered in consultation with the sponsor for subjects who develop high grade CRS with laboratory findings overlapping with HLH/MAS (including hyperferritinemia) that remains severe or life-threatening following prior therapies, including tocilizumab and corticosteroids.

**Table 23: Guidelines for the Management of Cytokine Release Syndrome**

<b>CRS Grade</b>	<b>Presenting Symptoms</b>	<b>Tocilizumab<sup>a</sup></b>	<b>Corticosteroids<sup>b</sup></b>
Grade 1	Temperature $\geq 38^{\circ}\text{C}/100.4^{\circ}\text{F}^{\text{c}}$	May be considered	N/A
Grade 2	Temperature $\geq 38^{\circ}\text{C}/100.4^{\circ}\text{F}^{\text{c}}$ with either:  Hypotension responsive to fluids and not requiring vasopressors.  Or, oxygen requirement of low-flow nasal cannula <sup>d</sup> or blow-by	Administer tocilizumab <sup>b</sup> 8 mg/kg intravenously over 1 hour (not to exceed 800 mg).  Repeat tocilizumab every 8 hours as needed if not responsive to intravenous fluids or increasing supplemental oxygen.  Limit to a maximum of 3 doses in a 24-hour period; maximum total of 4 doses.	Manage per guidance below if no improvement within 24 hours of starting tocilizumab.
Grade 3	Temperature $\geq 38^{\circ}\text{C}/100.4^{\circ}\text{F}^{\text{c}}$ with either:  Hypotension requiring one vasopressor with or without vasopressin.  Or, oxygen requirement of high-flow nasal cannula <sup>d</sup> , facemask, non-rebreather mask, or Venturi mask	Administer tocilizumab 8 mg/kg intravenously over 1 hour (not to exceed 800 mg).  Repeat tocilizumab every 8 hours as needed if not responsive to intravenous fluids or increasing supplemental oxygen.  Limit to a maximum of 3 doses in a 24-hour period; maximum total of 4 doses.	If no improvement, administer methylprednisolone 1 mg/kg intravenously twice daily or equivalent dexamethasone (eg, 10 mg intravenously every 6 hours).  Continue corticosteroids use until the event is Grade 1 or less, then taper over 3 days.
Grade 4	Temperature $\geq 38^{\circ}\text{C}/100.4^{\circ}\text{F}^{\text{c}}$ with either:  Hypotension requiring multiple vasopressors (excluding vasopressin).  Or, oxygen requirement of positive pressure (eg, CPAP, BiPAP, intubation, and mechanical ventilation)	Administer tocilizumab 8 mg/kg intravenously over 1 hour (not to exceed 800 mg).  Repeat tocilizumab every 8 hours as needed if not responsive to intravenous fluids or increasing supplemental oxygen.  Limit to a maximum of 3 doses in a 24-hour period; maximum total of 4 doses.	As above or administer methylprednisolone 1000 mg intravenously per day for 3 days per investigator discretion.  If no improvement or if condition worsens, consider alternate immunosuppressants. <sup>b</sup>

CPAP=Continuous Positive Airway Pressure; BiPAP=Bilevel Positive Airway Pressure

a: Refer to tocilizumab prescribing information for details

b: Monoclonal antibodies targeting cytokines may be considered based on institutional practice for unresponsive CRS.

c: Attributed to CRS. Fever may not always be present concurrently with hypotension or hypoxia as it may be masked by interventions such as antipyretics or anticytokine therapy (eg, tocilizumab or steroids).

d: Low-flow nasal cannula is  $\leq 6$  L/min, and high-flow nasal cannula is  $>6$  L/min.

Note: At first sign of CRS (such as fever) subjects should be immediately hospitalized for evaluation.

Supportive care for CRS (including but not limited to anti-pyretic agents, IV fluid support, vasopressors, supplemental oxygen, etc.) should be administered according to the clinical manifestations of the subject's illness. Similarly, ancillary testing such as B-type natriuretic peptide (BNP) assessment, echocardiograms, arterial blood gas, assessments of coagulation laboratory tests, etc. should be performed if clinically indicated.

## **6.2.2. Neurologic Toxicities**

Based on the specific mode-of-action of JNJ-68284528, severe or serious neurologic toxicities (including CAR-T cell-related neurotoxicity, eg, ICANS [Immune Effector Cell-Associated Neurotoxicity Syndrome]) may occur (Section 6.2.2.1). Additionally, subjects should be monitored for neurotoxicity for 1 year post JNJ-68284528 infusion (Section 6.2.2.2).

### **6.2.2.1. CAR-T Cell-related Neurotoxicity (Immune Effector Cell-Associated Neurotoxicity Syndrome [ICANS])**

Subjects should have the Immune Effector Cell-associated Encephalopathy (ICE) Assessment Tool (ICE-Tool; [Attachment 3](#)) performed at baseline (within 24 hours prior to infusion of JNJ-68284528 infusion) and daily after the first symptoms of CAR-T cell related neurotoxicity (eg, ICANS) are suspected and until resolution. Consider performing ICE-Tool more frequently until neurotoxicity symptoms resolve. Consider performing neuroimaging (eg, magnetic resonance imaging [MRI]) at screening and/or neurology consultation if pre-existing disease is suspected; see Section 9.7 Safety Evaluations.

Subjects should be monitored for neurologic toxicities including, but not restricted to: speech disorders, aphasia, convulsions, disturbances in consciousness, confusion, disorientation, or coordination and balance disorders. If these or other neurologic toxicities are observed, regardless of causality, then the sponsor's medical monitor must be consulted. Hospitalization is required for Grade 2, 3, or 4 CAR-T cell-related neurotoxicity (eg, ICANS) temporally associated to CRS.

At the first sign of neurotoxicity, neurology consultation and evaluation should be considered to rule out alternative etiologies including viral origin (ie, human herpesvirus [HHV] HHV-6, HHV-7). Subjects who have a lumbar puncture as part of their neurologic work up should have a sample of cerebral spinal fluid also sent to the sponsor for additional testing. For signs of seizures or raised intracranial pressure (ICP)/cerebral edema, consider neuroimaging (CT/MRI), transfer the subject to the intensive care unit (ICU) and treat according to institutional guidelines or practices.

General management for CAR-T cell-related neurotoxicity (eg, ICANS) with or without concurrent CRS is summarized in [Table 24](#). All neurologic adverse events, including CAR-T cell-related neurotoxicity (eg, ICANS), will be captured as an adverse event of special interest (see Section 12.3.3).

**Table 24: Guidelines for the Management of CAR-T Cell-related Neurotoxicity (eg, ICANS)**

ICANS Grade	Presenting Symptoms <sup>a</sup>	Concurrent CRS	No Concurrent CRS
Grade 1	ICE score 7-9 <sup>b</sup> or depressed level of consciousness <sup>c</sup> : awakens spontaneously.	Management of CRS as appropriate per <a href="#">Table 23</a> .	Monitor neurologic symptoms and consider neurology consultation and evaluation, per investigator discretion.
		Monitoring of neurologic symptoms and consider neurology consultation and evaluation, per investigator discretion.	Consider dexamethasone
		Consider non-sedating, anti-seizure medicines (eg, levetiracetam) for seizure prophylaxis.	
Grade 2	ICE score-3-6 <sup>b</sup> or depressed level of consciousness <sup>c</sup> : awakens to voice.	Administer tocilizumab per <a href="#">Table 23</a> for management of CRS.	Administer dexamethasone <sup>d</sup> 10 mg intravenously every 6 hours.
		If no improvement after starting tocilizumab, administer dexamethasone <sup>d</sup> 10 mg intravenously every 6 hours if not already taking other corticosteroids. Continue dexamethasone use until the event is Grade 1 or less, then taper.	Continue dexamethasone use until the event is Grade 1 or less, then taper.
		Consider non-sedating, anti-seizure medicines (eg, levetiracetam) for seizure prophylaxis. Consider neurology consultation and other specialists (ie, intensivists) for further evaluation, as needed.	
Grade 3	ICE score-0-2 <sup>b</sup> or depressed level of consciousness <sup>c</sup> : awakens only to tactile stimulus,  or seizures <sup>c</sup> , either: • any clinical seizure, focal or generalized, that resolves rapidly, or • non-convulsive seizures on EEG that resolve with intervention,  or raised ICP: focal/local edema on neuroimaging <sup>c</sup> .	Administer tocilizumab per <a href="#">Table 23</a> for management of CRS.	Administer dexamethasone <sup>d</sup> 10 mg intravenously every 6 hours.
		In addition, administer dexamethasone <sup>d</sup> 10 mg intravenously with the first dose of tocilizumab and repeat dose every 6 hours. Continue dexamethasone use until the event is Grade 1 or less, then taper.	Continue dexamethasone use until the event is Grade 1 or less, then taper.
		Consider non-sedating, anti-seizure medicines (eg, levetiracetam) for seizure prophylaxis. Consider neurology consultation and other specialists (ie, intensivists) for further evaluation, as needed.	

**Table 24: Guidelines for the Management of CAR-T Cell-related Neurotoxicity (eg, ICANS)**

ICANS Grade	Presenting Symptoms <sup>a</sup>	Concurrent CRS	No Concurrent CRS
Grade 4	ICE score-0 <sup>b</sup> or depressed level of consciousness <sup>c</sup> either: <ul style="list-style-type: none"> <li>• subject is unarousable or requires vigorous or repetitive tactile stimuli to arouse, or</li> <li>• stupor or coma,</li> </ul> or seizures <sup>c</sup> , either: <ul style="list-style-type: none"> <li>• life-threatening prolonged seizure (&gt;5 min), or</li> <li>• repetitive clinical or electrical seizures without return to baseline in between,</li> </ul> or motor findings <sup>c</sup> : <ul style="list-style-type: none"> <li>• deep focal motor weakness such as hemiparesis or paraparesis,</li> </ul> or raised ICP / cerebral edema <sup>c</sup> , with signs/symptoms such as: <ul style="list-style-type: none"> <li>• diffuse cerebral edema on neuroimaging, or</li> <li>• decerebrate or decorticate posturing, or</li> <li>• cranial nerve VI palsy, or</li> <li>• papilledema, or</li> <li>• Cushing's triad.</li> </ul>	Administer tocilizumab per <a href="#">Table 23</a> for management of CRS.  As above or consider administration of methylprednisolone 1000 mg intravenously per day with first dose of tocilizumab and continue methylprednisolone 1000 mg intravenously per day for 2 or more days, per investigator discretion.	As above or consider administration of methylprednisolone 1000 mg intravenously per day for 3 days; if improves, then manage as above.
		Consider non-sedating, anti-seizure medicines (eg, levetiracetam) for seizure prophylaxis. Consider neurology consultation and other specialists (ie, intensivists) for further evaluation, as needed.  In case of raised ICP/cerebral edema, refer to <a href="#">Table 25</a> for additional management guidelines.	

Abbreviations: CRS=cytokine release syndrome; EEG=electroencephalogram; ICANS= Immune Effector Cell-Associated Neurotoxicity Syndrome; ICE= Immune-effector Cell-associated Encephalopathy; ICP= intracranial pressure

<sup>a</sup> Management is determined by the most severe event, not attributable to any other cause

<sup>b</sup> If subject is arousable and able to perform Mental Status assessment, the following domains should be tested: orientation, naming, following commands, writing, and attention (see [Attachment 3](#); ICE-Tool).

<sup>c</sup> Attributable to no other cause.

<sup>d</sup> All references to dexamethasone administration are dexamethasone or equivalent

**Table 25: Guidelines for the Management of Raised ICP / Cerebral Edema<sup>a</sup>**

- Elevate head of patient's bed to an angle of 30 degrees.
- If patient has Ommaya reservoir, drain CSF to target opening pressure of <20 mmHg.
- Hyperventilation to achieve target partial pressure of arterial carbon dioxide (PaCO<sub>2</sub>) of 28–30 mmHg, but maintained for no longer than 24 hours.
- Consider neurology and/or neurosurgery consultation.
- Use high-dose corticosteroids with methylprednisolone IV 1 g/day, as recommended above.
- Hyperosmolar therapy with either mannitol (20 g/dL solution) or hypertonic saline (3% or 23.4%, as detailed below):
  - Mannitol: initial dose 0.5–1 g/kg; maintenance at 0.25–1 g/kg every 6 hours while monitoring metabolic profile and serum osmolality every 6 hours, and withhold mannitol if serum osmolality is  $\geq 320$  mOsm/kg, or the osmolality gap is  $\geq 40$ ,
  - Hypertonic saline: initial 250 mL of 3% hypertonic saline; maintenance at 50–75 ml/hr while monitoring electrolytes every 4 hours, and withhold infusion if serum Na levels reach  $\geq 155$  mEq/L,
  - For patients with imminent herniation: initial 30 mL of 23.4% hypertonic saline; repeat after 15 min, if needed.
- Consider IV anesthetics for burst-suppression pattern on electroencephalography.

a In addition to toxicity management guidelines provided in [Table 24](#): Guidelines for the Management of CAR-T Cell-Related Neurotoxicity (ie, ICANS)

### 6.2.2.2. Other Neurotoxicities

If any neurologic or psychiatric symptoms are noted (see below and Section 1.1.5), the medical monitor should be contacted, and the subject should be referred immediately to a neurologist for a full evaluation. Subjects should be monitored for neurotoxicity for 1-year post JNJ-68284528 infusion. Particular attention should be paid to the appearance of any of the following:

- personality changes
- lack of facial expression
- being less communicative
- difficulty with short term memory
- slowing of movements
- difficulty performing simple tasks
- difficulty swallowing
- joint or muscle stiffness or inflexibility
- shuffling gait
- restlessness
- micrographia

Early detection, workup and intervention, may be important to prevent neurologic toxicity from worsening. The following is a list of potential diagnostics that should be considered in subjects with new neurologic symptoms:



- Positron emission tomography/computerized tomography (PET/CT) of the brain and/or brain MRI with perfusion and an electroencephalogram (EEG).
- Lumbar puncture to rule out infection (in particular John Cunningham virus [JCV], herpes zoster virus [HZV], herpes simplex virus [HSV]-1/2, HHV-6, HHV-7, Epstein-Barr virus [EBV], cytomegalovirus [CMV]).
- Serologic testing for HHV-6 and HHV-7 by PCR for viremia.
- CSF flow cytometry and cytology should be considered to rule out leptomeningeal disease.
- Cerebral spinal fluid (CSF) analysis to rule out paraneoplastic syndromes.
- Thiamine level (consider empiric thiamine replacement while awaiting results) ([MD Anderson 2019](#))

Therapy directed at reduction or elimination of CAR-T cells, including chemotherapy, may be considered in consultation with the sponsor for subjects who develop neurotoxicity that remains unresponsive to other interventions.

Per Section 9.5 of the protocol, if cerebral spinal fluid (CSF) or other relevant biological sample analysis is clinically indicated, a sample of CSF will be requested for additional analysis by the sponsor.

### **6.2.3. Tumor Lysis Syndrome**

Although TLS is uncommon in subjects with multiple myeloma, one subject in the Phase 1 Legend-2 Study experienced fatal TLS. Subjects must be monitored closely for symptoms of TLS. Management of TLS, including hyperkalemia, hyperuricemia, hyperphosphatemia, and hypocalcemia, is highly recommended. It is also required that high-risk subjects, ie, those with a high tumor burden ( $\geq 60\%$  plasma cell infiltrate on the bone marrow biopsy or aspirate [whichever is higher] or a subject with multiple extramedullary disease sites or plasmacytomas), be treated prophylactically in accordance with local standards (eg, extra hydration; diuretics; allopurinol 300 mg daily and primary or secondary uricosuric agents, as indicated). TLS will be captured as an adverse event of special interest (see Section 12.3.3).

### **6.2.4. Second Primary Malignancy**

Second primary malignancy is a possibility due to the risk of viral insertion (DNA integration) of the lentiviral vector. Second primary malignancies should be managed per institutional standards. Second primary malignancies must be reported during the duration of the study, irrespective of when they occur, and subsequently will be collected in a long-term follow-up study yearly until 15 years post dosing of JNJ-68284528. A tumor sample should be collected and DNA, RNA, or protein analysis may be performed to investigate the presence of lentiviral elements.

### **6.2.5. Cytopenia**

Subjects may exhibit cytopenias for several weeks following lymphodepleting chemotherapy and JNJ-68284528 infusion. Frequent monitoring of hematological parameters and provide supportive care (eg, irradiated blood and thrombocyte concentrates, granulocyte-colony stimulating factor for neutropenia) as outlined by institutional guidelines. Pegylated myeloid growth factors



(ie, pegfilgrastim) are prohibited. Prolonged neutropenia may increase the risk of infection. Severe thrombocytopenia may increase the risk of bleeding. Blood counts should be monitored after JNJ-68284528 infusion. The use of myeloid growth factors, particularly granulocyte colony-stimulating factor (G-CSF), should be avoided during CRS.

Parvovirus B19 monitoring by PCR should be considered in subjects experiencing prolonged neutropenia or a decline in neutrophil counts following recovery.

For subjects in Cohort D and Cohort E, initiating lenalidomide after JNJ-68284528 may cause significant neutropenia and thrombocytopenia. For subjects in Cohort E, daratumumab may also increase the risk of cytopenia. Monitor complete blood counts (CBC) as specified in the Time and Events Schedule ([Table 3](#) and [Table 6](#)). A dose interruption and reduction may be required (refer to local prescription information). Guidance is provided in [Table 12](#), [Table 13](#), and [Table 14](#). Subjects should be monitored frequently for infection and bleeding. Supportive care should be provided per institutional standards.

### **6.2.6. Hypogammaglobulinemia**

Hypogammaglobulinemia may occur in subjects receiving JNJ-68284528. Monitor immunoglobulin levels after treatment as detailed in the Time and Events Schedule (see [Table 1](#), [Table 3](#), and [Table 6](#)) and more frequently if clinically indicated and treat according to local guidelines, including administration of immunoglobulin replacement and monitoring for infection. Vaccination with live virus vaccines is not permitted for at least 4 weeks prior to the start of the conditioning regimen and for 100 days after infusion of JNJ-68284528.

Subjects with IgG <400 mg/dL and recurrent infections may receive prophylactic IVIG as per institutional guidelines.

### **6.2.7. Infections**

Do not administer JNJ-68284528 to subjects with active infection. Administration of JNJ-68284528 may increase the risk of infection due to cytopenias or hypogammaglobulinemia. Subjects should be monitored frequently for infection and should have blood cultures obtained and empiric antibiotics administered per institutional standards. Immunocompromised patients are at risk for opportunistic infections; prophylactic use of antibiotics, antivirals, or antifungals should be considered. Extended use of anti-microbial therapies for at least 6 month (or longer per institutional guidelines) or consistent with post ASCT consensus guidelines after JNJ-68284528 dosing are recommended (see [Attachment 18](#)). Perform screening for hepatitis B virus (HBV), hepatitis B virus (HCV), and human immunodeficiency virus (HIV) and monitor as clinically indicated (see HBV monitoring recommendations in [Section 9.7](#) and [Attachment 10](#)), and initiate treatment as appropriate. HBV reactivation, in some cases resulting in fulminant hepatitis, hepatic failure and death, may occur in subjects treated with drugs directed against B cells such as JNJ-68284528. HBV reactivation has occurred in subjects receiving other CAR-T products who appear to have resolved hepatitis B infection. Routinely monitor HBV DNA and AST/ALT for subjects with risk of HBV reactivation.

## **6.2.8. Hypersensitivity Reactions**

Allergic reactions may occur with the infusion of JNJ-68284528. Serious hypersensitivity reactions including anaphylaxis, may be due to dimethyl sulfoxide (DMSO), dextran 40, or residual ampicillin or kanamycin in JNJ-68284528. Subjects should be treated urgently per institutional standards, avoiding corticosteroid use if possible. Subjects should receive premedication prior to JNJ-68284528 dosing as noted in Section 6.1.3.3.

For subjects in Cohort D and Cohort E please refer to the prescribing information for additional risks associated with lenalidomide.

## **6.2.9. Cohort E: Management of Infusion-related Reactions and Local Injection-site Reactions of Daratumumab SC**

### **6.2.9.1. Infusion-related Reactions**

Infusion-related reactions (IRRs) are systemic reactions related to daratumumab administration. Subjects should be observed carefully during daratumumab administrations. Trained study staff at the clinic should be prepared to intervene in case of any IRRs, and resources necessary for resuscitation (eg, agents such as epinephrine and aerosolized bronchodilator, medical equipment such as oxygen tanks, tracheostomy equipment, and a defibrillator) must be available at the bedside. Attention to staffing should be considered when multiple participants will be dosed at the same time.

If an IRR develops during daratumumab SC administration, then the administration should be temporarily interrupted. Participants who experience adverse events during daratumumab-SC administration must be treated for their symptoms. Participants should be treated with paracetamol (acetaminophen), antihistamine, or corticosteroids, as needed. Intravenous saline may be indicated. For bronchospasm, urticaria, or dyspnea, participants may require antihistamines, oxygen, corticosteroids, or bronchodilators. For hypotension, participants may require vasopressors. In the event of a life-threatening IRR (which may include pulmonary or cardiac events) or an anaphylactic reaction, daratumumab-SC should be discontinued.

### **Infusion-related Reactions of Grade 1 or Grade 2**

If the investigator assesses a Grade 1-2 IRR to be related to administration of study intervention, then the daratumumab-SC administration should be interrupted. When the participant's condition is stable, daratumumab-SC administration may be restarted at the investigator's discretion. Refer to the SIPPM for further details regarding continuation of daratumumab-SC administration.

If the participant experiences a Grade 2 or higher event of laryngeal edema, or a Grade 2 or higher event of bronchospasm that does not respond to systemic therapy and does not resolve within 6 hours from onset, then the participant must be permanently discontinued from daratumumab-SC treatment.

### **Infusion-related Reactions of Grade 3 or Higher**

For IRR adverse events (other than laryngeal edema or bronchospasm) that are Grade 3, the daratumumab-SC administration must be stopped, and the participant must be observed carefully until resolution of the adverse event or until the intensity of the event decreases to Grade 1, at which

point the daratumumab-SC administration may be restarted at the investigator's discretion. Refer to the SIPPMM for further details regarding continuation of daratumumab-SC administration.

If the intensity of the adverse event returns to Grade 3 after restart of the daratumumab-SC administration, then the participant must be permanently discontinued from daratumumab-SC treatment.

For IRR adverse events that are Grade 4, the daratumumab-SC administration must be stopped, and the participant permanently discontinued from daratumumab-SC treatment.

### **Recurrent Infusion-related Reactions**

If a Grade 3 IRR (or Grade 2 or higher event of laryngeal edema, or a Grade 2 or higher event of bronchospasm) recurs during or within 24 hours after a subsequent daratumumab-SC administration, the participant must be permanently discontinued from daratumumab-SC treatment.

#### **6.2.9.2. Injection-site Reactions**

In clinical studies, SC administration of daratumumab was associated with local injection site reactions, such as induration and erythema, in some subjects. The reactions usually resolved within 60 minutes. Local injection-site reactions should be managed per institutional standards.

## **7. TREATMENT COMPLIANCE**

Apheresis, administration of the conditioning regimen, and infusion of JNJ-68284528 will be done in the controlled environment of a qualified clinical site, under the direct observation of qualified study-site personnel. Subjects will be asked to return containers of lenalidomide at each study visit. Pill counts will be used to assess compliance. Alternative, site specific methods to ensure compliance with outpatient medication administration is permissible. Additional details are provided in the SIPPMM. The details of administration will be recorded in the eCRF (including date, dose of cells, start, and stop times of the IV infusion, and volume infused). Precautions associated with the use of the study treatment and concomitant medications will be reviewed by the sponsor.

Refer to the SIPPMM for a description of the chain of identity and chain of custody procedures associated with the apheresis product and JNJ-68284528.

Components of the D-VRd and D+R regimen will be administered or prescribed by qualified site staff, and the details of each administration will be recorded in the eCRF. Additional details are provided in the Non-JNJ-68284528 SIPPMM.

## **8. PRESTUDY AND CONCOMITANT THERAPY**

Throughout the study, investigators may prescribe concomitant medications or treatments (except for those listed in Section 8.3) deemed necessary to provide adequate supportive care. Medications (including prescription and over-the-counter products, and transfusions of blood products) different from the study treatment must be recorded throughout the study beginning with the signing of the ICF until 100 days after infusion of JNJ-68284528 (Cohorts A, B, and C), 30 days after the last dose of lenalidomide (Cohort D) or until the start of subsequent systemic anticancer

treatment, if earlier, and 30 days after the last dose of D+R (Cohort E) or until the start of subsequent systemic anticancer treatment, if earlier. All concomitant medications will be recorded during screening. Thereafter, selected concomitant medications will be reported. Selected concomitant medications consist of any medication given for an adverse event or serious adverse event, therapeutically or prophylactically, including, but not limited to:

- Anti-cytokine or anti-cytokine receptor therapies
- Anti-seizure medications
- Any medication given for prophylaxis or treatment of TLS
- Any medication given for prevention or treatment of thromboembolic events
- Corticosteroids (including prophylactically for blood product administrations, physiologic replacement doses, high or stress doses, etc.)
- Immunosuppressive agents
- Vaccinations
- Vasopressors and cardiac inotropic agents (For dose, record only maximum daily rate)
- Blood products
- Growth factors
- Systemic antimicrobials – given for prophylaxis or treatment
- Chemotherapy (given for any CAR-T cell related toxicity)

Other:

- Bisphosphonates
- Immunoglobulin therapy
- Medications listed as prohibited in the protocol
- Palliative radiation
- Pain medication
- Any treatment given for SPMs
- Any changes in doses from baseline or newly added concomitant medications to treat new or prior known co-morbidities

Recorded information will include a description of the type of the drug, dosing regimen, route of administration, duration of treatment, and its indication. Medications, including details of previous anticancer treatment, should be documented in the appropriate section of the eCRF.

Anti-myeloma therapy is permitted during bridging therapy (see Section 3.1).

The sponsor must be notified in advance (or as soon as possible thereafter) of any instances in which prohibited therapies are administered.

## 8.1. Prophylaxis for Herpes Zoster Reactivation

For subjects at risk for viral reactivation (all cohorts), prophylaxis for herpes zoster reactivation is strongly recommended. Acceptable antiviral therapy includes acyclovir (eg, 400 mg given orally 3 times a day, or 800 mg given orally 2 times a day or per institutional standards), famciclovir (eg, 125 mg given orally, twice a day or per institutional standards), or valacyclovir (eg, 500 mg given orally, twice a day or per institutional standards), initiated within 1 week after the start of study drug.

## 8.2. Permitted Medications

The following are examples of supportive therapies that may be used during the study:

- Standard supportive care therapies (antiemetics, antidiarrheals, anticholinergics, antispasmodics, antipyretics, antihistamines, analgesics, antibiotics and other antimicrobials, histamine receptor [H<sub>2</sub>] antagonists or proton pump inhibitors, and other medications intended to treat symptoms or signs of disease) and therapies intended to treat CAR-T cell related toxicity (ie, CRS) as clinically indicated, according to institutional standards and as deemed necessary by the investigator.
- Bisphosphonates may be initiated (if not already being administered) unless contraindicated within 1 week prior to the first dose of study treatment and continued until disease progression is established. In the case of severe adverse events such as hypercalcemia, bisphosphonates may be administered as clinically indicated, according to institutional standards and as deemed necessary by the investigator.
- Hematopoietic growth factor support and transfusions (irradiated blood products) are permitted to treat symptoms or signs of neutropenia, anemia or thrombocytopenia according to local standards of care. Non-pegylated myeloid growth factors are permitted up to 1 day prior to the start of the conditioning regimen (Section 6.1.2). For subjects in Cohort D and Cohort E, refer to the lenalidomide prescribing information for precautionary language for agents that may increase the risk of thrombosis.
- Documented infectious complications should be treated with oral or IV antibiotics or other anti-infective agents as considered appropriate by the treating investigator, according to standard institutional practice.
- Chemotherapy agents used to treat CAR-T cell related toxicity are permitted upon consultation with the sponsor (see Section 6.2).
- For subjects in Cohort D and Cohort E, see prescribing information for lenalidomide for additional supportive therapies, including consideration for thromboembolic prophylaxis depending on the subject's risk factors.

## 8.3. Prohibited Therapies

The following medications are prohibited during the study in addition to those therapies identified in Section 6.1.1 (prior to apheresis) and in Section 6.1.2.1 (prior to conditioning regimen). The sponsor must be notified in advance (or as soon as possible thereafter) of any instances in which prohibited therapies are (to be) administered

- Systemic corticosteroid use should be avoided, except for the treatment of CRS or CAR-T cell-related neurotoxicity (eg, ICANS), as described in Table 23 and Table 24. Alternative therapies, if feasible, should be given prior to corticosteroids. No cumulative dose of corticosteroids equivalent to  $\geq 70$  mg prednisone should be administered within the 7 days prior to apheresis (Section 4.1.2 [Cohort A], Section 4.2.2 [Cohort B], Section 4.3.2 [Cohort C], Section 4.4.2 [Cohort D] or within the 7 days prior to the conditioning regimen dosing (Section 6.1.2.1 [all Cohorts]).
- Any chemotherapy, anticancer immunotherapy (other than JNJ-68284528) including any BCMA-directed therapy, or experimental therapy not specified by the protocol (ie, except as described in Section 3.1 [bridging therapy] or protocol specific therapies which may be used in conjunction with JNJ-68284528).
- The use of RANK ligand inhibitors such as denosumab is prohibited due to their potential impact on immune function.
- While in follow-up, emergency orthopedic surgery or radiotherapy is generally prohibited, but may be allowed in the absence of disease progression. Cases must be discussed and approved by the sponsor. Such emergency radiotherapy may consist of localized radiotherapy for pain control or for stabilization of an extensive bone lesion at high risk of pathologic fracture or damage to surrounding tissues.
- Nonsteroidal anti-inflammatory agents should be avoided to minimize the risk of exacerbation of potential sub-clinical myeloma-related kidney disease. Based on the investigator's clinical judgement, low-dose aspirin may be continued for thromboprophylaxis (recommended for Cohort D). For subjects in Cohort D and Cohort E, refer to the lenalidomide prescribing information for recommendations for subjects with prior history of thrombosis.
- Other immunosuppressant agents unless used as protocol-specified pre- or post-treatment medications to treat an adverse event (eg, CRS).
- Vaccination with live, attenuated vaccine after signing consent and in the  $\leq 4$  weeks prior to the infusion of JNJ-68284528, and for 100 days after infusion of JNJ-68284528.
- The use of IV contrast infusions should be avoided to prevent myeloma-related kidney disease. If administration of IV contrast is necessary, then adequate precautions including hydration are indicated.
- Pegylated myeloid growth factors (ie, pegfilgrastim) are prohibited within the first 100 days after infusion of JNJ-68284528. For subjects in Cohort D, refer to the lenalidomide prescribing information for precautions with use of erythropoietic agents.

### 8.3.1. Additional Prohibited Therapies for Cohort E

Concomitant administration of strong CYP3A4 inducers is prohibited with the use of bortezomib. Administration of strong CYP3A4 inhibitors (eg, ketoconazole, ritonavir) should be avoided and is not recommended in subjects receiving bortezomib. If a strong CYP3A4 inhibitor must be given in combination with bortezomib, monitor subjects for signs of bortezomib toxicity and consider a bortezomib dose reduction. For an ongoing list of CYP3A inhibitors and inducers, please refer to <https://www.fda.gov/Drugs/DevelopmentApprovalProcess/DevelopmentResources/DrugInteractionsLabeling/ucm093664.htm>.



## 8.4. Subsequent Anticancer Therapy

Subsequent anticancer therapy administered after JNJ-68284528 (all cohorts) and after lenalidomide (Cohort D) or D+R (Cohort E), should be only administered after sponsor-confirmed PD per IMWG criteria and recorded in the eCRF. The type of high dose chemotherapy (for example, melphalan) should be documented in the eCRF as a subsequent therapy. The start and end date and best response should be documented in the eCRF, if available.

## 9. STUDY EVALUATIONS

### 9.1. Study Procedures

#### 9.1.1. Overview

The Time and Events Schedule summarizes the frequency and timing of procedures and assessments applicable to this study ([Table 1](#) to [Table 8](#)).

All planned assessments, including laboratory tests, on the day of JNJ-68284528 dosing must be completed and the results reviewed prior to the start of the infusion. Treatment decisions will be based on safety assessments performed at the local laboratory and disease assessments performed at the central laboratory.

If multiple assessments are scheduled for the same timepoint, it is recommended that procedures be performed in the following sequence: PRO questionnaires, electrocardiogram (ECG), vital signs, blood draw. Blood collections for biomarkers and pharmacokinetic assessments should be kept as close to the specified time as possible. Actual date and time of assessments will be recorded in the source documents and the laboratory requisition form. Within 24 hours of study treatment infusion, if the study treatment is infused peripherally, blood samples must be drawn from a vein contralateral to the arm into which JNJ-68284528 is infused. If the study treatment is infused via a central vein line, blood samples over the subsequent 24 hours must be drawn from a vein in either arm.

If the subject is unable to complete the PRO assessments, the reason for not completing the questionnaires will be documented. Refer to [Section 9.6](#) for details.

The approximate volume of blood drawn from each subject in Cohorts A, B, and C (up to 2 years' post-JNJ-68284528 treatment) is 924 mL, and for Cohort D is 1110 mL. In addition, women of child-bearing potential will have serum pregnancy tests prior to starting a new cycle of lenalidomide and within 4 weeks after stopping the treatment. This is an additional blood volume of 50 mL over the course of the 2-year treatment period. Repeat or unscheduled samples may be taken for safety reasons or for technical issues with the samples.

#### 9.1.2. Screening Phase

All subjects must sign an ICF prior to the conduct of any study-related procedures. The screening phase begins when the first screening assessment is performed. Screening procedures will be performed up to 28 days before apheresis or start of induction therapy for Cohort E. If an assessment was performed as part of the subject's routine clinical evaluation and not specifically

for this study, it does not need to be repeated after signed informed consent has been obtained provided the assessments fulfill the study requirements and are performed within the specified timeframe prior to the first dose of study treatment. Retesting of abnormal screening values that lead to exclusion are allowed only once during the screening phase (to reassess eligibility). The last result obtained prior to apheresis, or start of induction therapy for Cohort E, will be used to determine eligibility. Subjects who do not meet all inclusion criteria or who meet an exclusion criterion may, at the discretion of the investigator, be rescreened once upon the sponsor's written approval. Subjects who are to be rescreened must sign a new informed consent before rescreening. Rescreening and subsequent activities must be conducted in accordance with protocol defined time windows.

Subjects in Cohort E will receive induction treatment with D-VRd after screening and prior to apheresis as described in Section 6.1.5.

### **9.1.3. Apheresis**

Prior to apheresis, review of safety assessments should be completed per the Time and Events Schedule (Table 1, Table 3, and Table 5). Apheresis should be performed according to institutional standards, with a collection target of  $6 \times 10^9$  PBMCs (range: 2 to  $20 \times 10^9$  PBMCs); two apheresis collections may be performed to attain this target. Instructions for processing and shipping apheresis product are provided in the SIPPM.

### **9.1.4. Cyclophosphamide and Fludarabine Conditioning Regimen**

At the completion of manufacture and quality testing of JNJ-68284528, notification will be sent to the clinical site. Prior to dosing with cyclophosphamide and fludarabine, review of eligibility, safety assessments and disease characteristics should be completed per Section 6.1.2.1. The details regarding safety monitoring and study visits during this phase are included in the Time and Events Schedules (Table 1, Table 3, and Table 5).

A conditioning regimen of cyclophosphamide  $300 \text{ mg/m}^2$  and fludarabine  $30 \text{ mg/m}^2$  daily for 3 days must be administered prior to administration of JNJ-68284528. The dose of fludarabine should be reduced to  $24 \text{ mg/m}^2$  for subjects with an eGFR of 30 to  $70 \text{ mL/min/1.73m}^2$ . Sponsor approval must be obtained to modify the conditioning regimen schedule or dose.

### **9.1.5. JNJ-68284528 Administration**

Administration of JNJ-68284528 is fully described in Table 10.

### **9.1.6. Post-treatment Phase**

The post-treatment phase starts after the completion of JNJ-68284528 infusion and includes the post-infusion period and the post-treatment period.

#### **9.1.6.1. Post-infusion Period**

The post-infusion period starts after the completion of JNJ-68284528 infusion on Day 1 and lasts until Day 100. Any subject who receives an infusion of JNJ-68284528 should continue all subsequent post-infusion assessments as per the Time and Events Schedules (Table 1 to Table 8).



Dependent on the subject's status, medical history, concurrent comorbidities, adequate social support or potential risk factors for developing CAR-T toxicities, including CRS and neurotoxicity, it will be at the Investigator's discretion, subject's willingness, and sponsor approval, whether the subject will be admitted for inpatient monitoring or will receive JNJ-68284528 infusion as an outpatient (See [Table 10](#)).

Subjects will be asked to remain within 1 hour travel time of the hospital and in the company of a competent adult at all times for 1 additional week after hospital discharge, or until Study Day 21, whichever is sooner. Subjects will be provided a "patient ID card" with pertinent information about the study and be asked to carry this card with them for the duration of the post-infusion and post-treatment period.

#### **9.1.6.2. Post-treatment Period (Cohorts A, B, and C)**

The post-treatment period starts on Day 101 and lasts until study completion, defined as 2 years after the last subject in each cohort has received his or her initial dose of JNJ-68284528. Assessments are to be performed per the Time and Events Schedules ([Table 1](#) and [Table 2](#)) and include safety and disease assessments every 28 days up to 12 months then every 56 days thereafter.

After disease progression is documented, survival status and subsequent anticancer therapy will be obtained every 16 weeks until the end of study, unless the subject has died, is lost to follow-up, or has withdrawn consent. PRO assessments will continue to be collected in the post-treatment period even after disease progression is documented or subsequent anticancer therapy is initiated. If the information is obtained via telephone contact, written documentation of the communication must be available for review in the source documents. Survival status will also be collected within 4 weeks prior to the cut off of any planned efficacy analysis, in case a routine survival status collection does not occur within 4 weeks prior to the corresponding cut off. If the subject has died, the date and cause of death will be collected and documented on the eCRF, if or when available. Where allowed by local law, public records may be used to document death or to obtain survival status.

At the investigator discretion and with sponsor approval, study visits (for all cohorts) in the post-treatment part of the study, may be performed remotely via telemedicine technology that connects study subjects to their research coordinators and investigators, if not using the sponsor's telemedicine solution. Blood sample collection may be performed at the subject's home by mobile study personnel (ie, nurses and mobile phlebotomist), if not using the sponsor's home health nursing solution, in the post-treatment period, as early as after Day 100 after JNJ-68284528 infusion.

#### **9.1.6.3. Lenalidomide Treatment Period after JNJ-68284528 Infusion (Cohort D)**

Once introduced, the lenalidomide treatment period starts as early as Day 22 (if required hematological parameters are met) and lasts until study completion, defined as 2 ½ years after the last subject receives their initial dose of JNJ-68284528. Once the lenalidomide treatment is

initiated, subsequent assessment should be adjusted to at the start of each lenalidomide cycle according to Time and Event Schedule. Assessments are to be performed per the Time and Events Schedule (Table 3 and Table 4) and include safety and disease assessments. Safety assessments will be every 28 days for patients receiving lenalidomide. Disease assessments every 28 days for up to 12 months then every 56 days thereafter. Disease evaluations should continue to be performed until confirmed disease progression, death, start of a new anticancer treatment, withdrawal of consent for study participation, or study completion, whichever occurs first. Once disease progression is confirmed, subsequent disease assessment is not required.

After disease progression is documented, survival status and subsequent anticancer therapy will be obtained every 16 weeks until the end of study, unless the subject has died, is lost to follow-up, or has withdrawn consent. PRO assessments will continue to be collected in the post-treatment period even after disease progression is documented or subsequent anticancer therapy is initiated. If the information is obtained via telephone contact, written documentation of the communication must be available for review in the source documents. Survival status will also be collected prior to any planned efficacy analysis, in case a routine survival status collection does not occur within 4 weeks prior to the corresponding cut off. If the subject has died, the date and cause of death will be collected and documented on the eCRF, if or when available. Where allowed by local law, public records may be used to document death or to obtain survival status.

#### **9.1.6.4. D+R Consolidation Treatment Period after JNJ-68284528 Infusion (Cohort E)**

After infusion of JNJ-68284528, subjects in Cohort E will initiate D+R consolidation treatment (Section 6.1.5.2).

The lenalidomide treatment can be initiated a minimum of 21 days post JNJ-68284528 after resolution of cytokine release syndrome (CRS) or neurological toxicities associated with JNJ-68284528. Daratumumab treatment will be initiated approximately 60 to 100 days after JNJ-68284528 and upon adequate recovery from ongoing toxicities. The initiation of daratumumab should begin after at least 1 complete cycle of lenalidomide and coincide with the beginning of the next lenalidomide cycle.

Once lenalidomide treatment is initiated, subsequent safety, efficacy, biomarker, bioanalysis and PRO assessments should be adjusted to the start of the first lenalidomide cycle according to Time and Event Schedule. When possible, initiation of lenalidomide should start on a protocol required visit day to mitigate the need to adjust for subsequent assessments and increased visits at site. Safety assessments will occur every 28 days, at the beginning of each cycle, for subjects receiving lenalidomide or daratumumab, except for the hematology laboratory assessment which will be performed before each dose of daratumumab. Disease assessments will be performed every 28 days and should continue to be performed until confirmed disease progression, death, start of a new anticancer treatment, withdrawal of consent for study participation, or study completion, whichever occurs first.

After disease progression is documented, survival status and subsequent anticancer therapy will be obtained every 16 weeks until the end of study, unless the subject has died, is lost to follow-up, or has withdrawn consent. PRO assessments will continue to be collected in the post-treatment period even after disease progression is documented or subsequent anticancer therapy is initiated. If the information is obtained via telephone contact, written documentation of the communication must be available for review in the source documents. Survival status will also be collected prior to any planned efficacy analysis, in case a routine survival status collection does not occur within 4 weeks prior to the corresponding cut off. If the subject has died, the date and cause of death will be collected and documented on the eCRF, if or when available. Where allowed by local law, public records may be used to document death or to obtain survival status.

#### **9.1.6.5. Long-term Follow-up**

Second primary malignancies will be reported for the duration of the study. Following completion of this study, assessment for replication competent lentivirus (RCL) and second primary malignancies will be collected yearly until 15 years after the last dose with JNJ-68284528 on a follow-up study. In addition, subjects who received retreatment with JNJ-68284528 and are in follow-up at the end of the study will be monitored in this long-term follow-up study. A tumor sample should be collected and DNA, RNA, or protein analysis may be performed to investigate the presence of lentiviral elements.

### **9.2. Efficacy Evaluations**

Disease evaluations must be performed as specified in the Time and Events Schedule. Disease evaluations will be performed by a central laboratory (additional samples may be collected for analysis by the local laboratory) until disease progression, death, start of a new anticancer treatment, withdrawal of consent for study participation, or end of the study, whichever occurs first. If central laboratory data is not available for a particular timepoint, local efficacy data may be substituted. In these situations, the source laboratory data should be interpreted by the study investigator and retained in the site's records. Data will be entered in the eCRF (see Section 9.2.7). However, if a subject receives retreatment with JNJ-68284528 after a confirmed disease progression (see Section 3.1), then disease evaluations will continue according to the Time and Events schedule. This study will use the IMWG-based response criteria (Rajkumar 2011). If it is determined that the study treatment interferes with the immunofixation assay, CR will be defined as the disappearance of the original M-protein associated with multiple myeloma on immunofixation, and the determination of CR will not be affected by unrelated M-proteins secondary to the study treatment (Durie 2015).

For quantitative immunoglobulin (QIg) at baseline, M-protein, immunofixation, and free-light chain (FLC) measurements in serum and 24-hour urine, the investigator will use results provided by the central laboratory. Disease progression must be consistently documented across clinical study sites using the criteria in Attachment 1. The sponsor will use a validated computer algorithm to analyze response to treatment.

Subjects in Cohorts A, B, and C who did not meet the definition of measurable disease per inclusion criterion and were enrolled in the study based on disease that is followed by PET/CT or whole body MRI must also follow the efficacy assessments for all other disease compartments (ie, serum, urine, bone marrow, skeletal [bone] lesions) as described in the Time and Events Schedule.

### **9.2.1. Bone Marrow Examination for MRD Assessment**

MRD will be monitored in subjects using next generation sequencing (NGS) on bone marrow aspirate DNA by central laboratory. Baseline bone marrow aspirates will be used to define the myeloma clones, and post-treatment samples will be used to evaluate MRD negativity at predetermined intervals. Timepoints for the MRD assessment are reflected in the Time and Events schedule ([Table 1](#), [Table 3](#), [Table 5](#) and [Table 6](#)). In case the myeloma clone is not identified successfully from the baseline fresh bone marrow aspirate, the sponsor will ask for non-decalcified diagnostic tissue, such as non-decalcified slides (bone marrow aspirate or clot selection) or formalin-fixed, paraffin-embedded block (clot section only, no bone marrow biopsy).

### **9.2.2. Myeloma Protein Measurements in Serum and Urine**

Blood and 24-hour urine samples for M-protein measurements will be sent to and analyzed by a central laboratory. Only one serum and one 24-hour urine sample per time point are required by the central laboratory to perform the following tests. Assessments will be performed as specified in the Time and Events Schedule ([Table 1](#), [Table 3](#), [Table 5](#), and [Table 6](#)).

- Serum quantitative Ig
- Serum protein electrophoresis (SPEP)
- Serum immunofixation electrophoresis
- Serum FLC assay (for subject in suspected CR/sCR and every disease assessment for subjects with serum FLC only disease)
- 24-hour urine M-protein quantitation by electrophoresis (UPEP)
- Urine immunofixation electrophoresis
- Serum  $\beta$ 2-microglobulin

Blood and 24-hour urine samples will be collected as specified in the Time and Events Schedule ([Table 1](#), [Table 3](#), [Table 5](#), and [Table 6](#)) until the development of confirmed disease progression. Disease progression based on one of the laboratory tests alone must be confirmed by at least 1 repeat investigation. Disease evaluations will continue beyond relapse from CR until disease progression is confirmed. Serum and urine immunofixation and serum free light chain assays will be performed at screening, prior to the start of the conditioning regimen, and thereafter when a CR is suspected (when serum or 24-hour urine M-protein electrophoresis [by SPEP or UPEP] are 0 or non-quantifiable). For subjects with light chain multiple myeloma, serum and urine immunofixation tests will be performed routinely as per the Time and Events Schedule ([Table 1](#), [Table 3](#), [Table 5](#), and [Table 6](#)).

### **9.2.3. Serum Calcium Corrected for Albumin**

Blood samples for calculating serum calcium corrected for albumin will be collected and analyzed centrally until the development of confirmed disease progression. Development of hypercalcemia (corrected serum calcium >11.5 mg/dL [ $>2.9$  mmol/L]) may indicate disease progression or relapse if it is not attributable to any other cause (see disease response criteria in [Attachment 1](#)). Calcium binds to albumin and only the unbound (free) calcium is biologically active; therefore, the serum calcium level must be adjusted for abnormal albumin levels (“corrected serum calcium”). The formula for adjustment is presented in [Attachment 9](#).

### **9.2.4. Bone Marrow Examination**

Bone marrow aspirate or biopsy (acceptable if aspirate is not possible) will be performed for clinical assessments and biomarker evaluations, as indicated in the Time and Events schedule. Clinical staging (morphology, cytogenetics, and immunohistochemistry or immunofluorescence or flow cytometry) should be done by a local laboratory. A portion of the bone marrow aspirate will be sent to the central laboratory for immunophenotyping and to monitor BCMA, checkpoint ligand expression in CD138-positive multiple myeloma cells, and checkpoint expression on T cells. If feasible, bone marrow aspirate also will be performed to confirm CR and sCR and at disease progression. In addition, MRD will be evaluated as specified in Section 9.2.1 and the Time and Events schedules.

### **9.2.5. Skeletal Survey**

A skeletal survey (including skull, entire vertebral column, pelvis, chest, humeri, femora, and any other bones for which the investigator suspects involvement by disease) is to be performed during the screening phase and evaluated by the local laboratory by either roentgenography or low-dose computed tomography (CT) scans without the use of IV contrast. If a CT scan is used it must be of diagnostic quality. Following JNJ-68284528 infusion, and before disease progression is confirmed, X-rays or CT scans should be performed locally, whenever clinically indicated based on symptoms, to document response or progression. Magnetic resonance imaging (MRI) is an acceptable method for evaluation of bone disease and may be included at the discretion of the investigator; however, it does not replace the skeletal survey (see the disease response criteria in [Attachment 1](#)). If a radionuclide bone scan is used at screening, in addition to the complete skeletal survey, then both methods must be used to document disease status. These tests must be performed at the same time. Note: a radionuclide bone scan does not replace a complete skeletal survey.

If a subject presents with disease progression manifested by symptoms of pain due to bone changes, then disease progression may be documented by skeletal survey or other radiographs, depending on the symptoms that the subject experiences. If the diagnosis of disease progression is obvious by radiographic investigations, then no repeat confirmatory X-rays are necessary. If changes are equivocal, then a repeat X-ray is needed in 1 to 3 weeks after initial x-ray.

### **9.2.6. Documentation of Extramedullary Disease/ Extramedullary Plasmacytomas**

Sites of known extramedullary plasmacytomas must be documented  $\leq 14$  days prior to the first dose of the conditioning regimen (Cohorts A, B, and C) or at screening (Cohort D and Cohort E). Clinical examination or MRI may be used to document extramedullary sites of disease. CT scan evaluations are an acceptable alternative if there is no contraindication to the use of IV contrast. Positron emission tomography scan or ultrasound tests are not acceptable to document the size of extramedullary plasmacytomas. However, PET/CT fusion scans can be used to document extramedullary plasmacytomas if the CT component of the PET/CT fusion scan is of sufficient diagnostic quality.

Extramedullary plasmacytomas should be assessed for all subjects with a history of plasmacytomas or if clinically indicated at screening, by clinical examination or radiologic imaging. Assessment of measurable sites of extramedullary disease will be performed, measured, and evaluated locally every 4 weeks (for physical examination) for subjects with a history of plasmacytomas or as clinically indicated during treatment for other subjects until development of confirmed CR or confirmed disease progression. If assessment can only be performed radiologically, then evaluation of extramedullary plasmacytomas may be done every 12 weeks. The methodology used for evaluation of each disease site should be consistent across all visits. Irradiated or excised lesions will be considered not measurable, and will be monitored only for disease progression.

To qualify for VGPR, PR, or MR, the sum of products of the perpendicular diameters of the existing extramedullary plasmacytomas must have decreased by over 90% or at least 50%, respectively, and new plasmacytomas must not have developed (see the disease response criteria in [Attachment 1](#)). To qualify for disease progression, either the sum of products of the perpendicular diameters of the existing extramedullary plasmacytomas must have increased by at least 50%, or the longest diameter of previous lesion  $>1$ cm in short axis must have increased at least 50%, or a new plasmacytoma must have developed. When not all existing extramedullary plasmacytomas are reported, but the sum of products of the perpendicular diameters of the reported plasmacytomas have increased by at least 50%, then the criterion for disease progression is met.

These response criteria are applicable only to patients screened with measurable disease in serum or urine. For subjects in Cohorts A, B, and C without measurable disease in serum or urine, who screened via PET/CT or whole body MRI refer to [Attachment 22](#) for response characterization.

### **9.2.7. Local Laboratory Assessments**

All efforts should be made to collect efficacy data centrally. However, local laboratory data may be collected if central laboratory data are not available at a particular timepoint; this does not include screening assessments. Documentation of the local laboratory data should be sent to the Principal Investigator and filed in the medical record. It is the Principal Investigator's responsibility to ascertain that these results are reviewed and entered into the subject's medical record and the eCRF.



### 9.3. Pharmacokinetics and Immunogenicity

Serum and whole blood samples will be used to evaluate the pharmacokinetics of JNJ-68284528, as well as the immunogenicity of anti-JNJ-68284528 antibodies. Serum collected for pharmacokinetic and immunogenicity analyses may additionally be used to evaluate safety or efficacy aspects that address concerns arising during or after the study period. Genetic analyses will not be performed on these serum samples. Subject confidentiality will be maintained.

#### 9.3.1. Evaluations

Blood and serum samples will be collected for JNJ-68284528 pharmacokinetics, and immunogenicity (antibodies to JNJ-68284528) assessment as specified in the Time and Events Schedule (Table 2, Table 4, Table 7, and Table 8). Also, pharmacokinetic and immunogenicity samples will be collected at the time of onset of suspected CRS or CAR-T cell-related neurotoxicity (eg, ICANS) regardless of causality (specified in Table 2). The exact dates and times of blood sampling must be recorded on the laboratory requisition form. Refer to the Laboratory Manual for sample collection requirements. Collected samples must be stored under specified controlled conditions for the temperatures indicated in the Laboratory Manual.

Venous blood samples will be collected for measurement of CAR-T positive cellular concentration and transgene levels of JNJ-68284528. Bone marrow samples will be collected for measurement of transgene levels and cellular concentrations of JNJ-68284528 (see Time and Events schedule).

Blood samples will be collected for exploratory evaluations of soluble circulating BCMA (sBCMA). This data may be used for mechanistic pharmacokinetic/pharmacodynamic modeling. Subject confidentiality will be maintained. Additional information about the collection, handling, and shipment of biological samples can be found in the Laboratory Manual.

#### 9.3.2. Analytical Procedures

##### Pharmacokinetics

Blood and bone marrow samples will be analyzed to determine CAR-T positive cellular concentration and transgene levels of JNJ-68284528 using specific and sensitive assay methods that are validated by or under the supervision of the sponsor.

##### Immunogenicity

The detection and characterization of antibodies to JNJ-68284528 will be performed using a validated assay method by or under the supervision of the sponsor. Other analyses may be performed to characterize immunogenicity.

#### 9.3.3. Pharmacokinetic Parameters

Blood and bone marrow samples will be collected for the measurement of JNJ-68284528 cellular concentrations and transgene levels for pharmacokinetic analyses (Time and Events Schedule). Pharmacokinetic parameters will be estimated for individuals, and descriptive statistics will be calculated for each cohort. Correlation of  $C_{max}$  and AUC with dose may also be explored.

Pharmacokinetic parameters include, but are not limited to,  $AUC_{inf}$ ,  $AUC_{(0-t)}$ ,  $AUC_{tau}$ ,  $C_{max}$ , half-life, and  $T_{max}$  parameters will be calculated if sufficient data are available for estimation.

### **9.3.4. Immunogenicity Assessments/Antibodies to JNJ-68284528**

Anti-JNJ-68284528 antibodies will be evaluated in serum samples collected from all subjects according to the Time and Events Schedule. Additionally, serum samples should also be collected at the final visit from subjects who discontinued treatment or were withdrawn from the study. These samples will be tested by the sponsor or sponsor's designee.

Serum samples will be screened for antibodies binding to JNJ-68284528 and the titer of confirmed positive samples will be reported. Other analyses may be performed to verify the stability of antibodies to JNJ-68284528 or further characterize the immunogenicity of JNJ-68284528.

### **9.4. Pharmacokinetic/Pharmacodynamic Evaluations**

Pharmacokinetic/pharmacodynamic modeling will be explored to understand and characterize the dose-response relationship.

### **9.5. Biomarkers Evaluations**

Biomarker assessments will focus on several objectives: 1) evaluate apheresis and infused CAR-T cell subsets and activation markers including, but not limited to CD4+, CD8+, CD25+, central memory, effector memory cells; 2) serum or plasma proteomic profiling of cytokines (such as IL-6, IL-15, and IL-10) and other immune related proteins; 3) immunophenotyping of biomarkers of response/resistance on myeloma cells (such as BCMA and PD-L1); 4) determine the clinical benefit (ORR, DOR, TTR, PFS, and OS) of JNJ-68284528 in subjects with cytogenetic modifications (del17p, t(4;14), t(14;16), or other high-risk molecular subtypes) in all cohorts; and 5) immunophenotyping of immune cells subsets such as CD4+ and CD8+ T cells, regulatory T cells, B and NK cells. Additional biomarker samples may be collected to help understand an unexplained adverse event including but not limited to serum or PBMCs from whole blood. Additional sample(s) for cytokines will be collected as clinically indicated (Table 2).

The potential presence of RCL will be evaluated from whole blood samples of subjects treated with JNJ-68284528. RCL will be evaluated using a qPCR assay against the lentiviral vesicular stomatitis virus-G gene for 15 years in the follow-up study.

Peripheral blood mononuclear cells (PBMCs) will be retained for exploratory analysis of the immune system which may include retroviral insertion analysis, T cell receptor (TCR) analysis (both clonality or diversity of TCR), functional in vitro assays, or other. Biomarker analyses are dependent upon the availability of appropriate biomarker assays and clinical response rates. Biomarker analysis may be deferred or not performed, if during or at the end of the study, it becomes clear that the analysis will not have sufficient scientific value for biomarker evaluation, or if there are not enough samples or responders to allow for adequate biomarker evaluation. In the event the study is terminated early or shows poor clinical efficacy, completion of biomarker assessments is based on justification and intended utility of the data.



Based on emerging scientific evidence, the sponsor may request additional material from, including but not limited to, previously collected bone marrow samples, whole blood, bone marrow aspirate or biopsy, or cerebral spinal fluid, or other tissue sample during or after study completion for a retrospective analysis. For subjects diagnosed with a SPM, a tumor sample should be collected (See Section 6.2.4). Additionally, the sponsor will receive a sample of plasmacytoma if patient relapse is suspected. Subjects who have a lumbar puncture as part of their neurologic work up should have cerebral spinal fluid sent for additional tests by the sponsor. In all cases, such analyses would be specific to research related to the study treatment(s) or diseases being investigated. If a subject dies and an autopsy is performed, specimens may be requested by the sponsor for analysis.

### 9.5.1. Pharmacodynamic/Predictive Markers

The baseline of the JNJ-68284528 subsets and dynamic changes/persistence and activation of CAR-positive viable T cells may be associated with the depth and durability of response. An evaluation of these cell populations may be performed by flow cytometry or cytometry by time of flight (CyTOF) or both and correlated with response. Additional immunophenotyping may be performed on bone marrow aspirate and whole blood samples to evaluate expression of biomarkers on myeloma cells (such as BCMA and PD-L1) and immune cell populations (such as CD4<sup>+</sup> and CD8<sup>+</sup> T cells) by flow cytometry or CyTOF. TCR sequencing may be performed to study T cell clonality that may affect drug response. Samples may be characterized by gene expression profiling and somatic mutation analysis by next generation sequencing (whole exome and RNA sequencing) to evaluate potential biomarkers that may correlate with response. Samples may be evaluated by other similar technologies to evaluate protein or RNA expression or for somatic DNA analysis.

Circulating serum biomarkers present following chemotherapy conditioning and following infusion of CAR-T cells have been associated with response to some CAR-T cell-based therapies. Evaluation of cytokines (such as IL-6 and IFN- $\gamma$ ) and other circulating proteins (such as granzymes or perforin) will be analyzed to identify potential pharmacodynamic and predictive biomarkers of response or resistance.

An data review by the DMC will be conducted after the first 5 subjects are treated with JNJ-68284528 in Cohort E to determine if there are any potential pharmacodynamic interactions between D-VRd induction treatment and JNJ-68284528 (see Section 3.1.5).

### 9.6. Patient-Reported Outcome Assessments

The subjects' HRQoL (disease-related symptoms, functioning, and general well-being) will be captured using PRO measures. These measures will be administered according to the Time and Events Schedule (Table 1, Table 3, Table 5, and Table 6) for all cohorts; to be completed by the subjects before any clinical tests, procedures, or other consultations that would influence subject's perceptions of their current health state. The PRO measures will be provided in paper format, in the local language, with the site materials. If a subject requires assistance completing the PRO measures, a study coordinator may assist but should not prompt the subject in selecting their response or provide any interpretation of the questions or response options. The PRO measures

can be interviewer-administered, read verbatim, by telephone during the post-treatment period or completed during the home health visit by the mobile study nurse. When a subject completes the PRO measures, the study coordinator should check that the questionnaires are completed in full, or document why they are missing. Full training documentation will be provided to site coordinators before the start of data collection.

Samples of the PRO measures are provided in [Attachment 11](#):

- EORTC QLQ-C30
- MySim-Q
- PGIS
- PGIC
- PRO-CTCAE

The EORTC-QLQ-C30 version 3 includes 30 items in 5 functional scales (physical, role, emotional, cognitive, and social), 1 global health status scale, 3 symptom scales (pain, fatigue, nausea/vomiting), and 6 single symptom items (dyspnea, insomnia, appetite loss, constipation, diarrhea, and financial difficulties). The recall period is 1 week (“past week”) and responses are reported using a verbal rating scale. The item and scale scores are transformed to a 0 to 100 scale. A higher score represents greater HRQoL, better functioning, and more (worse) symptoms. The EORTC QLQ-C30 has been widely used among patients with multiple myeloma. Reliability, validity, and clinically meaningful change have been demonstrated ([Wisloff 1996](#); [Wisloff 1997](#)).

The Multiple Myeloma Symptom and Impact Questionnaire (MySim-Q) is a disease-specific PRO measure developed to assess symptoms and impacts important to patients with multiple myeloma. It includes 17 items resulting in a total symptom score and a total impacts score. The recall period is the “past 7 days” and responses are reported on a 5-point verbal rating scale. The MySim-Q is an optional assessment in this study.

The Patient Global Impression of Severity (PGIS) is a single item to assess severity of pain. Subjects are asked to rate the severity of their current pain on a 5-point verbal rating scale.

The Patient Global Impression of Change (PGIC) is a single item to assess the subject’s perception in change of their overall health status using a 7-point verbal rating scale. The PGIC is only administered post-infusion.

The National Cancer Institute’s Patient-Reported Outcomes version of the Common Terminology Criteria for Adverse Events (PRO-CTCAE) is an item library of common adverse events experienced by people with cancer that are appropriate for self-reporting and is an optional assessment in this study. Each symptom selected for inclusion can be rated by up to three attributes characterizing the presence/frequency, severity, and/or interference of the adverse event ([Trask 2018](#); [PRO-CTCAE 2019](#)). For subjects with multiple myeloma the following items were selected for inclusion: nausea, vomiting, diarrhea, shortness of breath, rash, dizziness, headache, and fatigue/tiredness/lack of energy. A 5-point verbal rating scale is used for subjects to select

their experience based on the last 7 days. Responses to the PRO-CTCAE will be kept separate from CTCAE data and sites will confirm that the questions are completed but will not have access to the subject's responses for real time review. Additionally, adverse event reporting will not be derived from the PRO data and safety analysis will not be performed using PRO data.

## 9.7. Safety Evaluations

Safety will be measured by adverse events, laboratory test results, vital sign measurements, physical examination findings (including neurologic examination), assessment of ICE-Tool scores, handwriting assessments, assessments of cardiac function, and assessment of ECOG performance status grade. Clinically relevant changes occurring during the study must be recorded on the adverse event section of the eCRF. Any clinically significant abnormalities persisting at the end of the study/early withdrawal will be followed by the investigator until resolution or until a clinically stable condition is reached. Safety monitoring assessments may be performed more frequently, if clinically indicated.

The study will include the following evaluations of safety and tolerability according to the time points provided in the Time and Events Schedule:

### Adverse Events

Adverse events (except for neurotoxicity events, second primary malignancy and HBV reactivation) will be reported by the subject (or, when appropriate, by a caregiver, surrogate, or the subject's legally acceptable representative) from the time a signed and dated informed consent is obtained until 100 days after infusion of JNJ-68284528 (Cohorts A, B, and C), and 100 days after infusion of JNJ-68284528, or 30 days after last dose of lenalidomide, whichever is later (Cohort D), or 30 days after last dose of D+R, whichever is later (Cohort E), or until the start of subsequent anticancer therapy, if earlier. Beyond the adverse event reporting period, adverse events that are considered related to study drug need to be reported until the end of the study. Second primary malignancies will be reported for the duration of the study and, subsequently, will be collected on a long-term follow-up study yearly until 15 years' post dosing of JNJ-68284528. Events of HBV reactivations, and new neurologic adverse events or exacerbation of existing neurologic adverse events will be reported during the first year post-dosing of JNJ-68284528.

Adverse events will be followed by the investigator as specified in Section 12, Adverse Event Reporting and graded according to the National Cancer Institute Common Terminology Criteria for Adverse Events (NCI-CTCAE Version 5.0), with the exception of CRS and CAR-T cell-related neurotoxicity (eg, ICANS). CRS should be evaluated according to the American Society for Blood and Bone Marrow Transplantation (ASBMT) consensus grading (Lee 2019) (Attachment 2). CAR-T cell-related neurotoxicity (eg, ICANS) should be graded using the ASBMT (American Society for Transplantation and Cellular Therapy [ASTCT]) consensus grading (Attachment 4). In addition to capturing ICANS and CRS adverse events (graded by ASTCT consensus grading), all individual symptoms of CRS (eg, fever, hypotension) and ICANS (eg, depressed level of consciousness, seizures) must be captured as individual adverse events and graded by CTCAE criteria. Neurotoxicity that is not temporally associated with CRS, or any other neurologic adverse events that do not qualify as ICANS, will be graded by CTCAE criteria. Events of neurotoxicity or

exacerbation of existing neurologic adverse events will be reported for 1-year post infusion of JNJ-68284528.

Changes in handwriting (ie, micrographia, dysgraphia, or agraphia) should be graded using the criteria outlined in [Attachment 17](#) and reported as an adverse event in the eCRF. Should a subject experience a serious CAR-T associated neurotoxicity (either ICANS or other neurotoxicity), then a copy of the handwriting assessment log should be submitted with the serious adverse event report.

Subjects with Grade 3 or higher toxicity or unresolved adverse events that lead to treatment discontinuation will continue to be assessed until recovery to Grade  $\leq 1$  or baseline, the event is deemed irreversible, the end of the study, or a maximum of 6 months, whichever comes first.

### **Clinical Laboratory Tests**

Blood samples for serum chemistry and hematology will be collected as shown in the Time and Events Schedule ([Table 1](#), [Table 3](#), [Table 5](#), and [Table 6](#)). Disease-related laboratory evaluations are detailed in [Section 9.2](#).

The investigator must review the laboratory results, document this review, and record any clinically relevant changes occurring during the study in the adverse event section of the CRF. Grade 3 or higher laboratory abnormalities should continue to be assessed until recovery to Grade  $\leq 1$  or baseline, the event is deemed irreversible, the end of study, or a maximum of 6 months, whichever comes first. Laboratory certificates or accreditation and normal ranges of the laboratory facility at the site must be submitted to the sponsor before the enrollment of any subject at the site. If the subject has the laboratory assessments conducted at a laboratory facility other than the one associated with the investigational site, the investigator must submit to the sponsor laboratory certificates or accreditation and normal ranges for that facility as well. The laboratory reports must be filed with the source documents.

The following tests will be performed by the local laboratory except for the calcium and albumin-adjusted calcium, which will be performed at the central laboratory:

<b>Hematology</b>	
Hemoglobin	Absolute lymphocyte count
White blood cell count	Platelet count
Absolute neutrophil count	
<b>Coagulation</b>	
Prothrombin time / International normalized ratio	Activated partial thromboplastin time
Fibrinogen	D-dimer
<b>Chemistry</b>	
Sodium	Total bilirubin <sup>a</sup>
Potassium	Alkaline phosphatase
Lactic acid dehydrogenase	Uric acid
Blood urea nitrogen or Urea	Calcium and albumin-adjusted calcium <sup>b</sup>
Creatinine	Phosphate
Glucose	Albumin
AST	Total protein
ALT	Magnesium
Gamma-glutamyl transpeptidase	Creatine phosphokinase (CPK)
Ferritin	C-reactive protein
eGFR <sup>c</sup>	Thyroid Function testing (Cohort D only) <sup>d</sup>
<b>Pregnancy Test</b>	
<b>Pregnancy Test:</b> serum (<5 IU/mL) $\beta$ -hCG	
<b>Tests at Screening only</b>	
<b>Serology:</b>	
- <b>Hepatitis B<sup>e</sup>:</b> HBsAg, anti-HBc, anti-HBs, HBV DNA quantification (for subjects who are anti-HBs positive without a history of vaccination or for subjects who are anti-HBc positive with or without anti-HBs positive) ( <a href="#">Attachment 10</a> )	
- <b>Hepatitis C:</b> HCV antibody, HCV RNA (for subjects who are anti HCV positive)	
- <b>HIV</b>	
<b>Tests at Apheresis only<sup>b</sup></b>	
HIV, Hepatitis B, Hepatitis C, HTLV and other infectious diseases as applicable per local regulations	

Abbreviations: ALT=alanine aminotransferase; AST=aspartate aminotransferase;  $\beta$ -hCG= $\beta$ -human chorionic gonadotropin; HBsAg=hepatitis B surface antigen; anti-HBc=anti-hepatitis B core antibody, anti-HBs=anti-hepatitis B surface antibody; HBV=hepatitis B virus; HCV=hepatitis C virus, HIV=human immunodeficiency virus; HTLV=human T-cell lymphotropic virus.

- Direct bilirubin if Gilbert's disease.
- Performed by central laboratory.
- See [Attachment 10](#) to determine eligibility for enrollment in the study and additional safety monitoring recommendations.
- Thyroid assessment: TSH only, if TSH is <LLN or >ULN need free T3 and free T4 testing
- Calculated using MDRD formula ([Attachment 8](#))

## Electrocardiogram (ECG)

12-lead ECGs will be performed as specified in the Time and Events Schedule ([Table 1](#), [Table 3](#), [Table 5](#), and [Table 6](#)). ECGs should be obtained prior to any other study procedures planned for the same day.

During the collection of ECGs, subjects should be in a quiet setting without distractions (eg, television, cell phones). Subjects should rest in a supine position for at least 5 minutes before ECG collection and should refrain from talking or moving arms or legs. If blood sampling or vital sign measurement is scheduled for the same time point as ECG recording, the procedures should be performed in the following order: ECG(s), vital signs, blood draw.

Additional cardiovascular assessments should be performed as clinically appropriate to ensure subject safety. The clinical investigator will review the results, including ECG morphology, for immediate management. Abnormalities noted at screening should be included in the medical history.

### **Echocardiogram or MUGA scan**

Assessment of cardiac function is required at screening using either echocardiogram or MUGA scan (results obtained  $\leq 8$  weeks before apheresis [All Cohorts] or Day 1 of D-VRd (Cohort E) are acceptable for determining eligibility). At a minimum, this will include assessment of left ventricular ejection fraction (LVEF) reported as a percentage. This value should be recorded in the eCRF. In addition, subjects who receive medications known to cause cardiotoxicity (per locally available prescribing information) in the bridging therapy should have a repeat assessment of cardiac function within 7 days prior to the start of the conditioning regimen.

### **Vital Signs**

Temperature, pulse/heart rate, respiratory rate, blood pressure and oxygen saturation monitoring will be performed as specified in the Time and Events Schedule (Table 1, Table 3, Table 5, and Table 6). Blood pressure and pulse/heart rate measurements should be preceded by at least 5 minutes of rest in a quiet setting without distractions (eg, television, cell phones). Blood pressure and pulse/heart rate measurements will be with a completely automated device, when available.

### **Physical Examination**

The screening physical examination will include, at a minimum, subject's height, general appearance, examination of the skin, ears, nose, throat, lungs, heart, abdomen, extremities, musculoskeletal system, nervous system, and lymphatic system. Thereafter, a symptom-directed physical examination will be conducted as clinically indicated at subsequent timepoints. Abnormalities will be recorded in the appropriate section of the eCRF. Body weight will be measured prior to infusion of JNJ-68284528 (see the Time and Events Schedule [Table 1, Table 3, Table 5, and Table 6]). Clinically significant post-baseline abnormalities should be recorded as adverse events.

### **ECOG Performance Status**

The ECOG performance status scale will be used to grade changes in the subject's daily living activities (Attachment 7) and will be assessed as noted in the Time and Events Schedule (Table 1, Table 3, Table 5, and Table 6).

### **Neurologic Examination**

Magnetic resonance imaging (MRI) at screening or neurology consultation should be considered if pre-existing disease is suspected. For subjects with prior pertinent neurologic disease (eg, stroke, encephalitis) consider baseline MRI of brain and an EEG. At the first sign of neurotoxicity, neurology consultation and evaluation should be considered. CAR-T cell-related neurotoxicity (eg, ICANS) should be graded using ASBMT (ASTCT) consensus grading. Other neurologic adverse events not associated with ICANs should be graded based on CTCAE version 5.0.



Findings from neurologic testing that support CAR-T cell-related neurotoxicity (eg, ICANS) should be reported in the CRF. Submission of neuroimaging scans may also be requested for sponsor review.

### ***Immune-effector Cell-associated Encephalopathy (ICE) Tool Scores***

The ICE test was developed to provide objectivity for the grading of multiple overlapping encephalopathy terms currently included on the approved CAR-T products (Lee 2019) (Attachment 3). The ICE tool will be collected as noted in the Time and Events Schedule (Table 1, Table 3, Table 5, and Table 6) to guide management throughout both phases of the study. It will also be used to grade the severity of ICANS (Attachment 4). All ICE scores must be reported in the eCRF.

### ***Handwriting Assessment***

Qualitative changes in handwriting since baseline are being explored by the sponsor as a potential early clinical predictive marker for neurotoxicity. Currently no standardized CTCAE toxicity gradings are available in the NCI-CTCAE v5.0. for these type of changes in handwriting. Therefore, the sponsor has developed a handwriting assessment criterion to assess subjects for occurrence of the following types of changes in handwriting: micrographia, dysgraphia, or agraphia, as potential early indicators for neurotoxicity (See Attachment 17).

Handwriting assessments will be collected on a writing log according to the Time and Events Schedules and as instructed by the sponsor. Subjects unable to write at baseline are excused from this assessment during study. The sponsor's medical monitor should immediately be notified when changes in handwriting are detected. This will prompt discussion about additional assessments to further evaluate for other neurotoxicity symptoms, further workup, as well as the potential initiation of interventions. All cases of handwriting abnormalities (ie, micrographia, dysgraphia, or agraphia) must be reported as an adverse event in the eCRF. Should a subject experience a serious CAR-T associated neurotoxicity (either ICANS or other neurotoxicity), then a copy of the handwriting assessment log should be submitted with the serious adverse event report. For visits post Day 100 that are completed remotely via Telemedicine, the handwriting assessment may be administered by the mobile study personnel.

### **Ocular Examination**

The ocular examination, for subjects in Cohort C who have received prior ADC, is to include best-corrected visual acuity (BCVA), slit lamp examination (with special focus on cornea), intraocular pressure, and dilated fundoscopic examination.

### **Indirect Antiglobulin Test (IAT) (Cohort E)**

Blood Type, Rh, and Indirect Antiglobulin Test (IAT) should be done before the first dose of daratumumab. Subject RBC phenotyping (standard or extended) is an alternative option to the IAT test, if locally required. Either method must be completed prior to first daratumumab administration.

Daratumumab interferes with the IAT, which is a routine pre-transfusion test performed to identify a patient's antibodies to minor antigens so that suitable donor blood can be given for transfusion. Daratumumab does not interfere with ABO/RhD typing. CD38 is expressed at very low-levels on erythrocytes. Daratumumab binds to the CD38 on erythrocytes, which results in a positive IAT (Indirect Coombs Test). This positive result masks the detection of antibodies to minor antigens and may prevent or delay blood banks from issuing donor blood for transfusion. This effect occurs during daratumumab treatment and for up to 6 months after treatment ends. Subjects will receive a patient identification wallet (study) card for the study that includes the blood profile (ABO, Rh, and IAT or phenotyping) determined before the first injection of daratumumab along with information on the IAT interference for healthcare providers/blood banks. Subjects are to carry this card throughout the treatment period and for at least 2 years after JNJ-68284528 infusion.

Blood banks can eliminate the daratumumab interference with IAT by treating reagent RBCs with dithiothreitol (DTT) ([Chapuy 2015](#); [Chapuy 2016](#)).

Possible methods for blood banks to provide safe RBCs for transfusion to subjects receiving daratumumab include:

- a) Providing ABO/RhD compatible, phenotypically (standard or extended phenotyping) or genotypically matched units
- b) Providing ABO/RhD compatible, K-negative units after ruling out or identifying alloantibodies using DTT-treated reagent RBCs

Uncrossmatched, ABO/RhD compatible RBC units should be administered if transfusion is needed emergently as per local blood bank practice.

Despite daratumumab binding to CD38 on erythrocytes, no indication of clinically significant hemolysis has been observed in daratumumab studies. For additional details, refer to the Daratumumab IB.

## 9.8. Sample Collection and Handling

The actual dates and times of sample collection must be recorded in the CRF or laboratory requisition form. Refer to the Time and Events Schedule ([Table 1](#) to [Table 8](#)) for the timing and frequency of all sample collections.

Instructions for the collection, handling, storage, and shipment of samples are found in the laboratory manual that will be provided. Collection, handling, storage, and shipment of samples must be under the specified, and where applicable, controlled temperature conditions as indicated in the laboratory manual.

## 9.9. Medical Resource Utilization

Health economics data such as medical resource utilization data, associated with medical encounters, will be collected in the CRF by the investigator and study-site personnel for all subjects throughout the study. Protocol-mandated procedures, tests, and encounters are excluded. Per the Time and Events Schedules ([Table 1](#), [Table 3](#), [Table 5](#), and [Table 6](#)), all medical care encounters



since the previous collection will be collected for all subjects. Medical resource evaluation data will be collected until Day 180 ( $\pm 7$  days). Health economics data such as costs associated with the medical encounters will be collected separately from the eCRF. All health economic data will be used only in a de-identified manner.

The data collected may be used to conduct exploratory economic analyses and will include:

- Number and duration of medical care encounters, including surgeries, and other selected procedures (inpatient and outpatient)
- Duration of hospitalization (total days length of stay, including duration by wards; eg, intensive care unit)
- Number and character of diagnostic and therapeutic tests and procedures
- Outpatient medical encounters and treatments (including physician or emergency room visits, tests and procedures, and medications)

## **10. SUBJECT COMPLETION/DISCONTINUATION OF STUDY TREATMENT/ WITHDRAWAL FROM THE STUDY**

### **10.1. Completion**

A subject will be considered to have completed the study if he or she dies before the end of the specific study cohort, has not been lost to follow-up or has not withdrawn consent for study participation before the end of the corresponding study cohort.

### **10.2. Discontinuation of Study Treatment**

A subject should not receive JNJ-68284528, or must be discontinued from lenalidomide treatment (Cohort D), or D-VRd and D+R treatment (Cohort E) if:

- The investigator believes that for safety reasons or tolerability reasons (eg, adverse event) it is in the best interest of the subject to discontinue study treatment
- Grade  $\geq 3$  nonhematologic toxicity related to cyclophosphamide and fludarabine occurs, and precludes retreatment with cyclophosphamide and fludarabine prior to JNJ-68284528 infusion per Section 6.1
- The subject received concurrent (non-protocol) anticancer treatment (with exception of sponsor-approved bridging therapy)
- Confirmed disease progression per IMWG criteria ([Attachment 1](#)) either between the time of conditioning therapy and infusion of JNJ-68284528, or during the post-JNJ-68284528 infusion treatment phase of Cohort D with lenalidomide (as applicable) and Cohort E with daratumumab and lenalidomide. For subjects in Cohorts A, B, and C without measurable disease in serum or urine at screening, progression on either PET/CT scan or Whole Body MRI (see [Attachment 22](#) for imaging response criteria) will be used.
- Subject refuses further study treatment
- Noncompliance with study treatment or procedure requirements

- The subject becomes pregnant prior to infusion, or during the post-JNJ-68284528 infusion lenalidomide treatment phase of Cohort D and Cohort E.

The primary reason for treatment discontinuation will be documented in the eCRF and source documents, and the subject should be followed per standard of care until recovery from bridging therapy or cyclophosphamide and fludarabine conditioning regimen. If a subject's study treatment is discontinued for any reason, this will not result in automatic withdrawal of the subject from the study.

### **10.3. Withdrawal from the Study**

A subject will be withdrawn from the study for any of the following reasons:

- Lost to follow-up
- Withdrawal of consent
- Failure to manufacture JNJ-68284528 after 2 apheresis attempts
- The sponsor discontinues the study

The reason(s) for subject withdrawal will be recorded on the eCRF and source documents. If a subject is lost to follow-up, every reasonable effort must be made by the study site personnel to contact the subject and determine the reason for discontinuation/withdrawal. The measures taken to follow up must be documented.

When a subject withdraws following dosing with JNJ-68284528, the reason for withdrawal is to be documented in the CRF and in the source document. Study assessments for the last visit in the post-infusion period (Time and Events Schedule: Day 100) should be completed prior to withdrawal, if feasible. If the reason for withdrawal from the study is withdrawal of consent then no additional assessments are allowed.

### **10.4. Withdrawal from the Use of Research Samples**

#### **Withdrawal from the Use of Samples in Future Research**

The subject may withdraw consent for use of samples for research (refer to Section 16.2.5, Long-Term Retention of Samples for Additional Future Research). In such a case, samples will be destroyed after they are no longer needed for the clinical study. Details of the sample retention for research are presented in the main ICF.

## **11. STATISTICAL METHODS**

Statistical analysis will be done by the sponsor or under the authority of the sponsor. A general description of the statistical methods to be used to analyze the efficacy and safety data is outlined below. Specific details will be provided in the Statistical Analysis Plan.

Statistical analysis will be performed for each cohort separately.

Continuous variables will be summarized using the number of observations, mean, standard deviation, coefficient of variation, median, and range as appropriate. Categorical values will be summarized using the number of observations and percentages as appropriate.

### 11.1. Subject Information

The analysis populations for this study are defined as follows:

- **Modified Intent-To-Treat (mITT) Analysis Set:**
  - For Cohorts A, B, C, and D: This set consists of subjects who received a JNJ-68284528 infusion at the target dose and will be considered as the primary analysis set for all efficacy summaries.
  - For Cohort E: This set consists of subjects who received at least a dose of D-VRd induction therapy and received a JNJ-68284528 infusion at the target dose, and will be considered as the primary analysis set for all efficacy summaries.
- **All Treated Analysis Set:** This set consists of subjects who received JNJ-68284528 infusion and will be considered as the primary analysis set for safety summaries.
- **Pharmacokinetic Analysis Set:** This set consists of all subjects who received JNJ-68284528 infusion and have at least 1 post-dose pharmacokinetic sample.
- **Immunogenicity Analysis Set:** This set consists of all subjects who received JNJ-68284528 infusion and have at least 1 post-dose immunogenicity sample.

### 11.2. Sample Size Determination

No formal statistical hypothesis testing will be performed. The sample size of each cohort is selected to collect necessary data on preliminary efficacy and safety.

### 11.3. Efficacy Analyses

Endpoint Definitions:

- Minimal residual disease (MRD) negative rate is defined as the proportion of subjects who have negative MRD by bone marrow aspirate at any time point
  - For Cohorts A, B, C, and D: after initial dose of JNJ-68284528 and before disease progression or starting subsequent therapy including retreatment of JNJ-68284528.
  - For Cohort E: after the first dose date of D-VRd and before disease progression or starting subsequent therapy

MRD negative is defined as less than 1 in  $10^5$  residual tumor cells detected in the bone marrow.

- MRD negative rate at 12 months for subjects who achieved a complete response (CR MRD neg 12 month) is defined as the proportion of subjects who are MRD negative by bone marrow aspirate and meet the IMWG criteria for complete response at 12 months (+/- 3 months) or for subjects without measurable disease in Cohorts A, B, and C complete response via imaging criteria ([Attachment 22](#)) at 12 months (+/- 3 months)

- For Cohorts A, B, C, and D: after initial dose of JNJ-68284528 and before disease progression or starting subsequent therapy including retreatment of JNJ-68284528.
- For Cohort E: after the first dose date of D-VRd and before disease progression or starting subsequent therapy.
- Time to MRD negativity will be calculated among subjects who are MRD negative by bone marrow aspirate
  - For Cohorts A, B, C, and D: from the date of the initial infusion of JNJ-68284528 to the initial date of reaching the MRD negative status.
  - For Cohort E: from the first dose date of D-VRd to the initial date of reaching the MRD negative status.

Duration of MRD negativity will be calculated among subjects who are MRD negative by bone marrow aspirate from the date of initial MRD negativity to the date when minimal residual disease is detected at the same threshold.

Overall response rate (ORR) is defined as the proportion of subjects who achieve a PR or better according to the IMWG criteria ([Durie 2006](#); [Durie 2015](#); [Rajkumar 2011](#); [Kumar 2016](#)). For subjects with neither serum nor urine measurable disease and using PET/CT or whole body MRI to satisfy the measurable disease criteria (Cohorts A, B, and C), see [Attachment 22](#) for adjudication of response criteria and [Attachment 23](#) for reporting considerations. Response to treatment will be analyzed by a validated computerized algorithm ([Dimopoulos 2016](#); [Palumbo 2016](#)).

VGPR or better response rate (sCR+CR+VGPR) is defined as the proportion of subjects who achieve a VGPR or better response according to the IMWG criteria ([Durie 2006](#); [Durie 2015](#); [Rajkumar 2011](#); [Kumar 2016](#)).

Duration of response (DOR) will be calculated among responders (with a PR or better response) from the date of initial documentation of a response (PR or better) to the date of first documented evidence of progressive disease, as defined in the IMWG criteria. Relapse from CR by positive immunofixation or trace amount of M-protein is not considered as disease progression. Disease evaluations will continue beyond relapse from CR until disease progression is confirmed. For subjects who have not progressed, data will be censored at the last disease evaluation before the start of any subsequent anti-myeloma therapy.

- Time to response (TTR) is defined as
  - For Cohorts A, B, C, and D: from the date of the initial infusion of JNJ-68284528 and the first efficacy evaluation that the subject has met all criteria for PR or better.
  - For Cohort E: from the first dose date of D-VRd to the first efficacy evaluation that the subject has met all criteria for PR or better.

Progression-free survival (PFS) is defined as the time from the date of the initial infusion of JNJ-68284528 for Cohorts A, B, C, and D, or the date of first dose of D-VRd for Cohort E to the date of first documented disease progression, as defined in the IMWG criteria, or death due to any cause, whichever occurs first. For subjects in Cohorts A, B, and C without measurable disease in

serum or urine at screening, PFS is defined as the time from the date of the initial infusion of JNJ-68284528 to the date of disease progression as identified by imaging response criteria in [Attachment 22](#) or death due to any cause, whichever occurs first. For subjects who have not progressed and are alive, data will be censored at the last disease evaluation before the start of any subsequent anti-myeloma therapy.

Overall survival (OS) is measured from the date of the initial infusion of JNJ-68284528 for Cohorts A, B, C, and D, or the date of first dose of D-VRd for Cohort E to the date of the subject's death. If the subject is alive or the vital status is unknown, then the subject's data will be censored at the date the subject was last known to be alive.

The primary analysis for the primary endpoint, the overall MRD negative rate, will be conducted approximately at 1 year after the last subject for each individual cohort has received his or her initial dose of JNJ-68284528, and will be based on the mITT analysis set. The analysis will be performed separately for each cohort. The MRD negative rate and its 2-sided 95% Clopper-Pearson exact CI will be presented. A sensitivity analysis of overall MRD negative rate will be performed based on the subjects in the mITT analysis set who received the JNJ-68284528 product that met all of the pre-specified release criteria.

The efficacy analyses will be provided for each cohort separately.

For time-to-event endpoints, such as DOR, PFS, and OS, the distributions will be provided using Kaplan-Meier estimates. Detailed analysis methods will be provided in the Statistical Analysis Plan.

#### **11.4. Pharmacokinetic Analyses**

All concentrations below the lowest quantifiable concentration or missing data will be labeled as such in the concentration database. Concentrations below the lowest quantifiable concentration will be treated as zero in the summary statistics. Descriptive statistics will be used to summarize CAR-T positive cell count and transgene level at each sampling timepoint. By dose level, individual pharmacokinetic parameters, including descriptive statistics ( $C_{max}$ ,  $t_{max}$ ,  $t_{last}$ ,  $AUC_{last}$ ,  $AUC_{\infty}$ ,  $\lambda_z$ ,  $t_{1/2}$ ) will be summarized in tables and listings.

If sufficient data are available, then other pharmacokinetic/pharmacodynamic modeling may be performed, including exploring the relationship between JNJ-68284528 cellular concentrations, transgene levels, pharmacodynamic markers (eg, sBCMA, M-protein) and endpoints of clinical efficacy and safety. If performed, details and results of the analysis will be presented in a separate report.

#### **11.5. Immunogenicity Analyses**

The incidence of anti-JNJ-68284528 antibodies will be summarized for all subjects who receive JNJ-68284528 and have appropriate samples for detection of antibodies to JNJ-68284528 (ie, subjects with at least 1 sample obtained after infusion of JNJ-68284528). The results will be

summarized by cohort for subjects with appropriate samples for the detection of antibodies to JNJ-68284528.

Immunogenicity analyses will be descriptive in nature and will include the number and percentage of subjects who developed anti-JNJ-68284528 antibodies. The effect of anti-JNJ-68284528 antibodies on pharmacokinetics, safety, and efficacy may also be evaluated.

### **11.6. Pharmacokinetic/Pharmacodynamic Analyses**

If sufficient data are available, then other pharmacokinetic/pharmacodynamic modeling may be performed, including exploring the relationship between JNJ-68284528 cellular concentrations, transgene levels, pharmacodynamic markers (eg, sBCMA, M-protein) and endpoints of clinical efficacy and safety. If performed, details and results of the analysis will be presented in a separate report.

### **11.7. Safety Analyses**

All safety analyses are to be performed on data from the all treated analysis set. The baseline value for safety assessment is defined as the value collected at the time closest to, but prior to, the start of JNJ-68284528 infusion for Cohorts A, B, C, and D or the first dose of D-VRd for Cohort E. The safety parameters to be evaluated are the incidence, severity, and type of adverse events, clinically significant changes in the subject's physical examination findings, vital signs measurements, and clinical laboratory results. Exposure to investigational product and reasons for discontinuation of study treatment will be tabulated. Adverse events will be summarized by system organ class, preferred term, worst grade experienced by the subject, and by cohort.

#### **Adverse Events**

The verbatim terms used in the CRF by investigators to identify adverse events will be coded using the Medical Dictionary for Regulatory Activities (MedDRA). Treatment-emergent adverse events are adverse events with onset after JNJ-68284528 infusion for Cohorts A, B, C, and D or after the first dose of D-VRd for Cohort E or that are a consequence of a pre-existing condition that has worsened since baseline. All reported treatment-emergent adverse events will be included in the analysis. For each adverse event, the percentage of subjects who experience at least 1 occurrence of the given event will be summarized by cohort. In addition, comparisons between cohorts will be provided if appropriate.

Summaries, listings, datasets, or subject narratives may be provided, as appropriate, for those subjects who die, who experience an adverse event of special interest, who discontinue study treatment due to an adverse event, or who experience a severe or a serious adverse event.

Adverse events that occur after administration of the conditioning regimen and before JNJ-68284528 infusion will be summarized and listed separately.

#### **Clinical Laboratory Tests**

Laboratory data will be summarized by type of laboratory test. Descriptive statistics will be calculated for each laboratory analyte at baseline and for observed values and changes from

baseline at each scheduled time point. Worst toxicity grade during treatment will be presented according to NCI-CTCAE Version 5.0. Change from baseline to the worst toxicity grade experienced by the subject during the study will be provided as shift tables.

### **Electrocardiogram (ECG)**

The interpretation of the ECGs as determined by a qualified physician (investigator or qualified designee) will be summarized at scheduled time points.

### **Vital Signs**

Descriptive statistics of temperature, pulse/heart rate, respiratory rate, and blood pressure (systolic and diastolic) values and changes from baseline will be summarized at each scheduled time point. The percentage of subjects with values beyond clinically important limits will be summarized.

## **11.8. Patient-reported Outcome Assessments**

The EORTC QLQ-C30, MySiM-Q, PGIC, and PGIS will be descriptively summarized at each time point. The proportion of subjects exceeding meaningful change thresholds will be evaluated to assess individual change from baseline ([EORTC Quality of Life 2018](#)). Frequency distributions of the PRO-CTCAE items by visit will be reported.

## **12. ADVERSE EVENT REPORTING**

Timely, accurate, and complete reporting and analysis of safety information from clinical studies are crucial for the protection of subjects, investigators, and the sponsor, and are mandated by regulatory agencies worldwide. The sponsor has established Standard Operating Procedures in conformity with regulatory requirements worldwide to ensure appropriate reporting of safety information; all clinical studies conducted by the sponsor or its affiliates will be conducted in accordance with those procedures.

### **12.1. Definitions**

#### **12.1.1. Adverse Event Definitions and Classifications**

##### **Adverse Event**

An adverse event is any untoward medical occurrence in a clinical study subject administered a medicinal (investigational or non-investigational) product. An adverse event does not necessarily have a causal relationship with the treatment. An adverse event can therefore be any unfavorable and unintended sign (including an abnormal finding), symptom, or disease temporally associated with the use of a medicinal (investigational or non-investigational) product, whether or not related to that medicinal (investigational or non-investigational) product. (Definition per International Conference on Harmonisation [ICH])

This includes any occurrence that is new in onset or aggravated in severity or frequency from the baseline condition, or abnormal results of diagnostic procedures, including laboratory test abnormalities.



Note: The sponsor collects adverse events starting with the signing of the ICF (refer to Section 12.3.1, All Adverse Events, for time of last adverse event recording).

### **Serious Adverse Event**

A serious adverse event based on ICH and EU Guidelines on Pharmacovigilance for Medicinal Products for Human Use is any untoward medical occurrence that at any dose:

- Results in death
- Is life-threatening  
(The subject was at risk of death at the time of the event. It does not refer to an event that hypothetically might have caused death if it were more severe.)
- Requires inpatient hospitalization that was not required by the protocol or prolongation of existing hospitalization
- Results in persistent or significant disability/incapacity
- Is a congenital anomaly/birth defect
- Is a suspected transmission of any infectious agent via a medicinal product
- Is Medically Important\*

\*Medical and scientific judgment should be exercised in deciding whether expedited reporting is also appropriate in other situations, such as important medical events that may not be immediately life threatening or result in death or hospitalization but may jeopardize the subject or may require intervention to prevent one of the other outcomes listed in the definition above. These should usually be considered serious.

If a serious and unexpected adverse event occurs for which there is evidence suggesting a causal relationship between the study treatment and the event (eg, death from anaphylaxis), the event must be reported as a serious and unexpected suspected adverse reaction even if it is a component of the study endpoint (eg, all-cause mortality).

### **Unlisted (Unexpected) Adverse Event/Reference Safety Information**

An adverse event is considered unlisted if the nature or severity is not consistent with the applicable product reference safety information. For JNJ-68284528 and daratumumab the expectedness of an adverse event will be determined by whether or not it is listed in the Investigator's Brochure. For dexamethasone, bortezomib, and lenalidomide, with a marketing authorization, the expectedness of an adverse event will be determined by whether or not it is listed in the respective USPI or SmPC.

### **Adverse Event Associated with the Use of the Drug**

An adverse event is considered associated with the use of the drug if the attribution is related by the definitions listed in Section 12.1.2, Attribution Definitions.



### 12.1.2. Attribution Definitions

#### Not Related

An adverse event that is not related to the use of the drug.

#### Related

An adverse event that might be due to the use of the drug.

### 12.1.3. Severity Criteria

An assessment of severity grade will be made by the investigator according to the NCI CTCAE Version 5.0, with the exception of CRS and CAR-T cell-related neurotoxicity (eg, ICANS). CRS should be evaluated according to the ASBMT (ASTCT) consensus grading (Lee 2019) (Attachment 2). CAR-T cell-related neurotoxicity (eg, ICANS) should be graded using the ASBMT consensus grading (Attachment 4). In addition to capturing ICANS and CRS adverse events (graded by ASTCT consensus grading), all individual symptoms of CRS (eg, fever, hypotension) and ICANS (eg, depressed level of consciousness, seizures) must be captured as individual adverse events and graded by CTCAE criteria. Other neurotoxicities will be graded by CTCAE criteria. Any adverse event or serious adverse event not listed in the NCI CTCAE Version 5.0 will be graded according to investigator clinical judgment by using the standard grades as follows:

- |                |   |
|----------------|---|
| <b>Grade 1</b> | Mild; asymptomatic or mild symptoms; clinical or diagnostic observations only; intervention not indicated.  |
| <b>Grade 2</b> | Moderate; minimal, local or noninvasive intervention indicated; limiting age-appropriate instrumental activities of daily living.*  |
| <b>Grade 3</b> | Severe or medically significant but not immediately life-threatening; hospitalization or prolongation of hospitalization indicated; disabling; limiting self-care activities of daily living.** |
| <b>Grade 4</b> | Life-threatening consequences; urgent intervention indicated.   |
| <b>Grade 5</b> | Death related to adverse event.   |

Activities of Daily Living (ADL)

\* Instrumental ADL refer to preparing meals, shopping for groceries or clothes, using the telephone, managing money, etc.

\*\* Self-care ADL refer to bathing, dressing and undressing, feeding self, using the toilet, taking medications, and not bedridden.

### 12.2. Special Reporting Situations

Safety events of interest on a sponsor study treatment that may require expedited reporting or safety evaluation include, but are not limited to:

- Overdose of a sponsor study treatment
- Suspected abuse/misuse of a sponsor study treatment
- Accidental or occupational exposure to a sponsor study treatment

- Medication error involving a sponsor product (with or without subject/patient exposure to the sponsor study treatment, eg, name confusion)
- Exposure to a sponsor study treatment from breastfeeding

Special reporting situations should be recorded in the CRF. Any special reporting situation that meets the criteria of a serious adverse event should be recorded on the serious adverse event page of the CRF.

## **12.3. Procedures**

### **12.3.1. All Adverse Events**

All adverse events (with the exception of neurotoxicity events, second primary malignancies, see Section 12.3.3, and HBV reactivation) and special reporting situations, whether serious or non-serious, will be reported from the time a signed and dated ICF is obtained until 100 days after infusion of JNJ-68284528 (Cohorts A, B, and C), and 100 days after infusion of JNJ-68284528 or 30 days after last dose of lenalidomide (Cohort D) or D+R (Cohort E), whichever is later. Events of HBV reactivations will be reported during the first year post-dosing of JNJ-68284528. New neurologic adverse events or exacerbation of existing neurologic adverse events will be reported for 1-year post infusion of JNJ-68284528. Beyond the adverse event reporting period, only adverse events that are considered related to study drug need to be reported. Adverse events of special interest are defined in Section 12.3.3.

Serious adverse events, including those spontaneously reported to the investigator within 100 days after infusion of JNJ-68284528 (Cohorts A, B, and C), and 100 days after infusion of JNJ-68284528 or 30 days after last dose of lenalidomide, whichever is later (Cohort D) or D+R (Cohort E) must be reported using the Serious Adverse Event Form. Beyond the serious adverse event reporting period, only serious adverse event that are considered related to study drug need to be reported. The sponsor will evaluate any safety information that is spontaneously reported by an investigator beyond the time frame specified in the protocol. All events that meet the definition of a serious adverse event will be reported as serious adverse events, regardless of whether they are protocol-specific assessments.

Adverse events and special reporting situations, whether serious or non-serious, will be collected for 30 days after the last study procedure for subjects who are enrolled and unable to be apheresed or receive bridging therapy, conditioning regimen, or JNJ-68284528; these subjects will continue to be followed for survival and subsequent anti-myeloma therapies until the end of the study.

All adverse events, regardless of seriousness, severity, or presumed relationship to study treatment, must be recorded using medical terminology in the source document and the CRF. Whenever possible, diagnoses should be given when signs and symptoms are due to a common etiology (eg, cough, runny nose, sneezing, sore throat, and head congestion should be reported as "upper respiratory infection"). The exceptions are CRS and JNJ-68284528-related neurotoxicity, all symptoms associated with these events will be collected in the eCRF. Investigators must record in the CRF their opinion concerning the relationship of the adverse event to study therapy. All

measures required for adverse event management must be recorded in the source document and reported according to sponsor instructions.

Exceptions:

- Progression of disease should not be considered nor should be reported as an adverse event (or serious adverse event). However, signs and symptoms of disease progression or of clinical sequelae resulting from disease progression/lack of efficacy that are determined by the investigator to be of clinical significance should be reported per the usual reporting requirements (Section 12.1).

All deaths not related to disease progression occurring at any time of the study after receiving JNJ-68284528, should be reported to the sponsor following expedited reporting procedures.

The sponsor assumes responsibility for appropriate reporting of adverse events to the regulatory authorities. The sponsor will also report to the investigator (and the head of the investigational institute where required) all suspected unexpected serious adverse reactions (SUSARs). Testing for RCL will be conducted by the sponsor. The sponsor assumes responsibility for appropriate reporting of RCL positive test results to regulatory authorities. The sponsor will notify the investigator of positive RCL test results in a timely manner. These RCL positive test results will not be collected as an adverse event in the eCRF.

The investigator (or sponsor where required) must report SUSARs to the appropriate Independent Ethics Committee/Institutional Review Board (IEC/IRB) that approved the protocol unless otherwise required and documented by the IEC/IRB.

For all studies with an outpatient phase, including open-label studies, the subject must be provided with a "patient ID card" and instructed to carry this card with them for at least 2 years after JNJ-68284528 infusion indicating the following:

- Study number
- Statement, in the local language(s), that the subject is participating in a clinical study
- Investigator's name and 24-hour contact telephone number
- Local sponsor's name and 24-hour contact telephone number (for medical staff only)
- Site number
- Subject number

### **12.3.2. Serious Adverse Events**

All serious adverse events occurring during the study must be reported to the appropriate sponsor contact person by study-site personnel within 24 hours of their knowledge of the event.

Information regarding serious adverse events will be transmitted to the sponsor using the Serious Adverse Event Form, which must be completed and signed by a physician from the study site, and

transmitted to the sponsor within 24 hours. The initial and follow-up reports of a serious adverse event should be made by facsimile (fax).

Events that require an escalation of care when the subject is already hospitalized should be recorded as a serious adverse event. Examples of such events include movement from routine care in the hospital to the ICU or if that event resulted in a prolongation of the existing planned hospitalization.

All serious adverse events that have not resolved by the end of the study, or that have not resolved upon discontinuation of the subject's participation in the study, must be followed until any of the following occurs:

- The event resolves
- The event stabilizes
- The event returns to baseline, if a baseline value/status is available
- The event can be attributed to agents other than the study treatment or to factors unrelated to study conduct
- It becomes unlikely that any additional information can be obtained (subject or health care practitioner refusal to provide additional information, lost to follow-up after demonstration of due diligence with follow-up efforts)

Suspected transmission of an infectious agent by a medicinal product will be reported as a serious adverse event. Any event requiring hospitalization (or prolongation of hospitalization) that occurs during the course of a subject's participation in a study must be reported as a serious adverse event, except hospitalizations for the following:

- Routine monitoring hospitalizations post-infusion required per protocol
- Hospitalizations not intended to treat an acute illness or adverse event (eg, social reasons such as pending placement in long-term care facility)
- Surgery or procedure planned before entry into the study (must be documented in the CRF). Note: Hospitalizations that were planned before the signing of the ICF, and where the underlying condition for which the hospitalization was planned has not worsened, will not be considered serious adverse events. Any adverse event that results in a prolongation of the originally planned hospitalization is to be reported as a new serious adverse event.
- The administration of blood or platelet transfusions. Hospitalization or prolonged hospitalization for a complication of such transfusion remains a reportable serious adverse event.
- The investigator may choose to hospitalize the subject as per institutional standards for CAR-T therapy and in accordance to the criteria provided in [Attachment 14](#).

For subjects in Cohort E, following treatment with daratumumab, if a subject is hospitalized overnight for observation but has not experienced a significant medical event, that hospitalization should not be reported as a serious adverse event.

### 12.3.3. Adverse Events of Special Interest

Cytokine release syndrome, neurotoxicity (including CAR-T cell-related neurotoxicity [eg, ICANS] and other neurotoxicities), tumor lysis syndrome, and second primary malignancy of any grade will be followed as part of standard safety monitoring activities by the sponsor, regardless of severity or causality. All events of CRS, and TLS (any grade) must be reported until 100 days after infusion of JNJ-68284528 (Cohorts A, B, and C), and 100 days after infusion of JNJ-68284528 or 30 days after last dose of lenalidomide (Cohort D) or D+R (Cohort E) or until the start of subsequent anticancer therapy, if earlier. New neurologic adverse events or exacerbation of existing neurologic adverse events will be reported for 1-year post infusion of JNJ-68284528.

Second primary malignancies must be reported for the duration of the study, irrespective of treatment emergent status, and subsequently will be collected in a long-term follow-up study yearly until 15 years post dosing of JNJ-68284528. Testing for RCL will be conducted by the sponsor. The sponsor assumes responsibility for appropriate reporting of RCL positive test results to regulatory authorities. The sponsor will notify the investigator of positive RCL test results in a timely manner. These RCL positive test results will not be collected as an adverse event in the eCRF. Although the risk for RCL is considered very low, there is a theoretical risk that RCL positivity may lead to genotoxicity and subsequent second primary malignancy.

The following events must be reported to the sponsor using the Serious Adverse Event Form within 24 hours of awareness of the event, irrespective of seriousness or causality (eg, serious and nonserious adverse events):

- $\geq$ Grade 3 CRS,
- $\geq$ Grade 3 neurotoxicity,
- $\geq$ Grade 3 tumor lysis syndrome,
- Any second primary malignancies.

Adverse events of special interest that are considered to be non-serious by the investigator are to be included on the serious adverse events form and in the eCRF. Grade 1 or 2 adverse events of special interest would not qualify for expedited reporting unless they meet serious adverse event criteria. All adverse events of special interest of any grade should be followed until recovery or until there is no further improvement.

Because of the embryo-fetal risk of lenalidomide, all subjects must adhere to the global REVLIMID<sup>®</sup> pregnancy prevention program or equivalent local lenalidomide pregnancy prevention program applicable in their region, whichever is more stringent. Investigators should comply with the lenalidomide Global Pregnancy Prevention Plan or with the respective country-specific REVLIMID<sup>®</sup> / Lenalidomide Risk Minimization Program (ie, pregnancy prevention program) as implemented in the post-marketing setting and ensure that all subjects adhere to these programs. When no REVLIMID<sup>®</sup> / Lenalidomide Risk Minimization Program exists, subjects must adhere to the lenalidomide Global Pregnancy Prevention Plan.

#### **12.3.4. Pregnancy**

All initial reports of pregnancy in female subjects or partners of male subjects must be reported to the sponsor by the study-site personnel within 24 hours of their knowledge of the event using the appropriate pregnancy notification form. Abnormal pregnancy outcomes (eg, spontaneous abortion, fetal death, stillbirth, congenital anomalies, ectopic pregnancy) are considered serious adverse events and must be reported using the Serious Adverse Event Form. Any subject who becomes pregnant during the study must discontinue further study treatment. Because the effect of the study treatment on sperm is unknown, pregnancies in partners of male subjects included in the study will be reported as noted above. Follow-up information regarding the outcome of the pregnancy and any postnatal sequelae in the infant will be required.

For subjects in Cohort D and Cohort E: lenalidomide is a thalidomide analogue and is contraindicated for use during pregnancy. Birth defects have been observed in preclinical studies of lenalidomide similar to thalidomide in humans. Therefore, strict monitoring for pregnancy must be conducted during Screening and throughout the Treatment Phase, as specified in the Time and Events Schedule. Investigators should comply with the lenalidomide Global Pregnancy Prevention Plan or with the respective country-specific REVLIMID®/ Risk Minimization Program (ie, Pregnancy prevention program), whichever is more stringent, as implemented in the post-marketing setting and ensure that all subjects adhere to these programs. When no REVLIMID® Risk Minimization Program exists, subjects must adhere to the lenalidomide Global Pregnancy Prevention Plan. If pregnancy does occur, then study treatment should be discontinued immediately, and the subject should be referred to an obstetrician experienced in reproductive toxicity for further evaluation and counseling.

#### **12.4. Contacting Sponsor Regarding Safety**

The names (and corresponding telephone numbers) of the individuals who should be contacted regarding safety issues or questions regarding the study are listed in the Contact Information page(s), which will be provided as a separate document.

### **13. PRODUCT QUALITY COMPLAINT HANDLING**

A product quality complaint (PQC) is defined as any suspicion of a product defect related to manufacturing, labeling, or packaging, ie, any dissatisfaction relative to the identity, quality, durability, or reliability of a product, including its labeling or package integrity. A PQC may have an impact on the safety and efficacy of the product. Timely, accurate, and complete reporting and analysis of PQC information from studies are crucial for the protection of subjects, investigators, and the sponsor, and are mandated by regulatory agencies worldwide. The sponsor has established procedures in conformity with regulatory requirements worldwide to ensure appropriate reporting of PQC information; all studies conducted by the sponsor or its affiliates will be conducted in accordance with those procedures.

#### **13.1. Procedures**

All initial PQCs must be reported to the sponsor by the study-site personnel within 24 hours after being made aware of the event.

If the defect is combined with a serious adverse event, the study-site personnel must report the PQC to the sponsor according to the serious adverse event reporting timelines (refer to Section 12.3.2, Serious Adverse Events). A sample of the suspected product should be maintained for further investigation if requested by the sponsor.

### **13.2. Contacting Sponsor Regarding Product Quality**

The names (and corresponding telephone numbers) of the individuals who should be contacted regarding product quality issues are listed in the Contact Information page(s), which will be provided as a separate document.

## **14. STUDY TREATMENT INFORMATION**

### **14.1. Physical Description of Study Treatment**

JNJ-68284528 therapy is a BCMA-directed genetically modified autologous T cell immunotherapy that involves reprogramming a subject's T cells with a transgene encoding a chimeric antigen receptor (CAR) to identify and eliminate BCMA-expressing malignant and normal cells. Upon binding to BCMA-expressing cells, the CAR transmits a signal to promote T cell expansion, activation, target cell elimination, and persistence of the JNJ-68284528.

### **14.2. Packaging**

JNJ-68284528 will be provided in an infusion bag with specific subject identifiers, this will include subject number and subject apheresis identification number or donor identification number (DIN), subject name and subject date of birth, as allowed by local regulations.

### **14.3. Labeling**

Study treatment labels will contain information to meet the applicable regulatory requirements.

### **14.4. Preparation, Handling, and Storage**

JNJ-68284528 is provided in a single-dose unit containing CAR-positive viable T cells based on the subject weight reported at the time of apheresis.

JNJ-68284528 therapy contains human cells genetically modified with a lentiviral vector. Follow local biosafety guidelines applicable for handling and disposal of such products. The product is prepared from autologous blood collected by apheresis. JNJ-68284528 may carry a risk of transmitting infectious viruses to healthcare professionals handling the product. Accordingly, healthcare professionals should employ universal precautions to avoid potential transmission of infectious diseases when handling the product.

Detailed instructions for storage conditions and handling will accompany clinical drug supplies to the clinical study sites. The storage conditions and expiry dates are indicated on the label. Refer to the IPPI for additional guidance on study treatment preparation, handling, and storage.

## 14.5. Drug Accountability

Information in this section relates to study treatment that is supplied to investigational sites from the study sponsor.

The investigator is responsible for ensuring that all sponsor-provided study treatment received at the site is inventoried and accounted for throughout the study. The study treatment administered to the subject must be documented on the treatment accountability form. All study treatment will be stored and disposed of according to the sponsor's instructions. Study-site personnel must not combine contents of the study treatment containers.

Study treatment must be handled in strict accordance with the protocol and the container label, and must be stored at the study site in a limited-access area or in a locked cabinet under appropriate environmental conditions. Unused study treatment must be available for verification by the sponsor's study site monitor during on-site monitoring visits. The return to the sponsor of unused study treatment will be documented on the treatment return form.

Potentially hazardous materials such as used ampules, needles, syringes and vials containing hazardous liquids, should be disposed of immediately in a safe manner and therefore will not be retained for treatment accountability purposes.

Study treatment should be dispensed under the supervision of the investigator or a qualified member of the study-site personnel, or by a hospital/clinic pharmacist. Study treatment will be supplied only to subjects participating in the study. The investigator agrees neither to dispense the study treatment from, nor store it at, any site other than the study sites agreed upon with the sponsor.

## 15. STUDY-SPECIFIC MATERIALS

The investigator will be provided with the following supplies:

- Study protocol
- Investigator's Brochure (JNJ-68284528)
- IPPI/SIPPM (includes apheresis and cell processing instructions)
- Laboratory manual
- Interactive web response system manual
- Printed PRO questionnaires and PRO Completion Guidelines/Training Materials
- Electronic data capture (eDC) Manual
- Sample ICF
- Subject diaries and instructions/educational materials
- Thermal printer and barcode scanner



## 16. ETHICAL ASPECTS

### 16.1. Study-specific Design Considerations

Potential subjects will be fully informed of the risks and requirements of the study and, during the study, subjects will be given any new information that may affect their decision to continue participation. They will be told that their consent to participate in the study is voluntary and may be withdrawn at any time with no reason given and without penalty or loss of benefits to which they would otherwise be entitled. Only subjects who are fully able to understand the risks, benefits, and potential adverse events of the study, and provide their consent voluntarily will be enrolled.

Despite significant progress in the treatment of patients with multiple myeloma, it still remains uncured, indicating the need for new therapeutic strategies for these patients. As discussed in Section 1.1.5, 74 subjects with relapsed or refractory multiple myeloma have received treatment with LCAR-B38M CAR-T cells in a clinical study setting. An analysis of safety data for these subjects demonstrated a manageable safety profile consistent with its known mechanism of action. In view of the Legend-2 study results and the prognosis for the subject population being considered for this study, a positive risk-benefit profile is anticipated. Subjects will be closely monitored throughout the study, as discussed throughout this protocol, for both safety and clinical benefit.

Apheresis risks may include hypotension, faintness, blurry vision, dizziness, coldness, sweating, infection, abnormal blood clotting, allergic reaction, bleeding, seizures, abdominal cramps, and tingling in the limbs. Subjects will be closely monitored during the procedure and will be evaluated for hospitalization in the case of CRS. All subjects (inpatient and outpatient, Table 10) will be hospitalized from Day 5 to Day 14 after JNJ-68284528 infusion (with potential discharge on Day 10 if there are no CRS, neurotoxicity or other clinically significant events).

The blood sample collection scheme was designed to collect the minimum number of blood samples that accurately and completely describe the pharmacokinetic/pharmacodynamic characteristics of the study treatment. This minimizes the number of venipunctures and the total volume of blood collected from each subject during the study. The volume of blood to be drawn is considered to be customary and acceptable for subjects participating in a cancer clinical study and is deemed reasonable over the timeframe of the study, based upon the standard of the American Red Cross.

### 16.2. Regulatory Ethics Compliance

#### 16.2.1. Investigator Responsibilities

The investigator is responsible for ensuring that the study is performed in accordance with the protocol, current ICH guidelines on Good Clinical Practice (GCP), and applicable regulatory and country-specific requirements.

Good Clinical Practice is an international ethical and scientific quality standard for designing, conducting, recording, and reporting studies that involve the participation of human subjects. Compliance with this standard provides public assurance that the rights, safety, and well-being of

study subjects are protected, consistent with the principles that originated in the Declaration of Helsinki, and that the study data are credible.

### **16.2.2. Independent Ethics Committee or Institutional Review Board**

Before the start of the study, the investigator (or sponsor where required) will provide the IEC/IRB with current and complete copies of the following documents (as required by local regulations):

- Final protocol and, if applicable, amendments
- Sponsor-approved ICF (and any other written materials to be provided to the subjects)
- Investigator's Brochure (or equivalent information) and amendments/addenda
- Sponsor-approved subject recruiting materials
- Information on compensation for study-related injuries or payment to subjects for participation in the study, if applicable
- Investigator's curriculum vitae or equivalent information (unless not required, as documented by the IEC/IRB)
- Information regarding funding, name of the sponsor, institutional affiliations, other potential conflicts of interest, and incentives for subjects
- Any other documents that the IEC/IRB requests to fulfill its obligation

This study will be undertaken only after the IEC/IRB has given full approval of the final protocol, amendments (if any, excluding the ones that are purely administrative, with no consequences for subjects, data or study conduct, unless required locally), the ICF, applicable recruiting materials, and subject compensation programs, and the sponsor has received a copy of this approval. This approval letter must be dated and must clearly identify the IEC/IRB and the documents being approved.

During the study the investigator (or sponsor where required) will send the following documents and updates to the IEC/IRB for their review and approval, where appropriate:

- Protocol amendments (excluding the ones that are purely administrative, with no consequences for subjects, data or study conduct)
- Revision(s) to ICF and any other written materials to be provided to subjects
- If applicable, new or revised subject recruiting materials approved by the sponsor
- Revisions to compensation for study-related injuries or payment to subjects for participation in the study, if applicable
- New edition(s) of the Investigator's Brochure and amendments/addenda
- Summaries of the status of the study at intervals stipulated in guidelines of the IEC/IRB (at least annually)
- Reports of adverse events that are serious, unlisted/unexpected, and associated with the study treatment

- New information that may adversely affect the safety of the subjects or the conduct of the study
- Deviations from or changes to the protocol to eliminate immediate hazards to the subjects
- Report of deaths of subjects under the investigator's care
- Notification if a new investigator is responsible for the study at the site
- Development Safety Update Report and Line Listings, where applicable
- Any other requirements of the IEC/IRB

For all protocol amendments (excluding the ones that are purely administrative, with no consequences for subjects, data or study conduct), the amendment and applicable ICF revisions must be submitted promptly to the IEC/IRB for review and approval before implementation of the change(s).

At least once a year, the IEC/IRB will be asked to review and reapprove this study, where required.

At the end of the study, the investigator (or sponsor where required) will notify the IEC/IRB about the study completion (if applicable, the notification will be submitted through the head of investigational institution).

### **16.2.3. Informed Consent**

Each subject (or a legally acceptable representative) must give written consent according to local requirements after the nature of the study has been fully explained. The ICF(s) must be signed before performance of any study-related activity. The ICF(s) that is/are used must be approved by both the sponsor and by the reviewing IEC/IRB and be in a language that the subject can read and understand. The informed consent should be in accordance with principles that originated in the Declaration of Helsinki, current ICH and GCP guidelines, applicable regulatory requirements, and sponsor policy.

Before enrollment in the study, the investigator or an authorized member of the study-site personnel must explain to potential subjects or their legally acceptable representatives the aims, methods, reasonably anticipated benefits, and potential hazards of the study, and any discomfort participation in the study may entail. Subjects will be informed that their participation is voluntary and that they may withdraw consent to participate at any time. They will be informed that choosing not to participate will not affect the care the subject will receive for the treatment of his or her disease. Subjects will be told that alternative treatments are available if they refuse to take part and that such refusal will not prejudice future treatment. Finally, they will be told that the investigator will maintain a subject identification register for the purposes of long-term follow up if needed and that their records may be accessed by health authorities and authorized sponsor personnel without violating the confidentiality of the subject, to the extent permitted by the applicable law(s) or regulations. By signing the ICF the subject or legally acceptable representative is authorizing such access, which includes permission to obtain information about his or her survival status. It also denotes that the subject agrees to allow his or her study physician to recontact the subject for the purpose of obtaining consent for additional safety evaluations, and subsequent disease-related

treatments, if needed. The physician may also recontact the subject for the purpose of obtaining consent to collect information about his or her survival status.

The subject or legally acceptable representative will be given sufficient time to read the ICF and the opportunity to ask questions. After this explanation and before entry into the study, consent should be appropriately recorded by means of either the subject's or his or her legally acceptable representative's personally dated signature. After having obtained the consent, a copy of the ICF must be given to the subject.

If the subject or legally acceptable representative is unable to read or write, an impartial witness should be present for the entire informed consent process (which includes reading and explaining all written information) and should personally date and sign the ICF after the oral consent of the subject or legally acceptable representative is obtained.

When prior consent of the subject is not possible and the subject's legally acceptable representative is not available, enrollment procedures should be described in the protocol with documented approval/favorable opinion by the IEC/IRB to protect the rights, safety, and well-being of the subject and to ensure compliance with applicable regulatory requirements. The subject or legally acceptable representative must be informed about the study as soon as possible and give consent to continue.

#### **16.2.4. Privacy of Personal Data**

The collection and processing of personal data from subjects enrolled in this study will be limited to those data that are necessary to fulfill the objectives of the study.

These data must be collected and processed with adequate precautions to ensure confidentiality and compliance with applicable data privacy protection laws and regulations. Appropriate technical and organizational measures to protect the personal data against unauthorized disclosures or access, accidental or unlawful destruction, or accidental loss or alteration must be put in place. For tracking and traceability of the apheresis material and investigational product, subject name and date of birth, as allowed by local regulations, will be collected to ensure chain of identity of the investigational product for the subject. Sponsor personnel whose responsibilities require access to personal data agree to keep the identity of subjects confidential.

The informed consent obtained from the subject (or his or her legally acceptable representative) includes explicit consent for the processing of personal data and for the investigator/institution to allow direct access to his or her original medical records (source data/documents) for study-related monitoring, audit, IEC/IRB review, and regulatory inspection. This consent also addresses the transfer of the data to other entities and to other countries.

The subject has the right to request through the investigator access to his or her personal data and the right to request rectification of any data that are not correct or complete. Reasonable steps will be taken to respond to such a request, taking into consideration the nature of the request, the conditions of the study, and the applicable laws and regulations.

Exploratory biomarker, pharmacokinetic and immunogenicity research is not conducted under standards appropriate for the return of data to subjects. In addition, the sponsor cannot make decisions as to the significance of any findings resulting from exploratory research. Therefore, exploratory research data will not be returned to subjects or investigators, unless required by law or local regulations. Privacy and confidentiality of data generated in the future on stored samples will be protected by the same standards applicable to all other clinical data.

#### **16.2.5. Long-term Retention of Samples for Additional Future Research**

Samples, including apheresis product, collected in this study and JNJ-68284528 that was manufactured but not administered to a subject may be stored for up to 15 years (or according to local regulations) for additional research. Samples will only be used to understand JNJ-68284528, to understand multiple myeloma, to understand differential intervention responders, and to develop tests/assays related to JNJ-68284528. The research may begin at any time during the study or the post-study storage period.

Stored samples will be coded throughout the sample storage and analysis process and will not be labeled with personal identifiers. Subjects may withdraw their consent for their samples to be stored for research (refer to Section 10.4, Withdrawal from the Use of Samples in Future Research).

#### **16.2.6. Country Selection**

This study will only be conducted in those countries where the intent is to launch or otherwise help ensure access to the developed product if the need for the product persists, unless explicitly addressed as a specific ethical consideration in Section 16.1, Study-Specific Design Considerations.

### **17. ADMINISTRATIVE REQUIREMENTS**

#### **17.1. Protocol Amendments**

Neither the investigator nor the sponsor will modify this protocol without a formal amendment by the sponsor. All protocol amendments must be issued by the sponsor, and signed and dated by the investigator. Protocol amendments must not be implemented without prior IEC/IRB approval, or when the relevant competent authority has raised any grounds for non-acceptance, except when necessary to eliminate immediate hazards to the subjects, in which case the amendment must be promptly submitted to the IEC/IRB and relevant competent authority. Documentation of amendment approval by the investigator and IEC/IRB must be provided to the sponsor. When the change(s) involves only logistic or administrative aspects of the study, the IEC/IRB (where required) only needs to be notified.

During the course of the study, in situations where a departure from the protocol is unavoidable, the investigator or other physician in attendance will contact the appropriate sponsor representative listed in the Contact Information page(s), which will be provided as a separate document. Except in emergency situations, this contact should be made before implementing any departure from the protocol. In all cases, contact with the sponsor must be made as soon as possible to discuss the

situation and agree on an appropriate course of action. The data recorded in the CRF and source documents will reflect any departure from the protocol, and the source documents will describe this departure and the circumstances requiring it.

## **17.2. Regulatory Documentation**

### **17.2.1. Regulatory Approval/Notification**

This protocol and any amendment(s) must be submitted to the appropriate regulatory authorities in each respective country, if applicable. A study may not be initiated until all local regulatory requirements are met.

### **17.2.2. Required Prestudy Documentation**

The following documents must be provided to the sponsor before shipment of study treatment to the study site:

- Protocol and amendment(s), if any, signed and dated by the principal investigator
- A copy of the dated and signed (or sealed, where appropriate per local regulations), written IEC/IRB approval of the protocol, amendments, ICF, any recruiting materials, and if applicable, subject compensation programs. This approval must clearly identify the specific protocol by title and number and must be signed (or sealed, where appropriate per local regulations) by the chairman or authorized designee.
- Name and address of the IEC/IRB, including a current list of the IEC/IRB members and their function, with a statement that it is organized and operates according to GCP and the applicable laws and regulations. If accompanied by a letter of explanation, or equivalent, from the IEC/IRB, a general statement may be substituted for this list. If an investigator or a member of the study-site personnel is a member of the IEC/IRB, documentation must be obtained to state that this person did not participate in the deliberations or in the vote/opinion of the study.
- Regulatory authority approval or notification, if applicable
- Signed and dated statement of investigator (eg, Form FDA 1572), if applicable
- Documentation of investigator qualifications (eg, curriculum vitae)
- Completed investigator financial disclosure form from the principal investigator, where required
- Signed and dated clinical trial agreement, which includes the financial agreement
- Any other documentation required by local regulations

The following documents must be provided to the sponsor before enrollment of the first subject:

- Completed investigator financial disclosure forms from all subinvestigators
- Documentation of subinvestigator qualifications (eg, curriculum vitae)

- Name and address of any local laboratory conducting tests for the study, and a dated copy of current laboratory normal ranges for these tests, if applicable
- Local laboratory documentation demonstrating competence and test reliability (eg, accreditation/license), if applicable

### **17.3. Subject Identification, Enrollment, and Screening Logs**

The investigator agrees to complete a subject identification and enrollment log to permit easy identification of each subject during and after the study. This document will be reviewed by the sponsor study-site contact for completeness.

The subject identification and enrollment log will be treated as confidential and will be filed by the investigator in the study file. To ensure subject confidentiality, no copy will be made. All reports and communications relating to the study will identify subjects by subject identification and date of birth (as allowed by local regulations). In cases where the subject is not enrolled into the study, the date seen and date of birth (as allowed by local regulations) will be used.

For tracking and traceability of the apheresis material and investigational product, subject name and date of birth, as allowed by local regulations, will be collected to ensure chain of identity of the investigational product for the subject.

The investigator must also complete a subject screening log, which reports on all subjects who were seen to determine eligibility for inclusion in the study.

### **17.4. Source Documentation**

At a minimum, source documents consistent in the type and level of detail with that commonly recorded at the study site as a basis for standard medical care must be available for the following: subject identification, eligibility, and study identification; study discussion and date of signed informed consent; dates of visits; results of safety and efficacy parameters as required by the protocol; record of all adverse events and follow-up of adverse events; concomitant medication; treatment receipt/dispensing/return records; study treatment administration information; and date of study completion and reason for early discontinuation of study treatment or withdrawal from the study, if applicable.

The author of an entry in the source documents should be identifiable.

Specific details required as source data for the study and source data collection methods will be reviewed with the investigator before the study and will be described in the monitoring guidelines (or other equivalent document).

The minimum source documentation requirements for Section 4, Inclusion Criteria and Exclusion Criteria that specify a need for documented medical history are as follows:

- Referral letter from treating physician or
- Complete history of medical notes at the site
- Discharge summaries

Inclusion and exclusion criteria not requiring documented medical history must be verified at a minimum by subject interview or other protocol required assessment (eg, physical examination, laboratory assessment) and documented in the source documents.

An electronic source system may be utilized, which contains data traditionally maintained in a hospital or clinic record to document medical care (eg, electronic source documents) as well as the clinical study-specific data fields as determined by the protocol. This data is electronically extracted for use by the sponsor. If the electronic source system is utilized, references made to the CRF in the protocol include the electronic source system but information collected through the electronic source system may not be limited to that found in the CRF. Data in this system may be considered source documentation.

### **17.5. Case Report Form Completion**

Case report forms are prepared and provided by the sponsor for each subject in electronic format. All CRF entries, corrections, and alterations must be made by the investigator or authorized study-site personnel. The investigator must verify that all data entries in the CRF are accurate and correct.

The study data will be transcribed by study-site personnel from the source documents onto an electronic CRF, if applicable. Study-specific data will be transmitted in a secure manner to the sponsor.

Worksheets may be used for the capture of some data to facilitate completion of the CRF. Any such worksheets will become part of the subject's source documents. Data must be entered into CRF in English. The CRF must be completed as soon as possible after a subject visit and the forms should be available for review at the next scheduled monitoring visit.

If necessary, queries will be generated in the eDC tool. If corrections to a CRF are needed after the initial entry into the CRF, this can be done in either of the following ways:

- Investigator and study-site personnel can make corrections in the eDC tool at their own initiative or as a response to an auto query (generated by the eDC tool).
- Sponsor or sponsor delegate can generate a query for resolution by the investigator and study-site personnel.

### **17.6. Data Quality Assurance/Quality Control**

Steps to be taken to ensure the accuracy and reliability of data include the selection of qualified investigators and appropriate study sites, review of protocol procedures with the investigator and study-site personnel before the study, periodic monitoring visits by the sponsor, and where applicable direct transmission of clinical laboratory data from a central laboratory into the



sponsor's data base. Written instructions will be provided for collection, handling, storage, and shipment of samples.

Guidelines for CRF completion will be provided and reviewed with study-site personnel before the start of the study. The sponsor will review CRF for accuracy and completeness during on-site monitoring visits and after transmission to the sponsor; any discrepancies will be resolved with the investigator or designee, as appropriate. After upload of the data into the study database they will be verified for accuracy and consistency with the data sources.

### **17.7. Record Retention**

In compliance with the ICH/GCP guidelines, the investigator/institution will maintain all CRF and all source documents that support the data collected from each subject, as well as all study documents as specified in ICH/GCP Section 8, Essential Documents for the Conduct of a Clinical Trial, and all study documents as specified by the applicable regulatory requirement(s). The investigator/institution will take measures to prevent accidental or premature destruction of these documents.

Essential documents must be retained until at least 2 years after the last approval of a marketing application in an ICH region and until there are no pending or contemplated marketing applications in an ICH region or until at least 2 years have elapsed since the formal discontinuation of clinical development of the investigational product. These documents will be retained for a longer period if required by the applicable regulatory requirements or by an agreement with the sponsor. It is the responsibility of the sponsor to inform the investigator/institution as to when these documents no longer need to be retained.

If the responsible investigator retires, relocates, or for other reasons withdraws from the responsibility of keeping the study records, custody must be transferred to a person who will accept the responsibility. The sponsor must be notified in writing of the name and address of the new custodian. Under no circumstance shall the investigator relocate or dispose of any study documents before having obtained written approval from the sponsor.

If it becomes necessary for the sponsor or the appropriate regulatory authority to review any documentation relating to this study, the investigator/institution must permit access to such reports.

### **17.8. Monitoring**

The sponsor will use a combination of monitoring techniques (central, remote, or on-site monitoring) to monitor this study.

The sponsor will perform on-site monitoring visits as frequently as necessary. The monitor will record dates of the visits in a study site visit log that will be kept at the study site. The first post-initiation visit will be made as soon as possible after enrollment has begun. At these visits, the monitor will compare the data entered into the CRF with the source documents (eg, hospital/clinic/physician's office medical records). The nature and location of all source documents will be identified to ensure that all sources of original data required to complete the

CRF are known to the sponsor and study-site personnel and are accessible for verification by the sponsor study-site contact. If electronic records are maintained at the study site, the method of verification must be discussed with the study-site personnel.

Direct access to source documents (medical records) must be allowed for the purpose of verifying that the recorded data are consistent with the original source data. Findings from this review will be discussed with the study-site personnel. The sponsor expects that, during monitoring visits, the relevant study-site personnel will be available, the source documents will be accessible, and a suitable environment will be provided for review of study-related documents. The monitor will meet with the investigator on a regular basis during the study to provide feedback on the study conduct.

In addition to on-site monitoring visits, remote contacts can occur. It is expected that during these remote contacts, study-site personnel will be available to provide an update on the progress of the study at the site.

## **17.9. Study Completion/Termination**

### **17.9.1. Study Completion/End of Cohort**

Cohorts A, B, and C will each be considered complete after the last subject has had two years of follow-up after the initial dose of JNJ-68284528. Cohort D and Cohort E will be considered complete 2 ½ years after the last subject receives their initial dose of JNJ-68284528.

The final data from the study site will be sent to the sponsor (or designee) after completion of the final subject assessment at that study site, in the time frame specified in the Clinical Trial Agreement.

### **17.9.2. Study Termination**

The sponsor reserves the right to close the study site or terminate the study at any time for any reason at the sole discretion of the sponsor. Study sites will be closed upon study completion. A study site is considered closed when all required documents and study supplies have been collected and a study-site closure visit has been performed.

The investigator may initiate study-site closure at any time, provided there is reasonable cause and sufficient notice is given in advance of the intended termination.

Reasons for the early closure of a study site by the sponsor or investigator may include but are not limited to:

- Failure of the investigator to comply with the protocol, the requirements of the IEC/IRB or local health authorities, the sponsor's procedures, or GCP guidelines
- Inadequate recruitment of subjects by the investigator
- Discontinuation of further study treatment development

### **17.10. On-site Audits**

Representatives of the sponsor's clinical quality assurance department may visit the study site at any time during or after completion of the study to conduct an audit of the study in compliance with regulatory guidelines and company policy. These audits will require access to all study records, including source documents, for inspection. Subject privacy must, however, be respected. The investigator and study-site personnel are responsible for being present and available for consultation during routinely scheduled study-site audit visits conducted by the sponsor or its designees.

Similar auditing procedures may also be conducted by agents of any regulatory body, either as part of a national GCP compliance program or to review the results of this study in support of a regulatory submission. The investigator should immediately notify the sponsor if he or she has been contacted by a regulatory agency concerning an upcoming inspection.

### **17.11. Use of Information and Publication**

All information, including but not limited to information regarding JNJ-68284528 or the sponsor's operations (eg, patent application, formulas, manufacturing processes, basic scientific data, prior clinical data, formulation information) supplied by the sponsor to the investigator and not previously published, and any data, including exploratory biomarker research data, generated as a result of this study, are considered confidential and remain the sole property of the sponsor. The investigator agrees to maintain this information in confidence and use this information only to accomplish this study, and will not use it for other purposes without the sponsor's prior written consent.

The investigator understands that the information developed in the study will be used by the sponsor in connection with the continued development of JNJ-68284528, and thus may be disclosed as required to other clinical investigators or regulatory agencies. To permit the information derived from the clinical studies to be used, the investigator is obligated to provide the sponsor with all data obtained in the study.

The results of the study will be reported in a Clinical Study Report generated by the sponsor and will contain data from all study sites that participated in the study as per protocol. Recruitment performance or specific expertise related to the nature and the key assessment parameters of the study will be used to determine a coordinating investigator for the study. Results of exploratory biomarker analyses performed after the Clinical Study Report has been issued will be reported in a separate report and will not require a revision of the Clinical Study Report. Study subject identifiers will not be used in publication of results. Any work created in connection with performance of the study and contained in the data that can benefit from copyright protection (except any publication by the investigator as provided for below) shall be the property of the sponsor as author and owner of copyright in such work.

Consistent with Good Publication Practices and International Committee of Medical Journal Editors guidelines, the sponsor shall have the right to publish such primary (multicenter) data and information without approval from the investigator. The investigator has the right to publish study

site-specific data after the primary data are published. If an investigator wishes to publish information from the study, a copy of the manuscript must be provided to the sponsor for review at least 60 days before submission for publication or presentation. Expedited reviews will be arranged for abstracts, poster presentations, or other materials. If requested by the sponsor in writing, the investigator will withhold such publication for up to an additional 60 days to allow for filing of a patent application. In the event that issues arise regarding scientific integrity or regulatory compliance, the sponsor will review these issues with the investigator. The sponsor will not mandate modifications to scientific content and does not have the right to suppress information. For multicenter study designs and substudy approaches, secondary results generally should not be published before the primary endpoints of a study have been published. Similarly, investigators will recognize the integrity of a multicenter study by not submitting for publication data derived from the individual study site until the combined results from the completed study have been submitted for publication, within 18 months after study end date, or the sponsor confirms there will be no multicenter study publication. Authorship of publications resulting from this study will be based on the guidelines on authorship, such as those described in the ICMJE Recommendations for the Conduct, Reporting, Editing and Publication of Scholarly Work in Medical Journals, which state that the named authors must have made a significant contribution to the conception or design of the work; or the acquisition, analysis, or interpretation of the data for the work; and drafted the work or revised it critically for important intellectual content; and given final approval of the version to be published; and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

### **Registration of Clinical Studies and Disclosure of Results**

The sponsor will register and disclose the existence of and the results of clinical studies as required by law.

---

**REFERENCES**

1. ACTEMRA® (tocilizumab). Prescribing Information. South San Francisco, CA: Genentech, Inc;2017.
2. Adams HC, Stevenaert F, Krejcik J, et al. High-Parameter Mass Cytometry Evaluation of Relapsed/Refractory Multiple Myeloma Patients Treated with Daratumumab Demonstrates Immune Modulation as a Novel Mechanism of Action. *Cytometry A*. 2019;95(3):279-289.
3. Adams H, Van der Borch K, Abraham Y, et al. Effects of daratumumab on the composition and activation status of immune-cell populations in Centaurus, a phase 2 randomized study of smoldering multiple myeloma (SMM) patients. *EHA*. 2018;S1577.
4. Attal M, Lauwers-Cances V, Hulin C, et al. Lenalidomide, Bortezomib, and Dexamethasone with Transplantation for Myeloma. *NEJM*. 2017;376(14):1311-1320.
5. Avery DT, Kalled SL, Ellyard JJ, et al. BAFF selectively enhances the survival of plasmablasts generated from human memory B cells. *J Clin Invest*. 2003;112(2):286-297.
6. Avet-Loiseau H, Casneuf T, Chiu C, et al. Evaluation of minimal residual disease (MRD) in relapsed/refractory multiple myeloma (RRMM) patients treated with daratumumab in combination with lenalidomide plus dexamethasone or bortezomib plus dexamethasone. *ASH Annual Meeting Abstracts*. 2016; Abstract 246. <http://ash.confex.com/ash/2016/webprogram/Paper96569.html>. Accessed on 08 May 2017.
7. Bossen C, Schneider P. BAFF, APRIL and their receptors: structure, function and signaling. *Semin Immunol*. 2006;18(5):263-275.
8. Burtis CA, Ashwood ER. *Tietz Textbook of Clinical Chemistry*, 3rd ed. Philadelphia; WB Saunders, 1998.
9. Carpenter RO, Evbuomwan MO, Pittaluga S, et al. B-cell maturation antigen is a promising target for adoptive T-cell therapy of multiple myeloma. *Clin Cancer Res*. 2013;19(8):2048-2060.
10. Cavo M, Terpos E, Nanni C, et al. Role of 18F-FDG PET/CT in the diagnosis and management of multiple myeloma and other plasma cell disorders: a consensus statement by the International Myeloma Working Group. *Lancet Oncol*. 2017;18(4):e206-e217.
11. Chapuy CI, Nicholson, RT, Aguad, MD, et al. Resolving the daratumumab interference with blood compatibility testing. *Transfusion*. 2015;55:1545-1554.
12. Chapuy CI, Aguad, MD, Nicholson, RT, et al. International validation of a dithiothreitol (DTT)-based method to resolve the daratumumab interference with blood compatibility testing. *Transfusion*. 2016;56:2964-2972.
13. Cheson BD, Pfistner B, Juweid ME, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol*. 2007;25(5):579-586.
14. Cho S-F, Anderson KC, Tai Y-T. BCMA CAR-T cell therapy arrives for multiple myeloma: a reality. *Ann Transl Med*. 2018a;6(Suppl 2):S93. doi: 10.21037/atm.2018.11.14.
15. Cho S-F, Anderson KC, Tai Y-T. Targeting B cell maturation antigen (BCMA) in multiple myeloma: potential uses of BCMA-based immunotherapy. *Front Immunol*. 2018b;10(9):1821. doi: 10.3389/fimmu.2018.01821. eCollection 2018.
16. Darce JR, Arendt BK, Chang SK, Jelinek DF. Divergent effects of BAFF on human memory B cell differentiation into Ig-secreting cells. *J Immunol*. 2007;178(9):5612-5622.
17. de Haart SJ, van de Donk NW, Minnema MC, et al. Accessory cells of the microenvironment protect multiple myeloma from T-cell cytotoxicity through cell adhesion-mediated immune resistance. *Clin Cancer Res*. 2013;19(20):5591-5601.
18. Dimopoulos MA, Oriol A, Nahi H, et al. Daratumumab, lenalidomide, and dexamethasone for multiple myeloma. *N Engl J Med*. 2016;375:1319-1331.
19. Durie BG, Harousseau JL, Miguel JS, et al. International uniform response criteria for multiple myeloma. *Leukemia*. 2006;20(9):1467-1473.
20. Durie BG, Miguel JF, Blade J, Rajkumar SV. Clarification of the definition of complete response in multiple myeloma. *Leukemia*. 2015;29(12):2416-2417.

21. Durie BG, Hoering A, Abidi MH, et al. Bortezomib with lenalidomide and dexamethasone versus lenalidomide and dexamethasone alone in patients with newly diagnosed myeloma without intent for immediate autologous stem-cell transplant (SWOG S0777): a randomised, open-label, phase 3 trial. *Lancet*. 2017;389:519-527.
22. Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1) *Eur J Cancer*. 2009;45:228-247.
23. EORTC Quality of Life Group. Guidelines for assessing quality of life in EORTC clinical trials. <http://groups.eortc.be/qol/manuals>. Accessed 22 July 2018.
24. Fraietta JA, Lacey SF, Orlando EJ, et al. Determinants of response and resistance to CD19 chimeric antigen receptor (CAR) T cell therapy of chronic lymphocytic leukemia. *Nat Med*. 2018;24(5):563-571.
25. Frigyesi I, Adolfsson J, Ali M, et al. Robust isolation of malignant plasma cells in multiple myeloma. *Blood*. 2014;123(9):1336-1340.
26. Gore L, Locatelli F, Zugmaier G, et al. Survival after blinatumomab treatment in pediatric patients with relapsed/refractory B-cell precursor acute lymphoblastic leukemia. *Blood Cancer J*. 2018;8(9):80. doi: 10.1038/s41408-018-0117-0.
27. Greipp PR, San Miguel J, Durie BGM, et al. International staging system for multiple myeloma. *J Clin Oncol*. 2005;23(15):3412-3420.
28. Hatzoglou A, Roussel J, Bourgeade MF, et al. TNF receptor family member BCMA (B cell maturation) associates with TNF receptor-associated factor (TRAF) 1, TRAF2, and TRAF3 and activates NF-kappa B, elk-1, c-Jun N-terminal kinase, and p38 mitogen-activated protein kinase. *J Immunol*. 2000;165(3):1322-1330.
29. Hillengass J, Usmani S, Rajkumar SV, et al. International Myeloma Working Group consensus recommendations on imaging in monoclonal plasma cell disorders. *Lancet Oncol*. 2019;20(6):e302-e312.
30. Hsi ED, Steinle R, Balasa B, et al. CS1, a potential new therapeutic antibody target for the treatment of multiple myeloma. *Clin Cancer Res*. 2008;14(9):2775-2784.
31. Hwang JP, Artz AS, Somerfield MR. Hepatitis B Virus Screening for Patients With Cancer Before Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update. *J Oncol Pract*. 2015;11(4):e487-489.
32. Jimenez-Zepeda VH, Reece DE, Trudel S, et al. Early relapse after single auto-SCT for multiple myeloma is a major predictor of survival in the era of novel agents. *Bone Marrow Transplantation*. 2015;50:204-208.
33. Kimberley FC, Hahne M, Medema JP. "APRIL hath put a spring of youth in everything": Relevance of APRIL for survival. *J Cell Physiol*. 2009;218(1):1-8.
34. Korde N, Kristinsson SY, Landgren O. Monoclonal gammopathy of undetermined significance (MGUS) and smoldering multiple myeloma (SMM): novel biological insights and development of early treatment strategies. *Blood*. 2011;117(21):5573-5581.
35. Krejcik J, Casneuf T, Nijhof IS, et al. Daratumumab depletes CD38+ immune regulatory cells, promotes T-cell expansion, and skews T-cell repertoire in multiple myeloma. *Blood*. 2016;128(3):384-394.
36. Kumar S, Paiva B, Anderson KC, et al. International Myeloma Working Group consensus criteria for response and minimal residual disease assessment in multiple myeloma. *Lancet Oncol*. 2016;17(8):e328-e346.
37. Kuramitsu S, Ohno M, Ohka F, et al. Lenalidomide enhances the function of chimeric antigen receptor T cells against the epidermal growth factor receptor variant III by enhancing immune synapses. *Cancer Gene Ther*. 2015;22(10):487-495.
38. Kurki S, Tamminen K, Miettinen T, Remes K. Prognostic comparison between ISS and R-ISS in real-life setting of myeloma patients: An example of utilization of electronic biobank database. *Blood*. 2016;128(22):5645.
39. Kyle RA, Rajkumar SV. Criteria for diagnosis, staging, risk stratification and response assessment of multiple myeloma. *Leukemia*. 2009;23(1):3-9.
40. Lahuerta JJ, Mateos MV, Martinez-Lopez J, et al. Influence of pre- and post-transplantation responses on outcome of patients with multiple myeloma: sequential improvement of response and achievement of complete response are associated with longer survival. *J Clin Oncol*. 2008;26:5575-5782.

41. Landgren O, Giral S. MRD-driven treatment paradigm for newly diagnosed transplant eligible multiple myeloma patients. *Bone Marrow Transplant*. 2016;51(7):913-914.
42. Lee DW, Gardner R, Porter DL, et al. Current concepts in the diagnosis and management of cytokine release syndrome. *Blood*. 2014;124(2):188-195.
43. Lee DW, Santomaso BD, Locke FL, et al. ASTCT consensus grading for cytokine release syndrome and neurological toxicity associated with immune effector cells. *Biol Blood Marrow Transplant*. 2019;25(4):625-638.
44. Levey AS, Coresh J, Greene T, et al. Using standardized serum creatinine values in the modification of diet in renal disease study equation for estimating glomerular filtration rate. *Ann Intern Med*. 2006;145:247-254.
45. Majithia N, Rajkumar SV, Lacy MQ, et al. Early relapse following initial therapy for multiple myeloma predicts poor outcomes in the era of novel agents. *Leukemia*. 2016;30:2208-2213.
46. MD Anderson Cancer Center. IEC Therapy Toxicity Assessment and Management. 2019. Appendix J. <https://www.mdanderson.org/documents/for-physicians/algorithms/clinical-management/clin-management-cytokine-release-web-algorithm.pdf>
47. Messiou C, Hillengass J, Delorme S, et al. Guidelines for Acquisition, Interpretation, and Reporting of Whole-Body MRI in Myeloma: Myeloma Response Assessment and Diagnosis System (MY-RADS). *Radiology*. 2019;291(1):5-13.
48. Moreau P, Joshua D, Chng WJ, et al. Impact of prior treatment on patients with relapsed multiple myeloma treated with carfilzomib and dexamethasone vs bortezomib and dexamethasone in the phase 3 ENDEAVOR study. *Leukemia*. 2017;31(1):115-122.
49. Mosteller RD. Simplified calculation of body-surface area. *NEJM*. 1987;317(17):1098.
50. Munshi NC, Avet-Loiseau H, Rawstron AC. Association of minimal residual disease with superior survival outcomes in patients with multiple myeloma (a meta-analysis). *JAMA Oncol*. 2016; doi:10.1001/jamaoncol.2016.3160.
51. Neelapu SS, Tummala S, Kebriaei P. Chimeric antigen receptor T-cell therapy – assessment and management of toxicities. *Nat Rev Clin Oncol*. 2018;15(1):47-62.
52. Ong SY, de Mel S, Chen YX, et al. Early relapse post autologous transplant is a stronger predictor of survival compared with pretreated patient factors in the novel agent era: analysis of the Singapore Multiple Myeloma Working Group. *Bone Marrow Transplantation*. 2016;51:933-937.
53. Oken MM, Creech RH, Tormey DC, et al. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *Am J Clin Oncol*. 1982;5(6):649-655.
54. Otahal P, Prukova D, Krai V, et al. Lenalidomide enhances antitumor functions of chimeric antigen receptor modified T cells. *Oncoimmunology*. 2016;5(4):e1115940.
55. Palaiologou M, Delladetsima I, Tiniakos D. CD138 (syndecan-1) expression in health and disease. *Histol Histopathol*. 2014;29(2):177-189.
56. Palumbo A, Anderson K. Multiple myeloma. *N Engl J Med*. 2011;364(11):1046-1060.
57. Palumbo A, Bringhen S, Mateos MV, et al. Geriatric assessment predicts survival and toxicities in elderly myeloma patients: an International Myeloma Working Group report. *Blood*. 2015;125(13):2068-2070.
58. Palumbo A, Chanan-Khan A, Weisel K, et al. Daratumumab, bortezomib, and dexamethasone for multiple myeloma. *N Engl J Med*. 2016;375:754-766.
59. Patel DR, Wallweber HJ, Yin J, et al. Engineering an APRIL-specific B cell maturation antigen. *J Biol Chem*. 2004;279(16):16727-16735.
60. PRO-CTCAE Instrument and Form Builder. <https://healthcaresdelivery.cancer.gov/pro-ctcae/instrument.html>. Accessed 11 April 2019.
61. Raje N, Berdeja J, Lin Y, et al. Anti-BCMA CAR T-cell therapy bb2121 in relapsed or refractory multiple myeloma. *NEJM*. 2019;380:1726-1737.

62. Rajkumar SV, Dimopoulos MA, Palumbo A, et al. International Myeloma Working Group updated criteria for the diagnosis of multiple myeloma. *Lancet Oncol.* 2014;15(12):e538-548.
63. Rajkumar SV, Harousseau JL, Durie B, et al. Consensus recommendations for the uniform reporting of clinical trials: report of the International Myeloma Workshop Consensus Panel 1. *Blood.* 2011;117(18):4691-4695.
64. Richardson PG, Oriol A, Beksac M, et al. Pomalidomide, bortezomib, and dexamethasone for patients with relapsed or refractory multiple myeloma previously treated with lenalidomide (OPTIMISMM): a randomised, open-label, phase 3 trial. *Lancet Oncol.* 2019;20(6):781-794.
65. Tai YT, Anderson KC. Targeting B-cell maturation antigen in multiple myeloma. *Immunotherapy.* 2015;7(11):1187-1199.
66. Topp MS, Duell J, Zugmaier G, et al. Evaluation of AMG 420, and anti-BCMA bispecific T-cell engager (BiTE) immunotherapy, in R/R multiple myeloma (MM) patients: Updated results of a first-in-human (FIH) phase I dose escalation study. *J Clin Oncol.* 2019;37(15)\_suppl:8007.
67. Trask PC, Dueck AC, Piau E, Campbell A. Patient-reported outcomes version of the common terminology criteria for adverse events: Methods for item selection in industry-sponsored oncology clinical trials. *Clin Trials.* 2018;15(6):616-623.
68. Trudel S, Lendvai N, Popat R, et al. Antibody-drug conjugate, GSK2857916, in relapsed/refractory multiple myeloma: and update on safety and efficacy from dose expansion phase 1 study. *Blood Cancer J.* 2019;9(4):37. doi: 10.1038/s41408-019-0196-6.
69. Usmani SZ, Nahi H, Mateos MV, et al. Subcutaneous delivery of daratumumab in relapsed or refractory multiple myeloma. *Blood.* 2019;134(8):668-677.
70. van de Donk N, Richardson PG, Malavasi F. CD38 antibodies in multiple myeloma: back to the future. *Blood.* 2018;131(1):13-29.
71. van de Donk NW, Adams H, Vanhoof G, et al. Daratumumab in Combination with Lenalidomide Plus Dexamethasone Results in Persistent Natural Killer (NK) Cells with a Distinct Phenotype and Expansion of Effector Memory T-Cells in Pollux, a Phase 3 Randomized Study. *Blood.* 2017;130(Suppl 1):3124.
72. Voorhees PM, Kaufman JL, Laubach JP, et al. Depth of response to daratumumab (DARA), lenalidomide, bortezomib, and dexamethasone (RVd) improves over time in patients (pts) with transplant-eligible newly diagnosed multiple myeloma (NDMM): Griffin study update. *ASH Annual Meeting.* 2019. Abstract 691.
73. Voorhees PM, Kaufman JL, Laubach JP, et al. Daratumumab, lenalidomide, bortezomib, and dexamethasone for transplant-eligible newly diagnosed multiple myeloma: GRIFFIN. *Blood.* 2020; 2020005288. doi: 10.1182/blood.2020005288. Online ahead of print.
74. Wang X, Walter M, Urak R, et al. Lenalidomide enhances the function of CS1 chimeric antigen receptor-redirected T cells against multiple myeloma. *Clin Cancer Res.* 2018;24(1):106-119.
75. Wisloff F, Eika S, Hippe E, et al. Measurement of health-related quality of life in multiple myeloma. *Nordic Myeloma Study Group. Br J Haematol.* 1996;92:604-613.
76. Wisloff F, Hjorth M. Health-related quality of life assessed before and during chemotherapy predicts for survival in multiple myeloma. *Nordic Myeloma Study Group. Br J Haematol.* 1997;97:29-37.



**Attachment 1: Criteria for Response to Multiple Myeloma Treatment**

<b>Response</b>	<b>Response Criteria</b>
Stringent complete response	<ul style="list-style-type: none"> <li>CR as defined below, <i>plus</i></li> <li>Normal FLC ratio, <i>and</i></li> <li>Absence of clonal PCs by immunohistochemistry or 2- to 4-color flow cytometry</li> </ul>
Complete response <sup>a</sup>	<ul style="list-style-type: none"> <li>Negative immunofixation of serum and urine, <i>and</i></li> <li>Disappearance of any soft tissue plasmacytomas, <i>and</i></li> <li>&lt;5% PCs in bone marrow</li> <li>No evidence of initial monoclonal protein isotype(s) on immunofixation of the serum and urine.<sup>b</sup></li> </ul>
Very good partial response <sup>a</sup>	<ul style="list-style-type: none"> <li>Serum and urine M-component detectable by immunofixation but not on electrophoresis, <i>or</i></li> <li>≥90% reduction in serum M-component plus urine M-component &lt;100 mg/24 hours</li> </ul>
Partial response	<ul style="list-style-type: none"> <li>≥50% reduction of serum M-protein and reduction in 24-hour urinary M-protein by ≥90% or to &lt; 200 mg/24 hours</li> <li>If serum and urine M-protein are not measurable, a decrease ≥50% in the difference between involved and uninvolved FLC levels is required in place of the M-protein criteria</li> <li>If serum and urine M-protein are not measurable, and serum free light assay is also not measurable, ≥50% reduction in bone marrow PCs is required in place of M-protein, provided baseline percentage was ≥30%</li> <li>In addition to the above criteria, if present at baseline, ≥50% reduction in the size of soft tissue plasmacytomas is also required.</li> </ul>
Minimal response	<ul style="list-style-type: none"> <li>≥25% but ≤49% reduction of serum M-protein and reduction in 24-hour urine M-protein by 50% to 89%</li> <li>In addition to the above criteria, if present at baseline, ≥50% reduction in the size of soft tissue plasmacytomas is also required</li> </ul>
Stable disease	<ul style="list-style-type: none"> <li>Not meeting criteria for sCR, CR, VGPR, PR, MR, or progressive disease</li> </ul>
Progressive disease <sup>c</sup>	<p>Any one or more of the following criteria:</p> <ul style="list-style-type: none"> <li>Increase of 25% from lowest response value in any of the following: <ul style="list-style-type: none"> <li>Serum M-component (absolute increase must be ≥0.5 g/dL), <i>and/or</i></li> <li>Urine M-component (absolute increase must be ≥200 mg/24 hours), <i>and/or</i></li> <li>Only in subjects without measurable serum and urine M-protein levels: the difference between involved and uninvolved FLC levels (absolute increase must be &gt; 10 mg/dL)</li> <li>Only in subjects without measurable serum and urine M-protein levels and without measurable disease by FLC levels, bone marrow PC percentage (absolute increase must be ≥10%)</li> </ul> </li> <li>Appearance of a new lesion(s), ≥50% increase from nadir in SPD of &gt;1 lesion, or ≥50% increase in the longest diameter of a previous lesion &gt;1 cm in short axis</li> <li>Definite development of new bone lesions or definite increase in the size of existing bone lesions</li> <li>≥50% increase in circulating plasma cells (minimum of 200 cells per μL) if this is the only measure of disease</li> </ul>

Key: CR=complete response; FLC=free light chain; PC=plasma cell; PR=partial response; sCR=stringent complete response; SPD=sum of the products of the maximal perpendicular diameters of measured lesions; VGPR=very good partial response.

<sup>a</sup> Clarifications to the criteria for coding CR and VGPR in subjects in whom the only measurable disease is by serum FLC levels: CR in such subjects indicates a normal FLC ratio of 0.26 to 1.65 (or reference range in testing laboratory) in addition to CR criteria listed above. VGPR in such subjects requires a ≥90% decrease in the difference between involved and uninvolved FLC levels. For patients achieving very good partial response by other criteria, a soft tissue plasmacytoma must decrease by more than 90% in the sum of the maximal perpendicular diameter (SPD) compared with baseline.

- b. In some cases it is possible that the original M protein light-chain isotype is still detected on immunofixation but the accompanying heavy-chain component has disappeared; this would not be considered as a CR even though the heavy-chain component is not detectable, since it is possible that the clone evolved to one that secreted only light chains. Thus, if a patient has IgA lambda myeloma, then to qualify as CR there should be no IgA detectable on serum or urine immunofixation; if free lambda is detected without IgA, then it must be accompanied by a different heavy chain isotype (IgG, IgM, etc.).
- c. Clarifications to the criteria for coding progressive disease: bone marrow criteria for progressive disease are to be used only in subjects without measurable disease by M-protein and by FLC levels; “25% increase” refers to M-protein, and FLC, and does not refer to bone lesions, or soft tissue plasmacytomas and the “lowest response value” does not need to be a confirmed value.

Notes: All response categories (CR, sCR, VGPR, PR, MR, and progressive disease) require 2 consecutive assessments made at any time before the institution of any new therapy; CR, sCR, VGPR, PR, MR, and stable disease categories also require no known evidence of progressive or new bone lesions if radiographic studies were performed. VGPR and CR categories require serum and urine studies regardless of whether disease at baseline was measurable on serum, urine, both, or neither.

Radiographic studies are not required to satisfy these response requirements. Bone marrow assessments need not be confirmed. For progressive disease, serum M-component increases of  $\geq 1$  g/dL are sufficient to define relapse if lowest M-component is  $\geq 5$  g/dL.

Source: Adapted from [Durie \(2015\)](#), [Rajkumar \(2011\)](#), [Kumar \(2016\)](#)

**Attachment 2: Cytokine Release Syndrome ASBMT (ASTCT) Consensus Grading System**

Grade	Toxicity
<b>Grade 1</b>	Fever <sup>a</sup> (Temperature $\geq 38^\circ$ )
<b>Grade 2</b>	Fever <sup>a</sup> (Temperature $\geq 38^\circ$ ) with either: <ul style="list-style-type: none"> <li>• Hypotension not requiring vasopressors</li> <li>• And/or<sup>c</sup> hypoxia requiring low-flow nasal cannula<sup>b</sup> or blow-by</li> </ul>
<b>Grade 3</b>	Fever <sup>a</sup> (Temperature $\geq 38^\circ$ ) with either: <ul style="list-style-type: none"> <li>• Hypotension requiring a vasopressor with or without vasopressin.</li> <li>• And/or<sup>c</sup> hypoxia requiring high-flow nasal cannula<sup>b</sup>, facemask, nonrebreather mask, or Venturi mask.</li> </ul>
<b>Grade 4</b>	Fever <sup>a</sup> (Temperature $\geq 38^\circ$ ) with either: <ul style="list-style-type: none"> <li>• Hypotension requiring multiple vasopressors (excluding vasopressin).</li> <li>• And/or<sup>c</sup> hypoxia requiring positive pressure (eg, CPAP, BiPAP, intubation and mechanical ventilation),</li> </ul>
<b>Grade 5</b>	Death

<sup>a</sup> Fever not attributable to any other cause. In patients who have CRS then receive antipyretics or anticytokine therapy such as tocilizumab or steroids, fever is no longer required to grade subsequent CRS severity. In this case, CRS grading is driven by hypotension and/or hypoxia.

<sup>b</sup> Low-flow nasal cannula is defined as oxygen delivered at  $\leq 6$  L/minute or blow-by oxygen delivery. High-flow nasal cannula is defined as oxygen delivered at  $>6$  L/minute.

<sup>c</sup> CRS grade is determined by the more severe event: hypotension or hypoxia not attributable to any other cause.

Note: Organ toxicities associated with CRS may be graded according to CTCAE v5.0 but they do not influence CRS grading.

Source: [Lee \(2019\)](#)

**Attachment 3: Immune Effector Cell-associated Encephalopathy (ICE) Tool**

<b>Immune Effector Cell-Associated Encephalopathy (ICE) Tool<sup>a</sup></b>
<p><b>Orientation:</b> Orientation to year, month, city, hospital:</p> <ul style="list-style-type: none"> <li>• <b>4 points</b></li> </ul> <p><b>Naming:</b> Name 3 objects (e.g., point to clock, pen, button):</p> <ul style="list-style-type: none"> <li>• <b>3 points</b></li> </ul> <p><b>Following commands:</b> (e.g., Show me 2 fingers or Close your eyes and stick out your tongue):</p> <ul style="list-style-type: none"> <li>• <b>1 point</b></li> </ul> <p><b>Writing:</b> Ability to write a standard sentence (e.g., Our national bird is the bald eagle):</p> <ul style="list-style-type: none"> <li>• <b>1 point</b></li> </ul> <p><b>Attention:</b> Count backwards from 100 by ten:</p> <ul style="list-style-type: none"> <li>• <b>1 point</b></li> </ul>
<p>a: ICE-Tool Scoring:</p> <ul style="list-style-type: none"> <li>• Score 10: No impairment</li> <li>• Score 7-9: Grade 1 ICANS</li> <li>• Score 3-6: Grade 2 ICANS</li> <li>• Score 0-2: Grade 3 ICANS</li> <li>• Score 0 due to patient unarousable and unable to perform ICE assessment: Grade 4 ICANS</li> </ul>

**Attachment 4: Immune Effector Cell-associated Neurotoxicity Syndrome (ICANS) ASBMT (ASTCT) Consensus Grading System<sup>a,b</sup>**

Neurotoxicity Domain	Grade 1	Grade 2	Grade 3	Grade 4
<b>ICE Score</b>	7-9	3-6	0-2	0 (patient is unarousable and unable to perform ICE).
<b>Depressed Level of Consciousness</b>	Awakens spontaneously.	Awakens to voice.	Awakens only to tactile stimulus.	Patient is unarousable or requires vigorous or repetitive tactile stimuli to arouse. Stupor or coma.
<b>Seizure</b>	N/A	N/A	Any clinical seizure, focal or generalized, that resolves rapidly; or Non-convulsive seizures on EEG that resolve with intervention.	Life-threatening prolonged seizure (>5 min); or Repetitive clinical or electrical seizures without return to baseline in between.
<b>Motor Findings</b>	N/A	N/A	N/A	Deep focal motor weakness such as hemiparesis or paraparesis.
<b>Raised Intracranial Pressure / Cerebral Edema</b>	N/A	N/A	Focal/local edema on neuroimaging.	Diffuse cerebral edema on neuroimaging; or Decerebrate or decorticate posturing; or Cranial nerve VI palsy; or Papilledema; or Cushing's triad.

a: Toxicity grading according to [Lee et al 2019](#)

b: ICANS grade is determined by the most severe event (ICE score, level of consciousness, seizure, motor findings, raised ICP/cerebral edema) not attributable to any other cause.

Note: all other neurologic adverse events (not associated with ICANS) should continue to be graded with CTCAE Version 5.0 during both phases of the study

**Attachment 5: International Myeloma Working Group Diagnostic Criteria**

Diagnostic criteria for myeloma must be met when the patient was diagnosed. Multiple myeloma is defined as clonal bone marrow plasma cells  $\geq 10\%$  or biopsy-proven bony or extramedullary plasmacytoma<sup>a</sup> and any one or more of the following myeloma defining events:

- Myeloma defining events:
  - Evidence of end organ damage that can be attributed to the underlying plasma cell proliferative disorder, specifically:
    - **C:** Hypercalcemia: serum calcium  $>0.25$  mmol/L ( $>1$  mg/dL) higher than the upper limit of normal or  $>2.75$  mmol/L ( $>11$  mg/dL)
    - **R:** Renal insufficiency: creatinine clearance  $<40$  mL per min<sup>b</sup> or serum creatinine  $>177$   $\mu$ mol/L ( $>2$  mg/dL)
    - **A:** Anemia: hemoglobin value of  $>20$  g/L below the lower limit of normal, or a hemoglobin value  $<100$  g/L
    - **B:** Bone lesions: one or more osteolytic lesions on skeletal radiography, CT, or PET-CT<sup>c,d</sup>
  - Any one or more of the following biomarkers of malignancy:
    - Clonal bone marrow plasma cell percentage<sup>a</sup>  $\geq 60\%$
    - Involved:uninvolved serum free light chain ratio<sup>e</sup>  $\geq 100$
    - $>1$  focal lesions on MRI studies<sup>f</sup>

<sup>a</sup> Clonality should be established by showing  $\kappa/\lambda$ -light-chain restriction on flow cytometry, immunohistochemistry, or immunofluorescence. Bone marrow plasma cell percentage should preferably be estimated from a core biopsy specimen; in case of a disparity between the aspirate and core biopsy, the highest value should be used.

<sup>b</sup> Measured or estimated by validated equations.

<sup>c</sup> If bone marrow has less than 10% clonal plasma cells, more than one bone lesion is required to distinguish from solitary plasmacytoma with minimal marrow involvement.

<sup>d</sup> PET-CT=<sup>18</sup>F-fluorodeoxyglucose PET with CT.

<sup>e</sup> These values are based on the serum Freelite assay (The Binding Site Group, Birmingham, UK). The involved free light chain must be  $\geq 100$  mg/L.

<sup>f</sup> Each focal lesion must be 5 mm or more in size.

Source: [Rajkumar \(2011\)](#)

**Attachment 6: Prior Cancer Therapy for Multiple Myeloma**

A line of therapy is defined as one or more cycles of a planned treatment program. This may consist of one or more planned cycles of single-agent therapy or combination therapy, as well as a sequence of treatments administered in a planned manner. For example, a planned treatment approach of induction therapy followed by autologous stem cell transplantation, followed by maintenance is considered one line of therapy. A new line of therapy starts when a planned course of therapy is modified to include other treatment agents (alone or in combination) as a result of disease progression, relapse, or toxicity. A new line of therapy also starts when a planned period of observation off therapy is interrupted by a need for additional treatment for the disease.

Source: [Rajkumar \(2011\)](#)

**Attachment 7: Eastern Cooperative Oncology Group Performance Status Grade**

<b>Grade</b>	<b>Eastern Cooperative Oncology Group Performance Status</b>
0	Fully active, able to carry on all pre-disease performance without restriction
1	Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature, eg, light house work, office work
2	Ambulatory and capable of all self-care but unable to carry out any work activities. Up and about more than 50% of waking hours
3	Capable of only limited self-care, confined to bed or chair more than 50% of waking hours
4	Completely disabled. Cannot carry on any self-care. Totally confined to bed or chair
5	Dead

Source: Eastern Cooperative Oncology Group, Robert Comis M.D., Group Chair ([Oken 1982](#)).



**Attachment 8: Formulas for Estimating Glomerular Filtration Rate*****Modified Diet in Renal Disease (MDRD) Formula***

For serum creatinine in **mg/dL**, the estimated glomerular filtration rate (eGFR) for the MDRD formula is:

$$\text{eGFR (MDRD) mL/min per } 1.73\text{m}^2 = 175 \times [\text{serum creatinine (mg/dL)}]^{-1.154} \times [\text{age}]^{-0.203} \times [1.212 \text{ if black}] \times [0.742 \text{ if female}]$$

For serum creatinine in **μmol/L**, the eGFR for the MDRD formula is:

$$\text{eGFR (MDRD) mL/min per } 1.73\text{m}^2 = 175 \times [\text{serum creatinine } (\mu\text{mol/L})/88.4]^{-1.154} \times [\text{age}]^{-0.203} \times [1.212 \text{ if black}] \times [0.742 \text{ if female}]$$

Source: [Levey \(2006\)](#)

**Attachment 9: Serum Calcium Corrected for Albumin**

If calcium is expressed in mg/dL and albumin is expressed in g/dL:

Corrected calcium (mg/dL) =

$$\text{Serum calcium (mg/dL)} + 0.8 \times (4 - \text{serum albumin [g/dL]})$$

If calcium is expressed in mmol/L and albumin is expressed in g/L:

Corrected calcium (mmol/L) =

$$\text{Serum calcium (mmol/L)} + 0.02 \times (40 - \text{serum albumin [g/L]})$$

Source: [Burtis \(1998\)](#)

**Attachment 10: Hepatitis B Virus Screening**

The following hepatitis B virus screening guide is to be used to determine subject eligibility for the study:

<b>Eligibility based on hepatitis B virus test results</b>			
<b>Action</b>	<b>Hepatitis B test result</b>		
	<b>Hepatitis B surface antigen (HBsAg)</b>	<b>Hepatitis B surface antibody (anti-HBs)</b>	<b>Hepatitis B core antibody (anti-HBc)</b>
<b>Exclude</b>	+	— <i>or</i> +	— <i>or</i> +
<b>Include</b>	—	—	—
	—	+ <sup>*#</sup>	+ <sup>#</sup>
	—	—	+ <sup>#</sup>
	—	+ <sup>*</sup>	—

\* Subjects who are anti-HBs positive and without history of vaccination, should have HBV-DNA quantification test. Subjects with positive HBV-DNA should be excluded. Subjects with negative HBV-DNA can be enrolled. If required by local country guidelines on HBV prevention, HBV-DNA and AST/ALT laboratories should be performed every 3 months for the first 12 months after JNJ-68284528 administration. If there is evidence of HBV reactivation initiate treatment as appropriate per institutional guidance.

# Subjects with positive anti-HBc and either positive or negative anti-HBs should have HBV-DNA quantification test. Subjects with positive HBV-DNA should be excluded. Subjects with negative HBV-DNA can be enrolled; however, HBV-DNA and AST/ALT laboratories should be performed every 3 months for the first 12 months after JNJ-68284528 administration. If there is evidence of HBV reactivation initiate treatment as appropriate per institutional guidance.

**Attachment 11: Patient Reported Outcomes (PRO) Measures****EORTC QLQ-C30****EORTC QLQ-C30 (version 3)**

We are interested in some things about you and your health. Please answer all of the questions yourself by circling the number that best applies to you. There are no "right" or "wrong" answers. The information that you provide will remain strictly confidential.

Please fill in your initials:

--	--	--	--	--

Your birthdate (Day, Month, Year):

--	--	--	--	--	--	--	--	--	--

Today's date (Day, Month, Year):

31 

--	--	--	--	--	--	--	--	--	--

	Not at All	A Little	Quite a Bit	Very Much
1. Do you have any trouble doing strenuous activities, like carrying a heavy shopping bag or a suitcase?	1	2	3	4
2. Do you have any trouble taking a <u>long</u> walk?	1	2	3	4
3. Do you have any trouble taking a <u>short</u> walk outside of the house?	1	2	3	4
4. Do you need to stay in bed or a chair during the day?	1	2	3	4
5. Do you need help with eating, dressing, washing yourself or using the toilet?	1	2	3	4

**During the past week:**

	Not at All	A Little	Quite a Bit	Very Much
6. Were you limited in doing either your work or other daily activities?	1	2	3	4
7. Were you limited in pursuing your hobbies or other leisure time activities?	1	2	3	4
8. Were you short of breath?	1	2	3	4
9. Have you had pain?	1	2	3	4
10. Did you need to rest?	1	2	3	4
11. Have you had trouble sleeping?	1	2	3	4
12. Have you felt weak?	1	2	3	4
13. Have you lacked appetite?	1	2	3	4
14. Have you felt nauseated?	1	2	3	4
15. Have you vomited?	1	2	3	4
16. Have you been constipated?	1	2	3	4

Please go on to the next page



MySIm Q**MULTIPLE MYELOMA SYMPTOM AND IMPACT  
QUESTIONNAIRE**

Instructions: The purpose of this questionnaire is to collect information about your experience with multiple myeloma. Please only consider your experiences related to your multiple myeloma when answering the following questions.

***People often experience changes in the severity of their symptoms from day to day or within a single day. When answering the next 4 questions, please only think about the time within the past 7 days when each symptom was at its worst.***

1. How would you rate the worst **pain in your back** within the past 7 days?

- No pain
- A little pain
- Moderate pain
- Quite a bit of pain
- Severe pain

2. How would you rate the worst **pain in your legs** within the past 7 days?

- No pain
- A little pain
- Moderate pain
- Quite a bit of pain
- Severe pain

3. How would you rate the **worst pain in areas other than your back or legs** within the past 7 days?

- No pain
- A little pain
- Moderate pain
- Quite a bit of pain
- Severe pain

4. How would you rate the **worst numbness or tingling in your hands or feet** within the past 7 days?

- No numbness or tingling
- A little numbness or tingling
- Moderate numbness or tingling
- Quite a bit of numbness or tingling
- Severe numbness or tingling

**For each question, select only 1 answer that best describes how often you experienced each issue within the past 7 days.**

5. How much did your **pain interfere** with your usual or daily activities within the past 7 days?

- Not at all
- A little bit
- Moderately
- Quite a bit
- Very much

6. How often did you have **low energy** within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

7. How often did you **tire easily** (for example, needing to rest during activities) within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

8. How often did you experience **muscle weakness** within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always



9. How often did you have **trouble with your sleep** (for example, difficulty falling asleep or staying sleep) within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

10. How often did you have a **poor appetite** within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

11. How often did you have **difficulty with your memory** within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

12. How often did you have **difficulty concentrating** on things (for example, reading a book or following a conversation) within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

13. How often were you **limited in doing your daily activities** within the past 7 days (for example, struggling or needing help with work or house chores)?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

14. How often did you have **difficulty walking** within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

15. How often were you **limited in your social life** (for example, activities with your friends or family) within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

***For each question, select only 1 answer that best describes how you felt within the past 7 days.***

16. How often have you felt **depressed about your multiple myeloma** within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

17. How often did you **worry that your multiple myeloma could get worse** within the past 7 days?

- Never
- Rarely
- Some of the time
- Most of the time
- Always

Patient Global Impression of Severity (PGIS)

**Patient's Global Impression of Severity (PGIS)  
Pain**

Overall, how would you rate the severity of your pain currently? (Please select one response)

- 1. None
- 2. Mild
- 3. Moderate
- 4. Severe
- 5. Very Severe

SPECIMEN

Patient Global Impression of Change (PGIC)

**Patient's Global Impression of Change (PGIC)  
of Overall Health**

Compared to before you received the CAR-T infusion in this study, how has your overall health changed? (Please select one response)

- 1. A lot better now
- 2. Moderately better now
- 3. A little better now
- 4. Neither better, nor worse (no change)
- 5. A little worse now
- 6. Moderately worse now
- 7. A lot worse now

SPECIMEN

## NCI PRO-CTCAE Items

**NCI PRO-CTCAE™ ITEMS**

Item Library Version 1.0

English

Form created on 6 September 2018

As individuals go through treatment for their cancer they sometimes experience different symptoms and side effects. For each question, please check or mark an  in the one box that best describes your experiences over the past 7 days...

1.	In the last 7 days, how OFTEN did you have NAUSEA?				
	<input type="radio"/> Never	<input type="radio"/> Rarely	<input type="radio"/> Occasionally	<input type="radio"/> Frequently	<input type="radio"/> Almost constantly
	In the last 7 days, what was the SEVERITY of your NAUSEA at its WORST?				
	<input type="radio"/> None	<input type="radio"/> Mild	<input type="radio"/> Moderate	<input type="radio"/> Severe	<input type="radio"/> Very severe

2.	In the last 7 days, how OFTEN did you have VOMITING?				
	<input type="radio"/> Never	<input type="radio"/> Rarely	<input type="radio"/> Occasionally	<input type="radio"/> Frequently	<input type="radio"/> Almost constantly
	In the last 7 days, what was the SEVERITY of your VOMITING at its WORST?				
	<input type="radio"/> None	<input type="radio"/> Mild	<input type="radio"/> Moderate	<input type="radio"/> Severe	<input type="radio"/> Very severe

3.	In the last 7 days, how OFTEN did you have LOOSE OR WATERY STOOLS (DIARRHEA/DIARRHOEA)?				
	<input type="radio"/> Never	<input type="radio"/> Rarely	<input type="radio"/> Occasionally	<input type="radio"/> Frequently	<input type="radio"/> Almost constantly

4.	In the last 7 days, what was the SEVERITY of your SHORTNESS OF BREATH at its WORST?				
	<input type="radio"/> None	<input type="radio"/> Mild	<input type="radio"/> Moderate	<input type="radio"/> Severe	<input type="radio"/> Very severe
	In the last 7 days, how much did your SHORTNESS OF BREATH INTERFERE with your usual or daily activities?				
	<input type="radio"/> Not at all	<input type="radio"/> A little bit	<input type="radio"/> Somewhat	<input type="radio"/> Quite a bit	<input type="radio"/> Very much

5.	In the last 7 days, did you have any RASH?	
	<input type="radio"/> Yes	<input type="radio"/> No

6.	In the last 7 days, what was the SEVERITY of your DIZZINESS at its WORST?				
	<input type="radio"/> None	<input type="radio"/> Mild	<input type="radio"/> Moderate	<input type="radio"/> Severe	<input type="radio"/> Very severe
	In the last 7 days, how much did DIZZINESS INTERFERE with your usual or daily activities?				
	<input type="radio"/> Not at all	<input type="radio"/> A little bit	<input type="radio"/> Somewhat	<input type="radio"/> Quite a bit	<input type="radio"/> Very much

The PRO-CTCAE™ items and information herein were developed by the NATIONAL CANCER INSTITUTE at the NATIONAL INSTITUTES OF HEALTH, in Bethesda, Maryland, U.S.A. Use of the PRO-CTCAE™ is subject to NCI's Terms of Use.



**NCI PRO-CTCAE™ ITEMS****Item Library Version 1.0****English****Form created on 6 September 2018**

7.	In the last 7 days, how OFTEN did you have a HEADACHE?				
	<input type="radio"/> Never	<input type="radio"/> Rarely	<input type="radio"/> Occasionally	<input type="radio"/> Frequently	<input type="radio"/> Almost constantly
	In the last 7 days, what was the SEVERITY of your HEADACHE at its WORST?				
	<input type="radio"/> None	<input type="radio"/> Mild	<input type="radio"/> Moderate	<input type="radio"/> Severe	<input type="radio"/> Very severe
	In the last 7 days, how much did your HEADACHE INTERFERE with your usual or daily activities?				
	<input type="radio"/> Not at all	<input type="radio"/> A little bit	<input type="radio"/> Somewhat	<input type="radio"/> Quite a bit	<input type="radio"/> Very much

8.	In the last 7 days, what was the SEVERITY of your FATIGUE, TIREDNESS, OR LACK OF ENERGY at its WORST?				
	<input type="radio"/> None	<input type="radio"/> Mild	<input type="radio"/> Moderate	<input type="radio"/> Severe	<input type="radio"/> Very severe
	In the last 7 days, how much did FATIGUE, TIREDNESS, OR LACK OF ENERGY INTERFERE with your usual or daily activities?				
	<input type="radio"/> Not at all	<input type="radio"/> A little bit	<input type="radio"/> Somewhat	<input type="radio"/> Quite a bit	<input type="radio"/> Very much

The PRO-CTCAE™ items and information herein were developed by the NATIONAL CANCER INSTITUTE at the NATIONAL INSTITUTES OF HEALTH, in Bethesda, Maryland, U.S.A. Use of the PRO-CTCAE™ is subject to NCI's Terms of Use.

**Attachment 12: Conversion Table for Glucocorticosteroid Dose**

Glucocorticoid	Approximate Equivalent Dose (mg)	Half-life (Biologic) hours
Intermediate-Acting		
Methylprednisolone	4	18-36
Prednisolone	5	18-36
Prednisone	5	18-36
Triamcinolone	4	18-36
Long-Acting		
Betamethasone	0.6 – 0.75	36-54
Dexamethasone	0.75	36-54



**Attachment 13: Anticipated Adverse Events**

Attachment Deleted per Amendment 3

**Attachment 14: JNJ-68284528 Outpatient Administration Guidelines**

When evaluating the suitability for outpatient administration, if allowed by local regulations and institutional guidance among other considerations, investigators should assess the subject's clinical status and the health care facility capability to safely manage outpatient logistics. General recommendations for each of these considerations are provided below; sponsor approval is required:

**1. Clinical consideration**

General guidance for clinical considerations for a subject that is suitable for outpatient administration and follow-up includes the following:

- Not requiring packed red blood cell or platelet transfusions more frequently than every 2 days
- No presence of an indwelling central line (with the exception of a PICC line) given risk of infection in the setting of cytopenia
- No fever or active infection (bacterial, fungal, viral) since study enrollment
- No Grade 3 or higher non-hematologic toxicities of cyclophosphamide and fludarabine including nausea, vomiting, and diarrhea
- No clinically significant coagulopathy that would increase the risk of bleeding in the setting of cytopenia
- No high tumor burden defined as at least 60% plasma cell infiltration of the marrow and/or the presence of extramedullary disease
- No risk factors for developing clinically significant tumor lysis syndrome and requiring management with increased hydration, allopurinol, or rasburicase. Patients who are receiving prophylactic treatment for TLS are eligible for outpatient infusion, if deemed stable by the investigator
- No rapidly progressing disease
- No deterioration in neurologic status, including mental status changes such as confusion or increased somnolence. The only exception is confusion or somnolence that has resolved and must be attributed to diphenhydramine premedication for JNJ-68284528.
- The following laboratory parameters:
  - Estimated glomerular filtration rate of  $\geq 40$  mL/min/1.73 m<sup>2</sup>
  - AST and ALT  $\leq 3$  times the upper limit of normal

**2. Logistical consideration for qualified healthcare facility**

Outpatient administration and post-JNJ-68284528 infusion follow-up must take place at a qualified healthcare facility.

The following should be considered for outpatient administration and follow-up until Day 4 prior to inpatient admission from Day 5 to Day 14 of JNJ-68284528 infusion:

- Site must discuss with subjects about how to recognize the presenting signs and symptoms of CAR-T associated toxicities (including but not limited to CRS, neurotoxicities, infections, etc.) as presented in the patient wallet card

- Site must provide patients with educational material including but not limited to emergency contact information
- Subject will receive daily phone call follow-ups from the hospital site staff (as required by the Time and Events Schedule and [Attachment 15](#)) during typical business hours
- Subject is required to stay within 30 minutes of transportation to the hospital and remain in the company of a competent adult at all times until the time of readmission on Day 5 after JNJ 68284528 infusion
- Subject must comply with all the protocol requirement procedures, including measuring and recording of body temperature twice per day, and coming to the site for safety assessments according to the Time and Events Schedule ([Table 1](#), [Table 3](#), [Table 5](#), and [Table 6](#)).
- Admission to the hospital is required at any time in the event of any presenting signs and symptoms of CRS and/or neurotoxicity even if these occur before Day 5. Even without symptoms of CRS or neurotoxicity, subject will be admitted for inpatient monitoring from Day 5 to Day 14 of JNJ-68284528 infusion
- If a subject does not develop symptoms of CRS or neurotoxicity or other clinically significant adverse event until Day 10 post JNJ-68284528 infusion, subject may be discharged with daily outpatient phone call follow-ups during business hours through study Day 14. Upon discharge from the hospital, the subject must stay locally within 1 hour of transportation to the hospital and remain in the company of a competent adult at all times for 1 additional week, or up to study Day 21, whichever is sooner
- Subjects that experience CRS or neurotoxicity, can be discharged from the hospital when they are afebrile for 24 hours and signs and symptoms of CRS and/or neurotoxicity or other clinically significant adverse event have resolved

**Attachment 15: Monitoring for Subjects Receiving JNJ-68284528 as Outpatient**

Subjects eligible to receive JNJ-68284528 as an outpatient (see recommendations in [Attachment 14](#)) in consultation with and approval of the sponsor

<b>Day 1</b>	JNJ-68284528 infusion
<b>Day 1 to 4</b>	<ul style="list-style-type: none"> <li>• Subject is required to stay within 30 minutes of transportation to the hospital and remain in the company of a competent adult at all times</li> <li>• Subject will receive daily phone call follow-ups from hospital staff during typical business hours</li> <li>• Admission to the hospital is required at any time in the event of any presenting signs and symptoms of CRS and/or neurotoxicity</li> </ul>
<b>Day 5 to 14</b>	<ul style="list-style-type: none"> <li>• Required inpatient admission</li> <li>• Potential discharge on Day 10 for subjects who do not develop symptoms of CRS, neurotoxicity, or other significant adverse events <ul style="list-style-type: none"> <li>– Subjects discharged on Day 10 will receive daily phone call follow-ups from the hospital staff during typical business hours through Day 14</li> </ul> </li> </ul>

**Attachment 16: Contraceptive and Barrier Guidance and Collection of Pregnancy Information****Woman of Childbearing Potential (WOCBP)**

A woman is considered fertile following menarche and until becoming postmenopausal unless permanently sterile (see below).

**Woman Not of Childbearing Potential*****Premenarchal***

A premenarchal state is one in which menarche has not yet occurred.

***Postmenopausal***

A postmenopausal state is defined as no menses for 24 months without an alternative medical cause. A high follicle stimulating hormone (FSH) level (>40 IU/L or mIU/mL) in the postmenopausal range may be used to confirm a postmenopausal state in women not using hormonal contraception or hormonal replacement therapy (HRT), however in the absence of 12 months of amenorrhea, a single FSH measurement is insufficient. If there is a question about menopausal status in women on HRT, the woman will be required to use one of the non-estrogen-containing hormonal highly effective contraceptive methods if she wishes to continue HRT during the study.

***Permanently sterile***

Permanent sterilization methods include hysterectomy, bilateral salpingectomy, bilateral tubal occlusion/ligation procedures, and bilateral oophorectomy.

Note: If the childbearing potential changes after start of the study (eg, a premenarchal woman experiences menarche) or the risk of pregnancy changes (eg, a woman who is not heterosexually active becomes active), a woman must begin a highly effective method of contraception, as described throughout the inclusion criteria.

If reproductive status is questionable, additional evaluation should be considered.

Contraceptive (birth control) use by men or women should be consistent with local regulations regarding the acceptable methods of contraception for those participating in clinical studies.

Typical use failure rates may differ from those when used consistently and correctly. Use should be consistent with local regulations regarding the use of contraceptive methods for subjects in clinical studies.

Because of the embryo-fetal risk of lenalidomide, all subjects treated with D-VRd or D+R must be enrolled in the lenalidomide pregnancy prevention program applicable in their region. Investigators should comply with the lenalidomide Global Pregnancy Prevention Plan or with the respective country-specific REVLIMID<sup>®</sup> / Lenalidomide Risk Minimization Program (ie, Pregnancy prevention program), whichever is more stringent, as implemented in the post-marketing setting and ensure that all subjects adhere to these programs. When no REVLIMID<sup>®</sup> /

Lenalidomide Risk Minimization Program exists, subjects must adhere to the Global lenalidomide Pregnancy Prevention Plan.

### Examples of Contraceptives

#### **EXAMPLES OF CONTRACEPTIVES<sup>a</sup> ALLOWED DURING THE STUDY INCLUDE: USER INDEPENDENT**

**Highly Effective Methods That Are User Independent** *Failure rate of <1% per year when used consistently and correctly.*

- Implantable progestogen-only hormone contraception associated with inhibition of ovulation<sup>b</sup>
- Intrauterine device (IUD)
- Intrauterine hormone-releasing system (IUS)
- Bilateral tubal occlusion
- Vasectomized partner  
*(Vasectomized partner is a highly effective contraceptive method provided that the partner is the sole sexual partner of the woman of childbearing potential and the absence of sperm has been confirmed. If not, additional highly effective method of contraception should be used. Spermatogenesis cycle is approximately 74 days.)*

#### **USER DEPENDENT**

**Highly Effective Methods That Are User Dependent** *Failure rate of <1% per year when used consistently and correctly.*

- Progestogen-only hormone contraception associated with inhibition of ovulation<sup>b</sup>
  - oral
  - injectable
- Sexual abstinence  
*(Sexual abstinence is considered a highly effective method only if defined as refraining from heterosexual intercourse during the entire period of risk associated with the study intervention. The reliability of sexual abstinence needs to be evaluated in relation to the duration of the study and the preferred and usual lifestyle of the subject.)*

#### **NOT ALLOWED AS SOLE METHOD OF CONTRACEPTION DURING THE STUDY (not considered to be highly effective - failure rate of $\geq 1\%$ per year)**

- Progestogen-only oral hormonal contraception where inhibition of ovulation is not the primary mode of action.
  - Male or female condom with or without spermicide<sup>c</sup>
  - Cap, diaphragm, or sponge with spermicide
  - A combination of male condom with either cap, diaphragm, or sponge with spermicide (double-barrier methods)<sup>c</sup>
  - Periodic abstinence (calendar, symptothermal, post-ovulation methods)
  - Withdrawal (coitus-interruptus)
  - Spermicides alone
  - Lactational amenorrhea method (LAM)
- a) Typical use failure rates may differ from those when used consistently and correctly. Use should be consistent with local regulations regarding the use of contraceptive methods for subjects in clinical studies.
- b) Hormonal contraception may be susceptible to interaction with the study intervention, which may reduce the efficacy of the contraceptive method. In addition, consider if the hormonal contraception may interact with the study intervention.
- c) Male condom and female condom should not be used together (due to risk of failure with friction).

**Pregnancy During the Study**

All initial reports of pregnancy in female subjects or partners of male subjects must be reported to the sponsor by the study-site personnel within 24 hours of their knowledge of the event using appropriate pregnancy notification form. Abnormal pregnancy outcomes (eg, spontaneous abortion, fetal death, stillbirth, congenital anomalies, and ectopic pregnancy) are considered serious adverse events and must be reported using the Serious Adverse Event Form. Any subject who becomes pregnant during the study must discontinue further study treatment. Because the effect of the study treatment on sperm is unknown, pregnancies in partners of male subjects included in the study will be reported as noted above. Follow-up information regarding the outcome of the pregnancy and any postnatal sequelae in the infant will be required.

**Attachment 17: Handwriting Adverse Event Toxicity Grading Criteria**

<b>Adverse Event Term</b>	<b>Grade 1</b>	<b>Grade 2</b>
<b>Micrographia:</b> abnormally small or cramped handwriting	Mildly smaller letters or reduced spacing (eg, <50% decrease from baseline)	Moderate to severely smaller letters or reduced spacing (eg, >=50% decrease from baseline)
<b>Dysgraphia:</b> illegible writing or writing that takes an unusually long time or great effort	Mildly slower writing, impaired straightness of line, difficulty in completing task from baseline; most words are legible	Moderate to severely slower writing, impaired straightness of line, difficulty in completing task from baseline; most words are illegible
<b>Agraphia:</b> pathologic loss of the ability to write	Able to write part of a sentence (3 or more words); noted change from baseline	Able to write just 1 to 2 words, or unable to write any words; noted change from baseline



**Attachment 18: Anti-microbial Prophylaxis Recommendations**

Subjects should receive antimicrobial prophylaxis as per recommendations below or per institutional standards.

Prophylaxis	Therapy	Start	Stop
<b>Anti-Bacterial</b>	Fluoroquinolones (Levofloxacin - 500 mg PO or IV daily, or equivalent)  <i>Suggested Alternative for subjects with allergy to quinolones:</i> Cefpodoxime - 200 mg PO twice a day	At neutropenia onset (ANC < 500/ $\mu$ L)	At Neutropenia resolution (for example, ANC $\geq$ 500/ $\mu$ L)
<b>Anti-Fungal</b>	Fluconazole - 400 mg daily (or equivalent)  <i>Alternatives:</i> Caspofungin or Micafungin  Prolonged neutropenia > 3 weeks – Consider switching to Posaconazole, <i>or</i> as per institutional guidelines	At neutropenia onset (ANC < 500/ $\mu$ L)	At Neutropenia resolution (for example, ANC $\geq$ 500/ $\mu$ L)
<b>Anti-Viral</b>	Acyclovir – 400 - 800 mg PO twice a day (dose to be adjusted as per institutional guidelines)  <i>Alternative:</i> Valacyclovir - 500 mg PO twice a day	Day 1 of infusion	Suggested for at least 12 months post infusion
<b><i>Pneumocystis</i> Pneumonia (PCP)</b>	Trimethoprim-sulfamethoxazole – 1 DS tablet PO daily or 1 SS tablet PO daily  <i>Alternatives:</i> Pentamidine (as per institutional guidelines), <i>or</i> Dapsone – 100 mg PO daily or 50 mg PO BID, <i>or</i> Atovaquone – 1500 mg PO daily	Day 1 of infusion	Suggested duration: 6 months OR until CD4 count $\geq$ 200 cells/ $\mu$ L, (whichever is longer)

**Attachment 19: International Staging System (ISS)**

Stage	Criteria
I	Serum $\beta$ 2-microglobulin <3.5 mg/L Serum albumin $\geq$ 3.5 g/dL
II	Not stage I or III*
III	Serum $\beta$ 2-microglobulin $\geq$ 5.5 mg/L

\*There are 2 categories for stage II: serum  $\beta$ 2-microglobulin <3.5 mg/L but serum albumin <3.5 g/dL; or serum  $\beta$ 2-microglobulin 3.5 to <5.5 mg/L irrespective of the serum albumin level.

Source: Adapted from [Greipp \(2005\)](#)

**Attachment 20: COVID-19 Guidance on Study Conduct****GUIDANCE ON STUDY CONDUCT DURING THE COVID-19 PANDEMIC**

It is recognized that the Coronavirus Disease 2019 (COVID-19) pandemic may have an impact on the conduct of this clinical study due to, for example, self-isolation/quarantine by participants and study-site personnel; travel restrictions/limited access to public places, including hospitals; study site personnel being reassigned to critical tasks.

In alignment with recent health authority guidance, the sponsor is providing options for study related participant management in the event of disruption to the conduct of the study. This guidance does not supersede any local or government requirements or the clinical judgement of the investigator to protect the health and well-being of participants and site staff.

Scheduled visits that cannot be conducted in person at the study site will be performed to the extent possible remotely/virtually or delayed until such time that on-site visits can be resumed. At each contact, participants will be interviewed to collect safety data. Key efficacy endpoint assessments should be performed if required and as feasible. Participants will also be questioned regarding general health status to fulfill any physical examination requirement.

Every effort should be made to adhere to protocol-specified assessments for participants on study intervention, including follow up. Modifications to protocol-required assessments may be permitted via COVID-19 Appendix after consultation with the participant, investigator, and the sponsor. Missed assessments/visits will be captured in the clinical trial management system for protocol deviations. Discontinuations of study interventions and withdrawal from the study should be documented with the prefix “COVID-19-related” in the case report form (CRF).

The sponsor will continue to monitor the conduct and progress of the clinical study, and any changes will be communicated to the sites and to the health authorities according to local guidance. If a participant has tested positive for COVID 19, the investigator should contact the sponsor’s responsible medical officer to discuss plans for study intervention and follow-up. Modifications made to the study conduct as a result of the COVID-19 pandemic should be summarized in the clinical study report.

---

**GUIDANCE SPECIFIC TO THIS PROTOCOL:**

These emergency provisions are meant to ensure subject safety on study while site capabilities are compromised by COVID-19-related restrictions. As restrictions are lifted and the acute phase of the COVID-19 pandemic resolves, sites should revert to original protocol conduct as soon as feasible.

**Study Visits and Assessments**

- At the discretion of the investigator and with sponsor approval, study visits may be performed remotely via telemedicine technology that connects study subjects to their research coordinators and investigators, if not using the sponsor's telemedicine solution. Blood sample collection may be performed at the subject's home by mobile study personnel (ie, nurses and mobile phlebotomist), if not using the sponsor's home health nursing solution. Home health nursing can be done via site contract with a visiting nurse service independent of the sponsor, if not using the sponsor's home health nursing solution. Patient reported outcome questionnaires can be completed by telephone or during the home health visit.
- For subjects who are unable to come to the site for scheduled visits and/or if site capabilities are compromised by COVID-19 related restrictions, contact (eg, telephone, videoconference, or other channels) with the subject should be made in advance, to collect information on the subject's current health status and any new or ongoing adverse events and concomitant medications. Normal study procedures should be followed for the applicable visits as closely as possible even if lab assessments and physical exams are performed locally. Where local laboratories are used, it is important to ensure appropriate documentation of laboratory reference ranges. The remote contact with the subject (as allowable per local regulations), the local laboratory results, and the sponsor discussion should be documented in the subject source record. Similarly, at a minimum, a comment must be entered in the Comments eCRF clearly designating as "COVID-19-related" and acknowledging the discussion between the investigator and the sponsor.
- For subjects eligible to receive retreatment, original protocol conduct must be followed until Day 100 after the CAR-T re-administration.
- All deviations from protocol-required assessments should be documented in detail within the subject's source record and should be clearly designated as "COVID-19-related". It must be documented if a visit is conducted remotely. Source documentation should detail how each assessment was collected (eg, remote vs. on-site, central vs. local laboratory, vital signs taken at home by caretaker vs. delegated in-home nursing).
- Consenting and re-consenting of subjects will be performed as applicable for the measures taken (including also remote consenting by phone or video consultation) and according to local guidance for informed consent applicable during the COVID-19 pandemic.

The above measures are recommended for consideration on a temporary basis during the COVID-19 pandemic to ensure that subject assessments continue as outlined in the protocol without imposing health risk to subjects, their families, and site staff. Every effort should be made to complete all protocol-required assessments.

**Attachment 21: Myeloma Frailty Score Calculator****MYELOMA FRAILITY SCORE CALCULATOR**

Developed by International Myeloma Working Group for the prognosis of elderly myeloma patients.

The score system (range 0-5), based on age, comorbidities, cognitive and physical conditions, developed by Palumbo A. et al<sup>1</sup>, identifies 3 groups of patients:

- fit (score=0)
- intermediate-fitness (score=1)
- frail (score $\geq$ 2)

The 3-year overall survival was 84% in fit patients, 76% in intermediate-fitness patients (HR 1.61, 95%CI 1.02-2.56, p=0.042) and 57% in frail patients (HR 3.57 CI 95% 2.37-5.39, p<0.001). The cumulative incidence of grade  $\geq$ 3 non-hematologic adverse events at 12 months was 22.2% in fit, 26.4% in intermediate-fitness (HR 1.23, 95%CI 0.89-1.71; p 0.217) and 34.0% (HR 1.74, 95%CI 1.28-2.38; p<0.001) in frail patients. The cumulative incidence of treatment discontinuation at 12 months was 16.5% in fit, 20.8% in intermediate-fitness (HR 1.41, 95%CI 1.00-2.01, p=0.052) and 31.2% (HR 2.21, 95%CI 1.57-3.09; p<0.001) in frail patients.

This frailty score predicts mortality and the risk of toxicity in elderly myeloma patients. The International Myeloma Working group proposes this score for the measurement of frailty in the treatment decision-making process and in designing future clinical trials.

<sup>1</sup>Palumbo A, Bringhen S, Mateos MV, et al. Geriatric assessment predicts survival and toxicities in elderly myeloma patients: an International Myeloma Working Group report. *Blood*. 2015 Mar 26;125(13):2068-7  
Supported by FO.NE.SA onlus

**Attachment 22: Characterization of Imaging Response****via PET/CT**

complete response: normalization of abnormal metabolic activity due to multiple myeloma disease activity

partial response:  $\geq 50\%$  decrease in sum of products of the longest perpendicular dimensions of dominant lesions

stable disease: not meeting criteria for partial response or progressive disease

progressive disease:  $\geq 50\%$  increase in product of longest perpendicular diameters of dominant lesions and/or new lesions

**via Whole Body MRI**

complete response: bone marrow highly likely to be responding and/or for soft-tissue disease, RECIST version 1.1 criteria for CR

partial response: bone marrow likely to be responding and/or for soft-tissue disease, RECIST version 1.1 criteria meeting requirements for PR

stable disease: not meeting criteria for partial response or progressive disease

progressive disease: bone marrow highly likely to be progressing and/or soft-tissue disease, RECIST version 1.1 criteria meeting requirements for PD

**References:**

[Cavo 2017](#)  
[Cheson 2007](#)  
[Eisenhauer 2009](#)  
[Messiou 2019](#)

**Attachment 23: Recommendations for the Reporting of Imaging Results****At Baseline**

A radiological report on whole-body imaging in patients with monoclonal plasma cell disorders should include:

- A. Infiltration and bone destruction pattern:
  - a. Minimal (normal appearing)
  - b. Focal lesions
  - c. Diffuse infiltration and bone destruction
  - d. Mixed (focal lesions on diffuse background)
- B. Absolute number of focal lesions:
  - a. For whole body MRI: 0, 1, 2–7, or >7
  - b. For PET/CT: 0, 1–3, or >3
- C. Number of fractures (new vs old, location, and likelihood of malignant vs benign cause)
- D. Extramedullary disease
- E. Soft tissue masses growing out of the bone marrow into the surrounding tissue
- F. Infiltration of the long bones
- G. Evidence of surgical procedures at the skeletal system
- H. Incidental findings

**In Remission**

Differentiate these findings with regards to response to therapy in imaging (guidelines papers for PET/CT and whole body MRI):

- A. Response
  - a. Normalization of bone marrow signal in previously affected areas
  - b. Decrease in the number and size of focal lesions
  - c. Resolution of severely infiltrated bone marrow infiltrate into focal lesions
  - d. Decrease in the of number and size of soft tissue tumors (paramedullary and extramedullary)
- B. No change
- C. Progression
  - a. Worsening of diffuse bone marrow signal or new appearance of infiltration in previously unaffected areas
  - b. Increase in the number and size of focal lesions
  - c. Merging of focal lesions into severely infiltrated bone marrow
  - d. Increase in the size or number of soft tissue tumors (paramedullary and extramedullary)
- D. Cystic or liquid transformation of focal lesions after therapy (for Whole Body MRI only)

[Hillengass 2019](#)

**Attachment 24: Asthma Guidelines**

Components of Severity		Classification of Asthma Severity												
		Intermittent			Persistent									
					Mild			Moderate			Severe			
		0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs	
<b>Impairment</b>  <small>Normal FEV<sub>1</sub>/FVC : 8-19 yr 85% 20-39 yr 80% 40-59 yr 75% 60-80 yr 70%</small>	Symptoms	≤ 2 days/week			more than 2 days/week but not daily			Daily			Throughout the day			
	Nighttime awakenings	0	≤ 2x/month		1-2x/month	3-4x/month		3-4x/month	> 1x/week but not nightly		> 1x/month	Often 7x/week		
	SABA use for symptom control (not prevention of EIB)	≤ 2 days/week			≤ 2 days/week but not daily			>2 days/week but not daily, and not more than 1x on any day	Daily			Several time per day		
	Interference with normal activity	None			Minor limitation			Some limitation			Extremely limited			
	Lung function	N/A	Normal FEV <sub>1</sub> between exacerbations	Normal FEV <sub>1</sub> between exacerbations	N/A	>80%	>80%	N/A	60-80%	60-80% Reduced	N/A	<60%	<60% Reduced	
FEV <sub>1</sub>		>80%	>80%		>80%	Normal		75-80%	5%		<75%	5%		
FEV <sub>1</sub> /FVC		>85%	Normal											



Components of Severity		Classification of Asthma Severity											
		Intermittent			Persistent								
					Mild			Moderate			Severe		
		0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs
<b>Risk</b>	Exacerbations requiring oral systemic corticosteroids	0-1/year			≥ 2 exacerbations in 6 months requiring oral steroids or >4 wheezing episodes/1-year lasting >1 day and risk factors for persistent asthma	≥2/year Relative annual risk may be related to FEV1.	≥2/year Relative annual risk may be related to FEV1.	≥ 2 exacerbations in 6 months requiring oral steroids or >4 wheezing episodes/1-year lasting >1 day and risk factors for persistent asthma	≥ 2/year Relative annual risk may be related to FEV1.	≥ 2/year Relative annual risk may be related to FEV1.	≥ 2 exacerbations in 6 months requiring oral steroids or >4 wheezing episodes/1-year lasting >1 day and risk factors for persistent asthma	≥2/year Relative annual risk may be related to FEV1.	≥2/year Relative annual risk may be related to FEV1.
		Consider severity and interval since last exacerbation. Frequency and severity may fluctuate over time for subjects in any severity category.											
<b>Recommended Step for Initiating Treatment</b>	Step 1			Step 2			Step 3 and consider short course of oral steroids	Step 3: medium dose ICS and consider short course of oral steroids	Step 3 and consider short course of oral steroids	Step 3 and consider short course of oral steroids	Step 3: medium dose ICS OR Step 4 and consider short course of oral steroids	Step 4 or 5 and consider short course of oral steroids	
	In 2-6 weeks, evaluate level of asthma control that is achieved. _____ 0-4 years: If no clear benefit is observed in 4-6 weeks, stop treatment and consider alternate diagnosis or adjusting therapy. 5-11 and 12+ years: adjust therapy accordingly.												

Components of Control		Classification of Asthma Control								
		Well Controlled			Not Well Controlled			Very Poorly Controlled		
		0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs
	Symptoms	≤ 2 days/week but not more than once on each day		≤ 2 days/week	> 2 days/week or multiple times on ≤ 2 days/week		> 2 days/week	Throughout the day		
<b>Impairment</b>	Nighttime awakenings	≤ 1x/month		≤ 2x/month	> 1x/month	≥ 2x/month	1-3x/week	> 1x/week	≥ 2x/week	≥ 4x/week
	Interference with normal activity	None			Some limitation			Extremely limited		
	SABA use for symptom control (not prevention of EIB)	≤ 2 days/week			> 2 days/week			Several times per day		
	Lung function FEV1 or peak flow FEV1/FVC	N/A	> 80% > 80%	> 80%	N/A	60-80% 75-80%	60-80%	N/A	< 60% < 75%	< 60%
	Validated questionnaires ATAQ ACQ ACT			0 ≤ 0.75 ≥ 20			1-2 ≥ 1.5 16-19			3-4 N/A ≤ 15
<b>Risk</b>	Exacerbations requiring oral systemic corticosteroids	0-1/year			≥ 2/year					
		Consider severity and interval since last exacerbation								
	Reduction in lung growth/ Progressive loss of lung function	Evaluation requires long-term follow-up								
		<ul style="list-style-type: none"> <li>Maintain current step</li> <li>Regular follow-up every 1-6 months</li> </ul>			Step up 1 step	Step up at least 1 step	<ul style="list-style-type: none"> <li>Step up 1 step</li> <li>Reevaluate</li> </ul>	<ul style="list-style-type: none"> <li>Consider short course of oral steroids</li> <li>Step up 1-2 steps</li> </ul>	<ul style="list-style-type: none"> <li>Consider short course of oral steroids</li> </ul>	

Components of Control	Classification of Asthma Control														
	Well Controlled			Not Well Controlled			Very Poorly Controlled								
	0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs	0-4 yrs	5-11 yrs	12 + yrs						
<b>Recommended Action for Treatment</b>	<ul style="list-style-type: none"> <li>Consider step down if well controlled for at least 3 months</li> </ul>			<ul style="list-style-type: none"> <li><b>Before step up:</b> Review adherence to medication, inhaler technique, and environmental control. If alternative treatment was used, discontinue it and use preferred treatment for that step.</li> <li><b>Reevaluate the level of asthma control in 2-6 weeks to achieve control.</b></li> <li>0-4 years: If no clear benefit is observed in 4-6 weeks, consider alternative diagnoses or adjusting therapy.</li> <li>5-11 years: Adjust therapy accordingly.</li> <li><b>For side effects,</b> consider alternative treatment options.</li> </ul>			<ul style="list-style-type: none"> <li>in 2-6 weeks</li> <li>For side effects, consider alternative treatment options</li> </ul>			<ul style="list-style-type: none"> <li><b>Before step up:</b> Review adherence to medication, inhaler technique, and environmental control. If alternative treatment was used, discontinue it and use preferred treatment for that step.</li> <li><b>Reevaluate the level of asthma control in 2-6 weeks to achieve control.</b></li> <li>0-4 years: If no clear benefit is observed in 4-6 weeks, consider alternative diagnoses or adjusting therapy.</li> <li>5-11 years: Adjust therapy accordingly.</li> <li><b>For side effects,</b> consider alternative treatment options.</li> </ul>			<ul style="list-style-type: none"> <li>Step up 1-2 steps</li> <li>Reevaluate in 2 weeks</li> <li>For side effects, consider alternative treatment options</li> </ul>		

**Attachment 25: Antihistamines That May Be Used Predose**

The following antihistamines may be used predose, before Dara-SC injection (including, but not limited to):

- Diphenhydramine
- Cetirizine
- Fexofenadine
- Loratadine
- Clemastine
- Dexchlorpheniramine
- Promethazine\*

\* The IV use of promethazine should be avoided.

**INVESTIGATOR AGREEMENT**

JNJ-68284528 (ciltacabtagene autoleucl)

Clinical Protocol 68284528MMY2003 Amendment 3/USA-1

**INVESTIGATOR AGREEMENT**

I have read this protocol and agree that it contains all necessary details for carrying out this study. I will conduct the study as outlined herein and will complete the study within the time designated.

I will provide copies of the protocol and all pertinent information to all individuals responsible to me who assist in the conduct of this study. I will discuss this material with them to ensure that they are fully informed regarding the study drug, the conduct of the study, and the obligations of confidentiality.

**Coordinating Investigator (where required):**

Name (typed or printed): \_\_\_\_\_

Institution and Address: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

(Day Month Year)

**Principal (Site) Investigator:**

Name (typed or printed): \_\_\_\_\_

Institution and Address: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Telephone Number: \_\_\_\_\_

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

(Day Month Year)

**Sponsor's Responsible Medical Officer:**Name (typed or printed): Sen Zhuang, MD PhD – VP Global Clinical Research & DevelopmentInstitution: Janssen Research & DevelopmentSignature:  \_\_\_\_\_ Date: 16 November 2020

(Day Month Year)

**Note:** If the address or telephone number of the investigator changes during the course of the study, written notification will be provided by the investigator to the sponsor, and a protocol amendment will not be required.

CONFIDENTIAL – FOIA Exemptions Apply in U.S.

261

Approved, Date: 14 November 2020

CONFIDENTIAL – FOIA Exemptions Apply in U.S.

261

Approved, Date: 14 November 2020