

### **CANCELING A TEST**

### **PRINCIPLE**

Follow this procedure to cancel a test that is inadvertently ordered in error, is a duplicate order or is cancelled after dispatching. Canceling with the appropriate comment is how the lab tracks specimens and remains compliant with CAP requirements.

#### **PROCEDURE**

Follow this procedure for any test that is dispatched but not collected or performed; and for tests ordered in duplicate. For example:

- ✓ Patient cannot collect
- ✓ Provider cancels a test after it has been dispatched
- ✓ Tests are dispatched and the member refuses testing
- ✓ An incorrect specimen was drawn and the patient needs to return
- ✓ Specimen integrity was not maintained and the specimen needs to be redrawn
- ✓ Multiple orders for the same test are seen in Collections Inquiry


# To cancel a test that has already been dispatched:

Example: All orders for an Oncology Patient are dispatched and the labels have printed. The patient then indicates that they do not need certain tests today; they are to be collected later.





- Click in the ACCN field and enter the ACCN, press the ENTER key
- 3. A list of tests on the ACCN will be displayed.
- 4. Highlight the test you want to cancel; only one test at a time can be cancelled.



Written: E. Gibson 10/20/10

Revised: E. Gibson 8/11, 11/11, 6/12,

12/12

5. Click on "Task" and then "Cancel Order", the Cancel Order pop up box will display.



6. Select the cancel reason from the list provided when you click on the down arrow button. Click on "Cancel Order."

NOTE: To add additional cancel information as free text, click on the "View" button and choose "Comments." Add additional information. For example: A patient presents in lab with a CBC order. The CBC has a comment "Platelet Clumper." When the CBC is cancelled and re-ordered, the comment must be entered for the re-ordered CBC.

- 7. Always remember to cancel the Auto Diff when canceling a CBC. Because an auto diff is a reflex test, it cannot be canceled using "re-order" codes. Use the code "duplicate" to cancel Auto Diffs.
- 8. Do not use the "Reschedule" option. Rescheduling a test(s) creates a transmission error that prevents the results from going back into HealthConnect and the provider's in-basket.
- 9. Do not order test(s) in DOE. The system will automatically re-order the tests for you.


## To cancel duplicate tests seen in Collections Inquiry:

When there are multiple orders of a test in Collections Inquiry, duplicates can be cancelled. Use caution when canceling tests; only cancel tests for which there are multiple orders i.e. multiple protimes ordered for the previous month or numerous lipid profiles for the same provider. Medicare regulations prohibit reimbursement for multiple tests drawn on the same day. For example - drawing two purple tops, testing and resulting two CBCs. There is potential for this to be viewed as fraud and Kaiser Permanente could be fined for Medicare fraud.

1. Click on the duplicate tests to highlight them

Revised: E. Gibson 8/11, 11/11, 6/12,

Written: E. Gibson 10/20/10



- Dispatch the test as if it were being ordered. Allow the labels to print.
- 3. Go to ORV and follow the procedure outlined above to cancel the orders; use the Duplicate Order cancelation code. **Do not** use any code with "reordered" as this will add the tests back into Millennium.

If it is busy and there is not enough time to cancel right away, it is acceptable to make a note of the patient's MRN and return later (same day) to cancel the duplicate testing. Canceling duplicates in a timely manner makes it easier to order the next time the patient comes to the lab.

### Canceling a STAT Test:

Providers who order STAT testing are expecting results back within a short time frame. Before canceling the test, notify the provider immediately that the STAT cannot be collected and there is a need to cancel the test.

### Canceling a Test That a Patient Refuses:

On occasion, a patient may refuse testing that has been ordered by a provider. It is within a patient's rights to refuse testing for any reason. In order to document that the testing was not collected due to patient refusal and not because of lab error, it is important to perform the following procedure:

- 1. Order the tests that the patient refused. Dispatch the labels.
- 2. Go into ORV and cancel the tests using the cancellation code cancelled, patient refused.
- 3. This comment will go to the patient chart and can be viewed at a later date if there is a question as to why testing was not performed.

Page 3 of 5

Written: E. Gibson 10/20/10

Revised: E. Gibson 8/11, 11/11, 6/12,

4. Place a call to the provider or nursing staff notifying them that the patient has refused testing. Add a test comment stating who was notified of the cancellation, at what time and your name.

## Canceling a CDiff as "Not indicated on formed stool":

When a stool is received for CDiff that is formed and does not meet the criteria for submission, it must be canceled with the appropriate code and the provider notified of the cancellation.

- 1. Dispatch the label for the CDiff.
- 2. Go into ORV and cancel the test using the cancelation code "Not indicated on formed stool." This comment will go to the patient chart.
- 3. Fill out a variance and place a call to the provider or nursing staff notifying them of the canceled test. Add a test comment stating who was notified of the cancelation, at what time and your name.

# Canceling a Quest test that is dispatched for collection:

Quest specimens require special handling when being canceled. If a Quest specimen is dispatched, placed on a transfer List and the list is put In Transit the specimen can be canceled on our end, but a call MUST be placed to Quest informing them of the cancelation. Once the Transfer List was placed In Transit, the specimens hit the Quest Pending Log. If tubes for other Quest tests are received. Quest will perform the canceled test if a call is not received informing them that the test is canceled.

## Canceling a specimen that does not have the correct identifiers:

When a specimen is received that does not have the correct identification it still needs to be tracked. In most cases, a reasonable effort can be made to identify the patient to whom the specimen belongs, but because the correct identification is lacking on the specimen, testing cannot be performed. If the requisition has both identifiers and the specimen is lacking two identifiers, you can proceed with the following process for cancelation.

- 1. Dispatch the test in CI or if the specimen was received with a requisition, order the test in DOE. Do not place the label on the specimen.
- 2. Cancel the specimen (test) using the cancel code "Improperly labeled specimen."

Page 4 of 5

- 3. Place a call to the provider or nursing staff notifying them of the canceled test. Add a test comment stating who was notified of the cancelation, at what time and your name.
  - If the specimen is from the TRiPS Dept fill out a variance and place a call to the Admin Specialists immediately for recollection.

Revised: E. Gibson 8/11, 11/11, 6/12,

Written: E. Gibson 10/20/10

### LIMITATIONS OF THE PROCEDURE

If the cancelled tests do not need to be re-ordered, do not choose a code that will automatically re-order. Always choose a code that reflects the reason why the tests are being cancelled. Providers have called to ask for reasons for canceled tests. See the list below for available codes.

### **CANCELLATION CODES THAT RE-ORDER TESTING**

Dispatched in error and re-ordered

Reordered, broken/spilled in transit

Reordered, clotted

Reordered, hemolyzed specimen

Reordered, incorrect test accessioned

Reordered, insufficient quantity

Reordered, no specimen received Reordered, patient not fasting

Reordered, physician request

Reordered, physician request

Reordered, test not due at this time

Reordered, sample not collected

Reordered, specimen contaminated

Reordered, specimen too viscous

Reordered, specimen unacceptable

Reordered, specimen unacceptable Reordered Reo

Reordered, specimen lost in transit

Reordered, specimen stability exceed

Reordered, wrong patient

## Miscellaneous codes for testing that does NOT need to be re-ordered

Cancelled by pathologist

Changed order

Changed from STAT to routine

Double labeled

GSM referral, Result viewable in chart

HCV titer below minimum level

Inconclusive, referred for sequencing

Not indicated

Not indicated on watery stool

Other

Referred to Children's for venipuncture

Specimen coll for elevated hct

Specimen QNS for required repeat

Test not performed

Test referred, see separate report

Quantity not sufficient

Cancelled after collected

Changed from routine to STAT

Confirmed by another method

Duplicate order

HCV not detected

Improperly labeled specimen

No source provided

Not indicated on formed stool

Nurse request

Patient declined due to cost of procedure

Referred to outside lab

Spec ID unclear

Test contraindicated

Test referred, result viewable in chart

Wrong encounter

Written: E. Gibson 10/20/10 Revised: E. Gibson 8/11, 11/11, 6/12,

12/12